


Newton's Law of Motion (II)

Physics 211
Syracuse University, Physics 211 Spring 2015
Walter Freeman


February 10, 2015

- Homework 3 due tomorrow


Exam 1 statistics



Exam 1 statistics




Exam 1 statistics



This exam was harder than earlier ones

Exam 1 statistics




Your average was 71%

This exam was harder than earlier ones

The historical average was 60%

Exam 1 statistics




This exam was harder than earlier ones

The historical average was 60%

Historically classes have had an extra week of prep!

Exam 1 statistics



This exam was harder than earlier ones

The historical average was 60%

Historically classes have had an extra week of prep!

I am very proud of you all!

Ask a Physicist: Water wheels and power stations

- Gravity

- Every object near Earth feels a force with magnitude mg directed downward. No exceptions!

- Normal forces

- Always directed normal (perpendicular) to a surface
- Magnitude is as large as it needs to be to stop objects from “crossing” ($a_{\perp} = 0$)
- Newton’s third law: if A pushes on B, B pushes back on A (the book problem)

- Tension

- The force transmitted through a rope from one thing to another
- Same on both sides of the rope (Newton’s 3rd...)

Force diagrams

- Accounting devices for your use, to keep straight forces for $\vec{F} = m\vec{a}$
- Some guidelines:
 - Draw a **separate diagram** for each object (book problem again!)
 - Each force gets a separate arrow
 - Draw them **big enough** that you can draw “component-triangles”
 - “Net force”, velocity, acceleration not forces; only physical agents are

(Examples on document camera)

Force is a vector; handle it like any other

One copy of Newton's second law in each direction (per object)

$$\vec{F} = m\vec{a} \rightarrow \begin{pmatrix} F_x = ma_x \\ F_y = ma_y \end{pmatrix}$$

Force is a vector; handle it like any other

One copy of Newton's second law in each direction (per object)

$$\vec{F} = m\vec{a} \rightarrow \begin{pmatrix} F_x = ma_x \\ F_y = ma_y \end{pmatrix}$$

Important: When dealing with inclines, choose your axes to align with the incline! (F_N is easy that way)

A problem-solving recipe (remember this!)

- **Accounting:** Draw force diagrams for every object
 - Work out components (trigonometry) of vectors in funny directions – no need for numbers

A problem-solving recipe (remember this!)

- **Accounting:** Draw force diagrams for every object
 - Work out components (trigonometry) of vectors in funny directions – no need for numbers
- **Physics:** Write down $\sum F = ma$ in each dimension, for each object

A problem-solving recipe (remember this!)

- **Accounting:** Draw force diagrams for every object
 - Work out components (trigonometry) of vectors in funny directions – no need for numbers
- **Physics:** Write down $\sum F = ma$ in each dimension, for each object
- **Math:** Put in the stuff you know, solve for the stuff you don't

A problem-solving recipe (remember this!)

- **Accounting:** Draw force diagrams for every object
 - Work out components (trigonometry) of vectors in funny directions – no need for numbers
- **Physics:** Write down $\sum F = ma$ in each dimension, for each object
- **Math:** Put in the stuff you know, solve for the stuff you don't
- **Kinematics:** Connect acceleration to motion

A problem-solving recipe (remember this!)

- **Accounting:** Draw force diagrams for every object
 - Work out components (trigonometry) of vectors in funny directions – no need for numbers
- **Physics:** Write down $\sum F = ma$ in each dimension, for each object
- **Math:** Put in the stuff you know, solve for the stuff you don't
- **Kinematics:** Connect acceleration to motion

It really is this easy; I promise!

A problem-solving recipe (remember this!)

- **Accounting:** Draw force diagrams for every object
 - Work out components (trigonometry) of vectors in funny directions – no need for numbers
- **Physics:** Write down $\sum F = ma$ in each dimension, for each object
- **Math:** Put in the stuff you know, solve for the stuff you don't
- **Kinematics:** Connect acceleration to motion

It really is this easy; I promise!
“Ask physics the question, don't tell it the answer”

Sample questions

A stone hangs from the roof of a car by a string; the car accelerates forward at 3 m/s^2 .

- What happens to the string?

A stone hangs from the roof of a car by a string; the car accelerates forward at 3 m/s^2 .

- What happens to the string?
- What angle does the string make with the vertical?

A stone hangs from the roof of a car by a string; the car accelerates forward at 3 m/s^2 .

- What happens to the string?
- What angle does the string make with the vertical?
- What is the tension in the string?

Sample questions

A cart slides down a frictionless track elevated at angle θ ; what is its acceleration?

- What happens to the string?

A cart slides down a frictionless track elevated at angle θ ; what is its acceleration?

- What happens to the string?
- What angle does the string make with the vertical?

A cart slides down a frictionless track elevated at angle θ ; what is its acceleration?

- What happens to the string?
- What angle does the string make with the vertical?
- What is the tension in the string?

Sample questions

Two masses of 20 and 40 kg hang from a massless pulley on either side. How do they move?

Sample questions

Two masses of m_1 and m_2 kg hang from a massless pulley on either side. How do they move?