

# 1D kinematics: solving problems

Physics 211  
Syracuse University, Physics 211 Spring 2022  
Walter Freeman

January 23, 2023

# On solving problems

You can recognize truth by its beauty and simplicity. When you get it right, it is obvious that it is right—at least if you have any experience—because usually what happens is that more comes out than goes in.... Inexperienced students make guesses that are very complicated, [but] the truth always turns out to be simpler than you thought.

—Richard Feynman, quoted by K. C. Cole, in *Sympathetic Vibrations: Reflections on Physics as a Way of Life* (1985)

Nature uses only the longest threads to weave her patterns, so each small piece of her fabric reveals the organization of the entire tapestry.

—Richard Feynman, *The Character of Physical Law* (1965)

- Homework 1 due Friday; homework 2 assigned tomorrow, due next Friday
- If at all possible, finish HW1 early and start HW2 this week
- I'll be in the Physics Clinic to help people with anything:
  - 2-4 PM today (Tuesday)
  - 2:15-4 PM Wednesday
  - 1:30-3:00 PM Thursday
- You can also ask your classmates and us for advice on Discord

## “Ask a Physicist”

“Why does gravity make all objects fall at the same rate, in light of Newton’s second law?”

## “Ask a Physicist”

“Why does gravity make all objects fall at the same rate, in light of Newton’s second law?”

Newton’s second law: **“It takes more force to make more massive things accelerate”**

## “Ask a Physicist”

“Why does gravity make all objects fall at the same rate, in light of Newton’s second law?”

Newton’s second law: **“It takes more force to make more massive things accelerate”**

We should look at this backwards: **We observe that all objects fall at the same rate. What must the law of gravity be to make this happen?**

## “Ask a Physicist”

“Why does gravity make all objects fall at the same rate, in light of Newton’s second law?”

Newton’s second law: **“It takes more force to make more massive things accelerate”**

We should look at this backwards: **We observe that all objects fall at the same rate. What must the law of gravity be to make this happen?**


The law of gravity: **“Gravity applies more force to more massive things”**

- Review material from last time
- Position, velocity, and acceleration graphs
- Problem-solving method for kinematics problems
- Sample problems




- Review material from last time
- Position, velocity, and acceleration graphs
- Problem-solving method for kinematics problems
- Sample problems
- Homework help?

# Last time: Position, velocity, and acceleration



# Last time: Position, velocity, and acceleration



- If we know acceleration as a function of time, how do we get from there to position vs. time?

- If we know acceleration as a function of time, how do we get from there to position vs. time?
- A. Look at the slope of the acceleration vs. time graph to get velocity, and then look at its slope to get position
- B. Look at the area under the curve of the acceleration vs. time graph to get velocity, and then look at the area under that graph to get position
- C. Take two derivatives of the acceleration vs. time graph to get position vs. time
- D. Take two integrals of the acceleration vs. time graph to get position vs. time

# The “kinematics equations”

$$v(t) = at + v_0$$

$$x(t) = \frac{1}{2}at^2 + v_0t + x_0$$

These equations are valid when...

- A. Acceleration is constant
- B. Velocity is constant
- C. The object moves in only one direction
- D. They are always valid

# The “kinematics equations”

$$v(t) = at + v_0$$

$$x(t) = \frac{1}{2}at^2 + v_0t + x_0$$

These equations are valid when...

- A. Acceleration is constant
  - B. Velocity is constant
  - C. The object moves in only one direction
  - D. They are always valid
- 
- A: these are the expressions for  $x(t)$  and  $v(t)$  when acceleration is constant!

How long does it take an object to fall from a height  $h$ ?



## Example problems

How long does it take an object to fall from a height  $h$ ?

First: Write the position and velocity equations, substituting in things you know.  
(Here, take ground level to be  $y = 0$ , and upward to be the positive direction.)

$$\text{A: } x(t) = \frac{1}{2}gt^2 + h$$

$$v(t) = -gt$$

$$\text{B: } x(t) = -\frac{1}{2}gt^2 + v_0t + h$$

$$v(t) = -gt$$

$$\text{C: } x(t) = -\frac{1}{2}gt^2 + h$$

$$v(t) = gt$$

$$\text{D: } x(t) = -\frac{1}{2}gt^2 + h$$

$$v(t) = -gt$$

$$\text{E: } x(t) = -\frac{1}{2}gt^2 + v_0t$$

$$v(t) = -gt$$

## Example problems

How long does it take an object to fall from a height  $h$ ?

## Example problems

How long does it take an object to fall from a height  $h$ ?

Second: Phrase the question in terms of your algebraic variables.

From the previous:  $x(t) = -\frac{1}{2}gt^2 + h$  and  $v(t) = -gt$ .

(Again, take ground level to be  $y = 0$ , and upward to be the positive direction.)

A: “What is the value of  $t$  when  $v = 0$ ?”

B: “What is the value of  $x$  when  $t = 0$ ?”

C: “What is the value of  $x$  when  $v = 0$ ?”

D: “What is the value of  $t$  when  $x = h$ ?”

E: “What is the value of  $t$  when  $x = 0$ ?”

## Example problems

How long does it take an object to fall from a height  $h$ ?

Third: Do the algebra your sentence tells you to do: “What is the value of  $t$  when  $x = 0$ ?”

From the previous:  $x(t) = -\frac{1}{2}gt^2 + h$  and  $v(t) = -gt$ .

(Again, take ground level to be  $y = 0$ , and upward to be the positive direction.)

A:  $\sqrt{2g/h}$

B:  $h/g$

C:  $\sqrt{2h/g}$

D:  $2h/g$

E:  $\sqrt{h/g}$

## Another example

You throw an object up with an initial speed of  $v_0$ . How much time does it take to reach a height  $h$ ?

## Another example

You throw an object up with an initial speed of  $v_0$ . How much time does it take to reach a height  $h$ ?

$$\begin{aligned}x(t) &= \frac{1}{2}at^2 + v_0t + x_0 \\h &= -\frac{1}{2}gt^2 + v_0t \\0 &= -\frac{1}{2}gt^2 + v_0t - h\end{aligned}$$

- $\rightarrow$  You need the quadratic formula for this – nonzero  $a$ ,  $v_0$ , and position
- The quadratic formula gives you two answers, but there's clearly only one
- In this case, both roots are positive. Do you want (A) the larger one, or (B) the smaller one?

## Another example

You throw an object up with an initial speed of  $v_0$ . How much time does it take to reach a height  $h$ ?

$$\begin{aligned}x(t) &= \frac{1}{2}at^2 + v_0t + x_0 \\h &= -\frac{1}{2}gt^2 + v_0t \\0 &= -\frac{1}{2}gt^2 + v_0t - h\end{aligned}$$

- $\rightarrow$  You need the quadratic formula for this – nonzero  $a$ ,  $v_0$ , and position
- The quadratic formula gives you two answers, but there's clearly only one
- In this case, both roots are positive. Do you want (A) the larger one, or (B) the smaller one?
- The homework asks you to address this idea.
- Hint: graph position vs. time, and interpret the question graphically
- What is the *mathematical* interpretation of the quadratic formula?

## Example problems

I am standing at the bottom of a hole of depth  $h$ . Someone throws a ball down to me at speed  $v_0$ . How fast is it going when it reaches me?

First: Write equations for  $x(t)$  and  $v(t)$ , putting in the things you know. (Here, take ground level as zero, and downward to be positive.)

$$\text{A: } x(t) = \frac{1}{2}gt^2$$

$$v(t) = -gt$$

$$\text{B: } x(t) = -\frac{1}{2}gt^2 + v_0t + h$$

$$v(t) = -gt$$

$$\text{C: } x(t) = \frac{1}{2}gt^2 + v_0t$$

$$v(t) = gt + v_0$$

$$\text{D: } x(t) = \frac{1}{2}gt^2 - v_0t$$

$$v(t) = -gt$$

$$\text{E: } x(t) = -\frac{1}{2}gt^2 - v_0t$$

$$v(t) = -gt$$



## Example problems

I am standing at the bottom of a hole of depth  $h$ . Someone throws a ball down to me at speed  $v_0$ . How fast is it going when it reaches me?

Second: Ask a question in terms of your algebraic variables. (Here, take ground level as zero, and downward to be positive.)

A: “What is  $v$  at the time when  $x$  is  $h$ ?”

B: “What is  $v$  at the time when  $x$  is  $0$ ?”

C: “What is  $t$  at the time when  $x$  is  $h$ ?”

D: “What is  $v$  at the time when  $x$  is  $-h$ ?”

E: “What is  $x$  at the time when  $v$  is  $0$ ?”

## Example problems

I am standing at the bottom of a hole of depth  $h$ . Someone throws a ball down to me at speed  $v_0$ . How fast is it going when it reaches me?

Third: Do the algebra. I'll demonstrate this on the document camera. This requires two steps: first find the time, then find  $v$ .

# Homework/recitation questions?

## Example problems

Some students get on the elevator in Bray Hall and press the button to go up. The elevator accelerates upward at  $0.5 \text{ m/s}^2$  for four seconds before it breaks down.

Thankfully, the emergency brake engages; the emergency brake decelerates the elevator at  $3 \text{ m/s}^2$  until it comes to a stop.

## Example problems

Some students get on the elevator in Bray Hall and press the button to go up. The elevator accelerates upward at  $0.5 \text{ m/s}^2$  for four seconds before it breaks down.

Thankfully, the emergency brake engages; the emergency brake decelerates the elevator at  $3 \text{ m/s}^2$  until it comes to a stop.

One of the students has a crowbar and pries the doors open, hoping to crawl out. How far above ground level is the elevator stuck?

- Can we use  $x(t) = \frac{1}{2}at^2 + v_0t + x_0$  for this? Why or why not?

## Example problems

Some students get on the elevator in Bray Hall and press the button to go up. The elevator accelerates upward at  $0.5 \text{ m/s}^2$  for four seconds before it breaks down.

Thankfully, the emergency brake engages; the emergency brake decelerates the elevator at  $3 \text{ m/s}^2$  until it comes to a stop.

One of the students has a crowbar and pries the doors open, hoping to crawl out. How far above ground level is the elevator stuck?

- Can we use  $x(t) = \frac{1}{2}at^2 + v_0t + x_0$  for this? Why or why not?
  - The acceleration isn't constant, but it's **piecewise constant**

## Example problems

Some students get on the elevator in Bray Hall and press the button to go up. The elevator accelerates upward at  $0.5 \text{ m/s}^2$  for four seconds before it breaks down.

Thankfully, the emergency brake engages; the emergency brake decelerates the elevator at  $3 \text{ m/s}^2$  until it comes to a stop.

One of the students has a crowbar and pries the doors open, hoping to crawl out. How far above ground level is the elevator stuck?

- Can we use  $x(t) = \frac{1}{2}at^2 + v_0t + x_0$  for this? Why or why not?
  - The acceleration isn't constant, but it's **piecewise constant**
  - Use one copy of the constant-acceleration relations for the first phase and another copy for the second phase

## Example problems

Some students get on the elevator in Bray Hall and press the button to go up. The elevator accelerates upward at  $0.5 \text{ m/s}^2$  for four seconds before it breaks down.

Thankfully, the emergency brake engages; the emergency brake decelerates the elevator at  $3 \text{ m/s}^2$  until it comes to a stop.

One of the students has a crowbar and pries the doors open, hoping to crawl out. How far above ground level is the elevator stuck?

- Can we use  $x(t) = \frac{1}{2}at^2 + v_0t + x_0$  for this? Why or why not?
  - The acceleration isn't constant, but it's **piecewise constant**
  - Use one copy of the constant-acceleration relations for the first phase and another copy for the second phase
  - Find the position and velocity at the end of the first phase
  - Those are the position and velocity at the beginning of the first phase



## Example problems

Some students get on the elevator in Bray Hall and press the button to go up. The elevator accelerates upward at  $0.5 \text{ m/s}^2$  for four seconds before it breaks down.

Thankfully, the emergency brake engages; the emergency brake decelerates the elevator at  $3 \text{ m/s}^2$  until it comes to a stop.

One of the students has a crowbar and pries the doors open, hoping to crawl out. How far above ground level is the elevator stuck?

- Can we use  $x(t) = \frac{1}{2}at^2 + v_0t + x_0$  for this? Why or why not?
  - The acceleration isn't constant, but it's **piecewise constant**
  - Use one copy of the constant-acceleration relations for the first phase and another copy for the second phase
  - Find the position and velocity at the end of the first phase
  - Those are the position and velocity at the beginning of the first phase

How long will it take before the administration sends an email out about the elevator?

## Example problems

- A bucket is being lowered from a cliff at a rate of  $10 \text{ m/s}$ . You drop a rock off the cliff when the bucket is  $10 \text{ m}$  beneath the top. How long does it take for the rock to land in the bucket?

Same idea as before; see example on the document camera.