

# Momentum

Physics 211  
Syracuse University, Physics 211 Spring 2022  
Walter Freeman

April 7, 2022

# Announcements

- Group Exam 3 tomorrow; Exam 3 next Tuesday
- Review notes posted on the website
- Extra homework help hours today: 9:45-10:45
- HW7 due Friday

# Extra exam review session

Extra exam review session will be held **Monday afternoon or evening**. We are still getting a time/place. (I am sadly booked all weekend.)

This will be a little more scripted than the last one, but I expect it should prepare everyone for the exam – instead of “cramming” on your own, come work with us.

To get the maximum benefit out of this, you should:

- Have done all your homework
- Go to the Physics Clinic after the group exam Friday and ask the coaches to help you with anything you couldn't do
- Be familiar with most of the recitation exercises we have done
- Have already attempted the practice exam over the weekend, and have looked at the solutions (posted Friday or Saturday)

# Recitation or homework questions?

# Recitation or homework questions?

Shall we talk about the “one with the truck”?

# Exam 3 Review

This unit is about *conservation laws*.

We have met two of them: *conservation of momentum* and the *work-energy theorem*.

These techniques let you analyze systems where you know something about “before” and “after” states.

General problem-solving techniques:

- **Draw cartoons** for critical parts of the motion (“before”, “after”, and “in between”)

# Exam 3 Review

This unit is about *conservation laws*.

We have met two of them: *conservation of momentum* and the *work-energy theorem*.

These techniques let you analyze systems where you know something about “before” and “after” states.

General problem-solving techniques:

- **Draw cartoons** for critical parts of the motion (“before”, “after”, and “in between”)
- **Choose a technique** to relate the things in each cartoon to the next

# Exam 3 Review

This unit is about *conservation laws*.

We have met two of them: *conservation of momentum* and the *work-energy theorem*.

These techniques let you analyze systems where you know something about “before” and “after” states.

General problem-solving techniques:

- **Draw cartoons** for critical parts of the motion (“before”, “after”, and “in between”)
- **Choose a technique** to relate the things in each cartoon to the next
- **Think carefully** about the terms (momentum, energy, work, etc.) that apply to each technique



# Exam 3 Review

This unit is about *conservation laws*.

We have met two of them: *conservation of momentum* and the *work-energy theorem*.

These techniques let you analyze systems where you know something about “before” and “after” states.

General problem-solving techniques:

- **Draw cartoons** for critical parts of the motion (“before”, “after”, and “in between”)
- **Choose a technique** to relate the things in each cartoon to the next
- **Think carefully** about the terms (momentum, energy, work, etc.) that apply to each technique
- **Write down the equations** for the work-energy theorem or conservation of momentum and do math

# Exam 3 Review

This unit is about *conservation laws*.

We have met two of them: *conservation of momentum* and the *work-energy theorem*.

These techniques let you analyze systems where you know something about “before” and “after” states.

General problem-solving techniques:

- **Draw cartoons** for critical parts of the motion (“before”, “after”, and “in between”)
- **Choose a technique** to relate the things in each cartoon to the next
- **Think carefully** about the terms (momentum, energy, work, etc.) that apply to each technique
- **Write down the equations** for the work-energy theorem or conservation of momentum and do math

# Review: energy methods

Use energy methods when:

- You have clear before and after states (draw your cartoons, dammit!)
- You can calculate the work done by the forces between them (**not collisions/explosions!**)
- You don't care about time

# Review: energy methods

Use energy methods when:

- You have clear before and after states (draw your cartoons, dammit!)
- You can calculate the work done by the forces between them (**not collisions/explosions!**)
- You don't care about time
- *Be careful with projectile motion!*
  - Energy methods can tell you “how fast” or “how high”
  - They cannot tell you “where does it land?”

## Review: The work-energy theorem

Work-energy theorem:  $\frac{1}{2}mv_f^2 - \frac{1}{2}mv_i^2 = \vec{F} \cdot \vec{d} = Fd \cos \theta$  (if this is constant)

Potential energy is an alternate way of keeping track of the work done by conservative forces:

- $PE_{\text{grav}} = mgh$
- $PE_{\text{spring}} = \frac{1}{2}kx^2$

# Review: Conservation of energy

$$PE_i + \frac{1}{2}mv_i^2 + W_{\text{other}} = PE_f + \frac{1}{2}mv_f^2$$

# Review: Conservation of energy

$$\begin{array}{ccccccccc} PE_i & + & \frac{1}{2}mv_i^2 & + & W_{\text{other}} & = & PE_f & + & \frac{1}{2}mv_f^2 \\ \text{(initial PE)} & + & \text{(initial KE)} & + & \text{(other work)} & = & \text{(final PE)} & + & \text{(final KE)} \end{array}$$

# Review: Conservation of energy

$$\begin{aligned} PE_i + \frac{1}{2}mv_i^2 + W_{\text{other}} &= PE_f + \frac{1}{2}mv_f^2 \\ \text{(initial PE)} + \text{(initial KE)} + \text{(other work)} &= \text{(final PE)} + \text{(final KE)} \\ \text{(total initial mechanical energy)} + \text{(other work)} &= \text{(total final mechanical energy)} \end{aligned}$$



## Review: Conservation of energy

$$\begin{aligned} PE_i + \frac{1}{2}mv_i^2 + W_{\text{other}} &= PE_f + \frac{1}{2}mv_f^2 \\ \text{(initial PE)} + \text{(initial KE)} + \text{(other work)} &= \text{(final PE)} + \text{(final KE)} \\ \text{(total initial mechanical energy)} + \text{(other work)} &= \text{(total final mechanical energy)} \end{aligned}$$

Since conservation of energy is the broadest principle in science, it's no surprise that we can do this!

# Review: Conservation of momentum

Use conservation of momentum when:

# Review: Conservation of momentum

Use conservation of momentum when:

- You have a collision (two objects collide and exchange forces for a brief time)
- You have an explosion (one object separates into two)
- Two objects exchange forces *only with each other*

# Review: Conservation of momentum

Use conservation of momentum when:

- You have a collision (two objects collide and exchange forces for a brief time)
- You have an explosion (one object separates into two)
- Two objects exchange forces *only with each other*

(that's it – the last one doesn't come up much, but sometimes it does: why does the Earth wobble?)

# Review: Conservation of momentum

To use conservation of momentum:

- Make sure you have your clear before/after cartoons
- $\sum \vec{p}_i = \sum \vec{p}_f$

# Review: Conservation of momentum

To use conservation of momentum:

- Make sure you have your clear before/after cartoons
- $\sum \vec{p}_i = \sum \vec{p}_f$
- **Momentum is a vector!**

# Review: Conservation of momentum

To use conservation of momentum:

- Make sure you have your clear before/after cartoons
- $\sum \vec{p}_i = \sum \vec{p}_f$
- **Momentum is a vector!**
- If you have motion in two dimensions, momentum in each direction is conserved separately:

$$\sum p_{x_i} = \sum p_{x_f}$$

$$\sum p_{y_i} = \sum p_{y_f}$$

Do not get lazy with your subscripts!

## Sample problems: an excited dog

A person of mass  $m$  is sitting in a tire swing with a string of length  $L$  when their dog (mass  $M$ ) runs and jumps horizontally into their lap.

If they swing up to an angle  $\theta$  above the horizontal, how fast was their dog running?



# What about “bouncing”?

Just knowing that momentum is conserved doesn't fully tell you what happens after a collision.

You also need to know how “sticky” or “bouncy” the objects are.

# What about “bouncing”?

Just knowing that momentum is conserved doesn't fully tell you what happens after a collision.

You also need to know how “sticky” or “bouncy” the objects are.

One way to describe this: *how does the total kinetic energy change?*

- Objects stick together: most kinetic energy lost (“inelastic collision”)

# What about “bouncing”?

Just knowing that momentum is conserved doesn't fully tell you what happens after a collision.

You also need to know how “sticky” or “bouncy” the objects are.

One way to describe this: *how does the total kinetic energy change?*

- Objects stick together: most kinetic energy lost (“inelastic collision”)
- Objects bounce a bit: some kinetic energy lost (“partially inelastic”)

# What about “bouncing”?

Just knowing that momentum is conserved doesn't fully tell you what happens after a collision.

You also need to know how “sticky” or “bouncy” the objects are.

One way to describe this: *how does the total kinetic energy change?*

- Objects stick together: most kinetic energy lost (“inelastic collision”)
- Objects bounce a bit: some kinetic energy lost (“partially inelastic”)
- Objects bounce “fully”: no kinetic energy lost (“elastic collision”)

# What about “bouncing”?

Just knowing that momentum is conserved doesn't fully tell you what happens after a collision.

You also need to know how “sticky” or “bouncy” the objects are.

One way to describe this: *how does the total kinetic energy change?*

- Objects stick together: most kinetic energy lost (“inelastic collision”)
- Objects bounce a bit: some kinetic energy lost (“partially inelastic”)
- Objects bounce “fully”: no kinetic energy lost (“elastic collision”)
- Kinetic energy *increases*: only possible with some energy source!

# What about “bouncing”?

Just knowing that momentum is conserved doesn't fully tell you what happens after a collision.

You also need to know how “sticky” or “bouncy” the objects are.

One way to describe this: *how does the total kinetic energy change?*

- Objects stick together: most kinetic energy lost (“inelastic collision”)
- Objects bounce a bit: some kinetic energy lost (“partially inelastic”)
- Objects bounce “fully”: no kinetic energy lost (“elastic collision”)
- Kinetic energy *increases*: only possible with some energy source!

# An application: neutron moderators

The only truly elastic collisions in nature are between particles. If we want *totally* elastic collisions, we should look to nuclear physics!

A note: this calculation we are going to do here demonstrates two things:

- How elastic collisions work
- ... and how *the art of approximation* is used in physics and engineering!

Recall how a nuclear reactor works:

- $^{235}\text{U}$  fissions when struck by neutrons with low energy (600 times more likely at low energy, less than 0.1 eV)
- When  $^{235}\text{U}$  fissions, it produces neutrons with 2 MeV of kinetic energy ( $v \approx 20$  million m/s)

How do we make these neutrons go from 2 million eV to 0.1 eV of kinetic energy so they can produce more fissions?

# Bounce it off of other atoms!

Compared to a neutron moving at  $2 \times 10^6$  m/s, other atoms appear to be holding still!

This leads us directly to something we know how to do:



# Bounce it off of other atoms!

Compared to a neutron moving at  $2 \times 10^6$  m/s, other atoms appear to be holding still!

This leads us directly to something we know how to do:

A neutron of mass  $m$  and kinetic energy  $E_0$  strikes another atom of mass  $M$ . They collide elastically. What is the neutron's energy after the collision?

# Bounce it off of other atoms!

Compared to a neutron moving at  $2 \times 10^6$  m/s, other atoms appear to be holding still!

This leads us directly to something we know how to do:

A neutron of mass  $m$  and kinetic energy  $E_0$  strikes another atom of mass  $M$ . They collide elastically. What is the neutron's energy after the collision?

$$v_f = -v_0 \frac{M - m}{m + M}$$

# Bounce it off of other atoms!

Compared to a neutron moving at  $2 \times 10^6$  m/s, other atoms appear to be holding still!

This leads us directly to something we know how to do:

A neutron of mass  $m$  and kinetic energy  $E_0$  strikes another atom of mass  $M$ . They collide elastically. What is the neutron's energy after the collision?

$$v_f = -v_0 \frac{M - m}{m + M}$$

How much kinetic energy is lost?

# Bounce it off of other atoms!

Compared to a neutron moving at  $2 \times 10^6$  m/s, other atoms appear to be holding still!

This leads us directly to something we know how to do:

A neutron of mass  $m$  and kinetic energy  $E_0$  strikes another atom of mass  $M$ . They collide elastically. What is the neutron's energy after the collision?

$$v_f = -v_0 \frac{M - m}{m + M}$$

How much kinetic energy is lost?

The fraction of kinetic energy lost is

$$4 \frac{m}{M}.$$

# So what atoms can we use as moderators?

They have to scatter neutrons more readily than they absorb them (hydrogen so-so, oxygen/carbon/heavy hydrogen great)

They have to be lightweight (that's what we just found)

They have to not be chemically grouchy (no hydrogen or oxygen by themselves!)

- $\text{H}_2\text{O}$  (light water: most of the world, not the best moderator)
- $\text{D}_2\text{O}$  (heavy water: Canadians)
- $\text{CO}_2$  (carbon dioxide: British)
- Pure carbon (Graphite: Soviets)