

PHYSICS 307 PROJECT 3

Due Tuesday, 3 October, at 11 AM

1. First we will study a DE that we can solve analytically so we can compute error as a function of stepsize.

Newton's law of cooling says that the rate of temperature loss is proportional to the temperature difference between the cooling object and the ambient temperature T_a ; that is,

$$\frac{\partial T}{\partial t} = -k(T - T_a) \quad (1)$$

where k is a proportionality constant that depends on the materials involved and their surface area. The analytic solution to this DE is

$$T(t) = T_a + (T(0) - T_a)e^{-kt} \quad (2)$$

Consider a cooling cup of tea. When the water is removed from the heat, it is at 100° . Suppose that room temperature is 30° , and that $k = 0.1\text{min}^{-1}$.

- (a) Write a program that solves the differential equation above for the temperature T as a function of time using Euler's method. Plot this function vs. time for stepsizes dt of 1 minute, 10 seconds, and 1 second, along with the analytic solution. Does your numeric solution do a good job of capturing the behavior of the system?
 - (b) Now, investigate the error in a rigorous way. Using a range of stepsizes dt , compute the temperature after five minutes and compute the error in each value. Make a log-log plot of the error vs. the stepsize. Is the scaling what you expect it to be? Note: You should actually solve the DE numerically using Euler's method; the exact solution given above is just so you can compute the error. This is the second-to-last thing we will study in this class that has an easy, obvious analytic solution!
2. Repeat the preceding problem with the second-order Runge-Kutta method.
 3. Now, using RK2, answer the question "How long does it take the tea to cool to $T_f = 70^\circ$?" Make a plot of the error in your result vs. stepsize and verify that you're seeing second-order scaling.

Note that in order to achieve second-order accuracy, you will need to correct for "overshoot", as we will discuss in class.

Briefly, if you discover that the temperature is T_1 (greater than 70 degrees) at t_1 , and T_2 (less than 70 degrees) at t_2 , in order to achieve second-order precision you need to

interpolate to figure out at what time you actually hit 70 degrees. That time is given by

$$t_1 + \frac{t_1 - t_2}{T_1 - T_2}(T_1 - T_f) \tag{3}$$

or alternatively

$$t_1 + \left(\frac{\partial T}{\partial t} \Big|_{t_1} \right)^{-1} (T_1 - T_f) \tag{4}$$

(I find the second more intuitive and easier to apply, since you already have been calculating $\frac{\partial T}{\partial t}$ many times already.)