Analysis of Phase Jump Occurrence in IGS Station Data

Zijun Liu and Brian Breitsch
Email: zili5742@colorado.edu

Introduction

I had a chance to work with graduate mentor for research project at the spring semester of 2019. It was such a great opportunity to practice with solving practical problems with everything that I have learned. After discussed with graduate mentor, we decided to focus on what factors that casuse TEC line jump points. The reason that I chose this topic, because it's basic for aerospace area and it can be used for Machine Learning to collect the data and analyze why it causes these jump points. About what is TEC is Total Electron Context. Why we want to do this, because the global positional system is a wide applied tool and it investigates the earth's ionosphere. That's why we want to do this.

As for the *Figure 1*, this is the TEC line jump point picture from Sep 1st, 2018 in G02 satellite. As for x-axis is the Time, y-axis is the TEC. As seen from the Figure, the orange scattered point is the TEC line jump point at the end of the the line.



Figure 1

Research Objectives

My research topic is Analysis of Phase Jump Occurrence in IGS Station Data. I'm going to focus on factors about why jump point occurred. As seen from Figure 1, I used the difference between Pseudorange for dual frequency.

Equations

Method

For this research, there are four subsections for this research.
First, I called it as **Plot Jump Point**.
The first part is about plot the TEC line and check what is be the reasons cause it. I use the difference between dual frequency signal from satellite receiver and use the equation above to plot it. Here is the example of the plot. The left one is satellite G01 from Sep 3rd, 2018 (*see Figure 2*). The right one satellite G05 from Sep 3rd, 2018 (*see Figure 3*).



Figure 2



Figure 3

The second part is about **auto download the data and plot the data**. In this research project, I have to check many data for the similarity. For example, I go through the data from satellite G02 about 1 or 2 year's data. it will take long if I plot each data manually, so that's why we work on how to auto download the data from the website to get the observation data and how to auto plot it. After that, I will automatically save it in my laptop as picture. And then, I will check for difference and similarity.

The next step is that I have to check how many jump point occurring at each satellite in whole year. I called this step as **count for How many Jump point**. The same problem is that I have to check the data as much as possible. It could automatically do the calculation on what I will work on it. I use threshold to compare with each data and then export it as .csv file. It's very important because we have to use these data to find out the Azimuth and elevation to make sky plot. Here is the example about how the table in .csv file like (*see Figure 4*).

A	B
Name_of_Satellite_Statio	Number_of_Jump_Point
20170821_G10	7
20170821_G13	10
20170821_G15	7
20170821_G16	6
20170821_G18	5
20170821_G20	2
20170821_G21	1
20170821_G24	10
20170821_G27	11
20170821_G29	2
20170821_G32	1
20170821_G08	8
20170821_G14	15
20170821_G11	18
20170821_G31	2
20170821_G01	15
20170821_G25	10
20170821_G22	3
20170821_G12	3
20170821_G03	8
20170821_G26	5
20170821_G23	0
20170821_G09	5

(*see Figure 4*)

In the third step, I'm going to use Azimuth and Elevation to get the sky Plot. I called it as **Sky Plot by Azimuth and Elevation**. I'm going to get the Azimuth and Elevation for each station and export it in the .csv file too (see Figure 5).

A	B	C	D	E
date_str	station_id	satellite_id	azimuth	elevation
20170801	nist	G01	-82.11344601	40.31870634
20170801	nist	G01	-130.5054476	15.09977074
20170801	nist	G01	-132.0046807	13.4566465
20170801	nist	G01	-132.4480758	12.99539411
20170801	nist	G01	-132.7421779	12.63842732
20170801	nist	G01	-133.0351353	12.31192551
20170801	nist	G01	-133.1811982	12.14885878
20170801	nist	G01	-133.6177884	11.66044162
20170801	nist	G01	-134.6278479	10.52585338
20170801	nist	G01	-135.0573582	10.04200552
20170801	nist	G01	-135.7692146	9.239154641
20170801	nist	G01	-135.9110309	9.079150033
20170801	nist	G01	-136.6176024	8.282141334
20170801	nist	G01	37.30568197	6.174084087
20170801	nist	G03	-54.52626504	7.077193841
20170801	nist	G03	-54.00587228	8.141795886
20170801	nist	G03	-53.58448261	9.035815362
20170801	nist	G03	-53.0138667	10.29793155
20170801	nist	G03	-52.69821366	11.02464144
20170801	nist	G03	-52.23937367	12.12220806
20170801	nist	G03	-51.65583811	13.59961487
20170801	nist	G03	-51.03968045	15.28072659
20170801	nist	G03	-49.18531427	21.61365644
20170801	nist	G03	-49.05151087	22.20212638
20170801	nist	G03	-48.93403879	22.79263748
20170801	nist	G03	-169.9708051	5.614171974
20170801	nist	G03	-170.3331587	4.891932846
20170801	nist	G03	33.31001092	5.750732988
20170801	nist	G05	62.6161067	5.088237382
20170801	nist	G05	-48.62137304	9.095059696
20170801	nist	G05	-53.6371151	9.398497836

(*see Figure 5*)

And then, I'm going to use it to make sky plot to show the jump point in the sky plot. It can directly show where jump occurrence in the universe (see Figure 6). This figure is for station yell, which located on Yellowknife city in Canada, for 2017 whole year. We can see the jump point occur in the similarity on the different dates.



(*see Figure 6*)