

Part 9: Adult Advanced Life Support: 2025 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care

Jane G. Wigginton, MD, MSCS, Vice Chair; Sachin Agarwal, MD, MPH; Jason A. Bartos, MD, PhD; Ryan A. Coute, DO; Ian R. Drennan, ACP, PhD; Ameera Haamid, MD; Peter J. Kudenchuk, MD; Mark S. Link, MD; Ashish R. Panchal, MD, PhD; Michele M. Pelter, RN, PhD; Marina Del Rios, MD, MS; Amber J. Rodriguez, PhD; Sarah M. Perman, MD, MSCE; Stephen Sanko, MD; Pavitra Kotini-Shah, MD; Michael C. Kurz, MD, MS, Chair

ABSTRACT: In these 2025 Advanced Life Support Guidelines, the American Heart Association provides comprehensive recommendations for the resuscitation and management of adults experiencing cardiac arrest, respiratory arrest, and life-threatening cardiovascular emergencies. Based on structured evidence reviews and the latest clinical research, these guidelines offer evidence-based strategies to optimize survival and patient outcomes. The 2025 guidelines provide guidance for the treatment of cardiac arrest, including ventricular fibrillation, pulseless ventricular tachycardia, asystole, and pulseless electrical activity, as well as peri-arrest conditions such as atrial fibrillation and flutter with rapid ventricular response. Recommendations are made for defibrillation, electrical cardioversion, advanced airway management, drug therapies, and intravenous access. Additionally, guidelines are provided for the use of double sequential defibrillation, head-up cardiopulmonary resuscitation, and point-of-care ultrasound in the advanced life support setting. Termination of resuscitation rules have been refined to guide decision-making based on the emergency medical services professional's scope of practice. Finally, these guidelines also underscore the importance of identifying causative versus secondary arrhythmias to inform the appropriate timing of therapeutic strategies.

Key Words: AHA Scientific Statements ■ advanced cardiac life support ■ airway management ■ arrhythmias, cardiac ■ cardiopulmonary resuscitation ■ drug therapy ■ electric countershock ■ emergency medical services

TOP 10 TAKE-HOME MESSAGES FOR ADULT ADVANCED LIFE SUPPORT

1. A rapid assessment of clinical stability is essential to direct the appropriate advanced life support (ALS) treatment, and these guidelines go into greater depth to describe how poor organ perfusion manifests as instability.
2. Higher first-shock energy settings (≥ 200 J) are preferable to lower settings for cardioversion of atrial fibrillation and atrial flutter.
3. Updated termination of resuscitation (TOR) guidelines emphasize rule application based on emergency medical services (EMS) scope of practice (basic life support [BLS], ALS, or universal TOR rule [UTOR]), and that end-tidal carbon dioxide (ETCO_2) should not be used in isolation to end resuscitative efforts.
4. The usefulness of vector change (VC) and double sequential defibrillation (DSD) has not been established as therapies for shock-refractory ventricular fibrillation (VF); however, further investigation of these techniques, patient candidacy, and the development of new technology to optimize shock delivery are necessary.
5. Head-up cardiopulmonary resuscitation (CPR) use is discouraged outside of the setting of rigorous clinical trials with appropriate subject protections.
6. Recommendations regarding outdated or extraordinary procedures that have been replaced by

modern equivalents with better efficacy (eg, administration of intra-arrest medications via an in-place endotracheal tube) have been removed.

7. Use of point of care ultrasonography (POCUS) by experienced professionals during cardiac arrest may be considered to diagnose reversible causes if it can be done without interrupting resuscitative efforts (ie, CPR).
8. Polymorphic ventricular tachycardia is always unstable and should be treated immediately with defibrillation, because delays in shock delivery worsen outcomes.
9. Intravenous (IV) access remains the first-line choice for drug administration during cardiac arrest; however, intraosseous (IO) access is a reasonable alternative if IV access is not feasible or delayed.
10. Arrhythmias can be both the cause of and a manifestation of clinical instability. Evaluating the proximal cause of that instability will direct professionals to the most judicious use of these guidelines.

PREAMBLE

In the United States, cardiac arrest remains a significant public health challenge. The incidence of EMS-treated OHCA is 83.4 individuals per 100 000 population annually.¹ Despite advancements in resuscitation science, training, and systems of care, survival to hospital discharge after EMS-treated OHCA remains low, at approximately 10%. Immediate interventions, such as high-quality CPR and timely defibrillation, are crucial for improving outcomes.

In-hospital cardiac arrest (IHCA) occurs in approximately 1 in 100 hospitalized adults. Data indicates that about 72.2% of these patients achieve return of spontaneous circulation (ROSC), with survival to hospital discharge rates around 24.2%. Among those who survive to discharge, approximately 85% have favorable neurological outcomes. These statistics underscore the importance of ALS interventions, including advanced airway management, pharmacologic therapies, and coordinated postresuscitation care, in both prehospital and in-hospital settings.

The International Liaison Committee on Resuscitation (ILCOR) emphasizes the “Formula for Survival,” which identifies 3 critical components for improved outcomes: guidelines based on sound science, effective education of resuscitation professionals, and well-functioning systems of care.² The 2025 ALS guidelines build upon these principles by incorporating the latest evidence, promoting high-quality education for ALS professionals, and describing the critical decisions required to provide life-saving emergency cardiovascular care.

This document outlines the American Heart Association (AHA) recommendations for ALS interventions, drawing from rigorous evidence evaluation and expert consensus. Recommendations are tailored to the scope and practice of ALS

professionals, focusing on enhancing survival and functional recovery. The 2025 ALS guidelines aim to provide a framework for continued advancements in resuscitation science while highlighting areas for future research and innovation.¹

REFERENCES

1. Martin SS, Aday AW, Allen NB, Almarzooq ZI, Anderson CAM, Arora P, Avery CL, Baker-Smith CM, Bansal N, Beaton AZ, et al; American Heart Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics Committee. 2025 Heart disease and stroke statistics: a report of US and global data from the American Heart Association. *Circulation*. 2025;151:e41–e660. doi: 10.1161/CIR.0000000000001303
2. Soreide E, Morrison L, Hillman K, Monsieurs K, Sunde K, Zideman D, Eisenberg M, Sterz F, Nadkarni VM, Soar J, et al; Utstein Formula for Survival Collaborators. The formula for survival in resuscitation. *Resuscitation*. 2013;84:1487–1493. doi: 10.1016/j.resuscitation.2013.07.020

INTRODUCTION TO THE 2025 ALS GUIDELINES

Scope of the Guidelines

The 2025 ALS guidelines provide evidence-based recommendations for health care professionals managing adult cardiac arrest and life-threatening cardiovascular emergencies. These guidelines are tailored to professionals with advanced training in resuscitation techniques, incorporating a wide range of pharmacologic and device-based therapies and critical interventions to optimize survival and neurological outcomes.

This document complements the separately published 2025 BLS guidelines, which address foundational resuscitation practices for lay rescuers and BLS- and ALS-trained health care professionals. Importantly, the ALS guidelines address prehospital resuscitation, in-hospital resuscitation, and TOR recommendations when resuscitation has been unsuccessful. There is a recognition that in the literature there is differential use of the terms *return of spontaneous circulation* (ROSC) versus *return of circulation*. For the purposes of these guidelines, after CPR, when return of circulation is achieved by mechanical means (eg, extracorporeal membrane oxygenation), we use return of circulation; when due to cardiac function, we use ROSC. Specific recommendations about the training of ALS resuscitation professionals are provided in Part 12: Resuscitation Education Science,¹ recommendations about incorporating ALS professionals or procedures into systems of care are provided in Part 4: Systems of Care,² recommendations about caring for the adult patient following ROSC are provided in Part 11: Post-Cardiac Arrest Care,³ and recommendations about ethical considerations in delivering or withholding ALS care are provided in Part 3: Ethical Considerations of Resuscitation.⁴

Organization of the Writing Group

The ALS Writing Group included experts in emergency medicine, EMS, cardiology, critical care, public health, nurs-

ing, and education, along with AHA staff and science editors. Members disclosed potential conflicts of interest before and every 90 days during appointment, aligning with the AHA's policies to ensure unbiased guideline development. Disclosure information for writing group members is in Appendix 1.

METHODOLOGY AND EVIDENCE REVIEW

These guidelines are based on evidence evaluation conducted by the ALS Writing Group.⁵ Evidence review processes included systematic reviews, scoping reviews, and annual focused evidence updates, composed by ILCOR, affiliates, and individual Writing Group Members, to support robust and transparent guideline development. Each of these resulted in a description of the literature and exhaustive discussion among Writing Group members that facilitated guideline development. Details of evidence evaluation can be found in Part 2: Evidence Evaluation and Guidelines Development.⁶

Class of Recommendation and Level of Evidence

Each recommendation in the 2025 ALS guidelines is assigned a Class of Recommendation (COR) to reflect its strength and a Level of Evidence (LOE). The LOE is based on the quality, quantity, relevance, and consistency of the available evidence. In determining the COR, the writing group considered the LOE and other factors, including systems issues, economic factors, and ethical factors such as equity, acceptability, and feasibility. Criteria for each COR and LOE are described in Table 1. For further explanation, refer to Part 2: Evidence Evaluation and Guidelines Development.⁶

Guideline Structure

The 2025 ALS guidelines are organized into knowledge chunks, which are modular, and organized into sections addressing specific clinical topics.⁷ Each module includes a table of recommendations, a brief synopsis, supporting rationale, and references. Flow diagrams and tables clarify the cascade of clinical decision-making, and hyperlinked references provide access to studies and resources.

Document Review and Approval

Each section of the 2025 ALS guidelines underwent rigorous review by subject matter experts and independent peer reviewers. All guidelines were reviewed and approved for publication by the AHA Science Advisory and Coordinating Committee and the AHA Executive Committee. Disclosure information for peer reviewers is listed in Appendix 2.

REFERENCES

1. Donoghue A, Auerbach M, Banerjee A, Blewer A, Cheng A, Kadlec K, Lin Y, Diederich E, Sawyer T, Stallings D, et al. Part 12: resuscitation educa-

tion science: 2025 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2025;152(suppl 2):S719–S750. doi: 10.1161/CIR.0000000000001374

2. Dezfulian C, Cabañas JG, Buckley JR, Cash RE, Crowe RP, Drennan IR, Mahgoub M, Mannarino CN, May T, Salcido DD, et al. Part 4: systems of care: 2025 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2025;152(suppl 2):S353–S384. doi: 10.1161/CIR.0000000000001378

3. Hirsch KG, Amorim E, Coppler PJ, Drennan IR, Elliott A, Gordon AJ, Jentzer JC, Johnson NJ, Moskowitz A, et al. Part 11: post-cardiac arrest care: 2025 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2025;152(suppl 2):S673–S718. doi: 10.1161/CIR.0000000000001375

4. Elmer J, Atkins DL, Daya MR, Del Rios M, Fry JT, Henderson CM, Lewis-Newby M, Madrigal VN, Marco CA, Ornato JP, et al. Part 3: ethics: 2025 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2025;152(suppl 2):S323–S352. doi: 10.1161/CIR.0000000000001371

5. Berg KM, Bray JE, Ng KC, Liley HG, Greif R, Carlson JN, Morley PT, Drennan IR, Smyth M, Scholefield BR, et al; Collaborators. 2023 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations: Summary From the Basic Life Support; Education, Implementation, and Teams; and First Aid Task Forces. *Circulation*. 2023;148:e187–e280. doi: 10.1161/CIR.0000000000001179

6. Panchal AR, Bartos JA, Wyckoff MH, Drennan IR, Mahgoub M, Schexnayder SM, Rodriguez AJ, Sasson C, Wright JI, Brooks SC, et al. Part 2: evidence evaluation and guidelines development: 2025 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2025;152(suppl 2):S313–S322. doi: 10.1161/CIR.0000000000001373

7. Levine GN, O’Gara PT, Beckman JA, Al-Khatib SM, Birtcher KK, Cigarroa JE, de las Fuentes L, Deswal A, Fleisher LA, Gentile F, et al. Recent Innovations, Modifications, and Evolution of ACC/AHA Clinical Practice Guidelines: An Update for Our Constituencies: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. 2019;139:e879–e886. doi: 10.1161/CIR.0000000000000651

DEFIBRILLATION

Recommendations for Defibrillation Indication, Type, and Energy		
COR	LOE	Recommendations
1	B-NR	1. In adult patients with cardiac arrest, defibrillators (using biphasic or monophasic waveforms) are recommended to treat tachyarrhythmias requiring a shock such as ventricular fibrillation or pulseless ventricular tachycardia.
2a	B-R	2. Biphasic waveform defibrillation is preferred over monophasic waveform defibrillation for treatment of tachyarrhythmias in adult patients with cardiac arrest.
2a	B-NR	3. A single shock strategy is preferred to stacked shocks for defibrillation in adult patients with cardiac arrest.
2a	C-LD	4. It is reasonable that selection of fixed versus escalating energy levels for subsequent shocks for adults in cardiac arrest with presumed shock-refractory arrhythmias be based on the specific manufacturer’s instructions for that waveform.
2b	B-R	5. If using a defibrillator capable of escalating energies, higher energy for second and subsequent shocks may be considered if the initial shock fails to restore a perfusing rhythm in adult patients with cardiac arrest.
2b	C-LD	6. In the absence of conclusive evidence that one biphasic waveform is superior to another in termination of ventricular fibrillation, it may be reasonable to use the manufacturer’s recommended energy dose for the first shock in adult patients with cardiac arrest. If this is not known, defibrillation at the maximal dose may be considered.

Table 1. Applying Class of Recommendation and Level of Evidence to Clinical Strategies, Interventions, Treatments, or Diagnostic Testing in Patient Care (updated December 2024)*

CLASS (STRENGTH) OF RECOMMENDATION	LEVEL (QUALITY) OF EVIDENCE‡
Class 1 (STRONG) Benefit >>> Risk Suggested phrases for writing recommendations: - Is recommended - Is indicated/useful/effective/beneficial - Should be performed/administered/other - Comparative-Effectiveness Phrases†: - Treatment/strategy A is recommended/indicated in preference to treatment B - Treatment A should be chosen over treatment B	Level A - High-quality evidence± from more than 1 RCT - Meta-analyses of high-quality RCTs - One or more RCTs corroborated by high-quality registry studies
	Level B-R (Randomized) - Moderate-quality evidence± from 1 or more RCTs - Meta-analyses of moderate-quality RCTs
Class 2a (MODERATE) Benefit >> Risk Suggested phrases for writing recommendations: - Is reasonable - Can be useful/effective/beneficial - Comparative-Effectiveness Phrases†: - Treatment/strategy A is probably recommended/indicated in preference to treatment B - It is reasonable to choose treatment A over treatment B	Level B-NR (Nonrandomized) - Moderate-quality evidence± from 1 or more well-designed, well-executed nonrandomized studies, observational studies, or registry studies - Meta-analyses of such studies
Class 2b (WEAK) Benefit ≥ Risk Suggested phrases for writing recommendations: - May/might be reasonable - May/might be considered - Usefulness/effectiveness is unknown/unclear/uncertain or not well-established	Level C-LD (Limited Data) - Randomized or nonrandomized observational or registry studies with limitations of design or execution - Meta-analyses of such studies - Physiological or mechanistic studies in human subjects
Class 2b (WEAK) Benefit ≥ Risk Suggested phrases for writing recommendations: - May/might be reasonable - May/might be considered - Usefulness/effectiveness is unknown/unclear/uncertain or not well-established	Level C-EO (Expert Opinion) Consensus of expert opinion based on clinical experience
Class 3: No Benefit (MODERATE) Benefit = Risk (Generally, LOE A or B use only) Suggested phrases for writing recommendations: - Is not recommended - Is not indicated/useful/effective/beneficial - Should not be performed/administered/other	COR and LOE are determined independently (any COR may be paired with any LOE). A recommendation with LOE C does not imply that the recommendation is weak. Many important clinical questions addressed in guidelines do not lend themselves to clinical trials. Although RCTs are unavailable, there may be a very clear clinical consensus that a particular test or therapy is useful or effective. * The outcome or result of the intervention should be specified (an improved clinical outcome or increased diagnostic accuracy or incremental prognostic information). † For comparative-effectiveness recommendations (COR 1 and 2a; LOE A and B only), studies that support the use of comparator verbs should involve direct comparisons of the treatments or strategies being evaluated. ‡ The method of assessing quality is evolving, including the application of standardized, widely-used, and preferably validated evidence grading tools; and for systematic reviews, the incorporation of an Evidence Review Committee.
Class 3: HARM (STRONG) Risk > Benefit Suggested phrases for writing recommendations: - Potentially harmful - Causes harm - Associated with excess morbidity/mortality - Should not be performed/administered/other	COR indicates Class of Recommendation; EO, expert opinion; LD, limited data; LOE, Level of Evidence; NR, nonrandomized; R, randomized; and RCT, randomized controlled trial.

Synopsis

Along with CPR, early defibrillation is critical to survival when sudden cardiac arrest is caused by ventricular fibrillation or pulseless ventricular tachycardia (VF/pVT).^{1,2} Defibrillation is most successful when administered as soon as possible after the onset of VF/pVT. Conversely, when VF/pVT is prolonged, depletion of the heart's energy reserves can compromise the efficacy of defibrillation unless replenished by a prescribed period of CPR before rhythm analysis. Figure 1 describes the algorithm for performing shocks and other ALS interventions. Minimizing disruptions in CPR surrounding shock administration is a high priority. Currently marketed defibrillators use proprietary shock waveforms that differ in their electrical characteristics. These deliver different peak currents even at the same programmed energy setting, making comparisons of shock efficacy between devices chal-

lenging. Technologies have been developed to diagnose the underlying cardiac rhythm during CPR and to derive prognostic information from the ventricular waveform that may guide patient management. These technologies require further validation before routine use.

While investigations continue to evaluate the optimal waveform and current, escalating energy may also be effective in termination of VF or pVT. Furthermore, optimized pad placement for defibrillation is an important factor in shock success (see Recommendation-Specific Supportive Text, items 4 and 5, and the section Vector Change and Double Sequential Defibrillation).

Recommendation-Specific Supportive Text

1. Emergent electric defibrillation is highly effective at terminating VF/pVT and other hemodynamic destabilizing tachyarrhythmias (please




Figure 1. Adult Cardiac Arrest Circular Algorithm.

CPR indicates cardiopulmonary resuscitation; ET, endotracheal; IO, intraosseous; IV, intravenous; pVT, pulseless ventricular tachycardia; and VF, ventricular fibrillation.

see sections on wide-complex and narrow-complex tachycardias or atrial fibrillation/flutter as appropriate).

2. Biphasic waveform defibrillators (which deliver pulses of opposite polarity) expose patients to a much lower peak electric current with equivalent or greater efficacy for terminating atrial⁹ and ventricular tachyarrhythmias than monophasic (single polarity) defibrillators.^{4–10} These potential differences in safety and efficacy favor preferential use of a biphasic defibrillator, when available. Biphasic defibrillators have largely replaced monophasic shock defibrillators which were last commercially manufactured in the late 1990s, however, some may still be in use.
3. The rationale for a single shock strategy, in which CPR is immediately resumed after the first shock rather than after serial “stacked” shocks (if required) is based upon several considerations. These include the high success rate of the first shock with biphasic waveforms (lessening the need for successive shocks), the declining success of immediate second and third serial shocks when the first shock has failed,¹¹ and the protracted interruption in CPR required for a series of stacked shocks. A single shock strategy results in shorter interruptions in CPR and a significantly improved survival to hospital admission and discharge (although not 1-year survival) compared with serial “stacked” shocks.^{12–14} It is unknown whether stacked shocks or single shocks are more effective in settings of a monitored witnessed arrest, specifically, an in-patient cardiac arrest or cardiac arrest after cardiac surgery where the rhythm change is monitored in real time. (See the section on cardiac arrest after cardiac surgery in Part 10.¹⁵)

4 and 5. Commercially available defibrillators either provide fixed energy settings or allow for escalating energy settings; both approaches are highly effective in terminating VF/pVT.¹⁶ An optimal energy setting for initial or subsequent biphasic defibrillation, whether fixed or escalating, has not been identified and is best deferred to the defibrillator's manufacturer. When a manufacturer's specified setting is unknown, another approach is to apply the maximum dose setting for that device. A randomized trial comparing fixed 150 J biphasic defibrillation with escalating higher shock energies (200–300–360 J) observed similar rates of successful defibrillation and conversion to an organized rhythm after the first shock. However, among patients who required multiple shocks, escalating shock energy resulted in a significantly higher rate of

conversion to an organized rhythm, although overall survival did not differ between the 2 treatment groups.¹⁷ An observational study comparing fixed 200 J biphasic defibrillation against escalating (200–300–360 J) shocks had similar findings.¹⁸ Different strategies to increase current delivery from biphasic waveform shock have been described, including optimizing pad-skin contact, applying manual pressure to pads during shock delivery (with appropriate self-protective insulation precautions), or vector change of pads. Vector change and double sequential defibrillation are described below.¹⁹

6. There is no conclusive evidence of superiority of one biphasic shock waveform over another for defibrillation.²⁰ Given the variability in electric characteristics between proprietary biphasic waveforms, energy settings are prespecified by the manufacturer for each specific device.

REFERENCES

1. Swor RA, Jackson RE, Cynar M, Sadler E, Basse E, Boji B, Rivera-Rivera EJ, Maher A, Grubb W, Jacobson R. Bystander CPR, ventricular fibrillation, and survival in witnessed, unmonitored out-of-hospital cardiac arrest. *Ann Emerg Med*. 1995;25:780–784. doi: 10.1016/s0196-0644(95)70207-5
2. Larsen MP, Eisenberg MS, Cummins RO, Hallstrom AP. Predicting survival from out-of-hospital cardiac arrest: a graphic model. *Ann Emerg Med*. 1993;22:1652–1658. doi: 10.1016/s0196-0644(05)81302-2
3. Inácio JF, da Rosa Mdos S, Shah J, Rosário J, Vissoci JR, Manica AL, Rodrigues CG. Monophasic and biphasic shock for transthoracic conversion of atrial fibrillation: Systematic review and network meta-analysis. *Resuscitation*. 2016;100:66–75. doi: 10.1016/j.resuscitation.2015.12.009
4. White RD, Hankins DG, Bugliosi TF. Seven years' experience with early defibrillation by police and paramedics in an emergency medical services system. *Resuscitation*. 1998;39:145–151. doi: 10.1016/s0300-9572(98)00135-x
5. van Alem AP, Chapman FW, Lank P, Hart AA, Koster RW. A prospective, randomised and blinded comparison of first shock success of monophasic and biphasic waveforms in out-of-hospital cardiac arrest. *Resuscitation*. 2003;58:17–24. doi: 10.1016/s0300-9572(03)00106-0
6. Schneider T, Martens PR, Paschen H, Kuisma M, Wolcke B, Gliner BE, Russell JK, Weaver WD, Bossaert L, Chamberlain D. Multicenter, randomized, controlled trial of 150-J biphasic shocks compared with 200- to 360-J monophasic shocks in the resuscitation of out-of-hospital cardiac arrest victims. Optimized Response to Cardiac Arrest (ORCA) Investigators. *Circulation*. 2000;102:1780–1787. doi: 10.1161/01.cir.102.15.1780
7. Morrison LJ, Dorian P, Long J, Vermeulen M, Schwartz B, Sawadsky B, Frank J, Cameron B, Burgess R, Shield J, et al; Steering Committee, Central Validation Committee, Safety and Efficacy Committee. Out-of-hospital cardiac arrest rectilinear biphasic to monophasic damped sine defibrillation waveforms with advanced life support intervention trial (ORBIT). *Resuscitation*. 2005;66:149–157. doi: 10.1016/j.resuscitation.2004.11.031
8. Leng CT, Paradis NA, Calkins H, Berger RD, Lardo AC, Rent KC, Halperin HR. Resuscitation after prolonged ventricular fibrillation with use of monophasic and biphasic waveform pulses for external defibrillation. *Circulation*. 2000;101:2968–2974. doi: 10.1161/01.cir.101.25.2968
9. Higgins SL, O'Grady SG, Banville I, Chapman FW, Schmitt PW, Lank P, Walker RG, Iliina M. Efficacy of lower-energy biphasic shocks for transthoracic defibrillation: a follow-up clinical study. *Prehosp Emerg Care*. 2004;8:262–267. doi: 10.1016/j.prehosp.2004.02.002
10. Didon JP, Fontaine G, White RD, Jekova I, Schmid JJ, Cansell A. Clinical experience with a low-energy pulsed biphasic waveform in out-of-hospital cardiac arrest. *Resuscitation*. 2008;76:350–353. doi: 10.1016/j.resuscitation.2007.08.010
11. Koster RW, Walker RG, Chapman FW. Recurrent ventricular fibrillation during advanced life support care of patients with pre-hospital cardiac arrest. *Resuscitation*. 2008;78:252–257. doi: 10.1016/j.resuscitation.2008.03.231

12. Rea TD, Helbock M, Perry S, Garcia M, Cloyd D, Becker L, Eisenberg M. Increasing use of cardiopulmonary resuscitation during out-of-hospital ventricular fibrillation arrest: survival implications of guideline changes. *Circulation*. 2006;114:2760–2765. doi: 10.1161/CIRCULATIONAHA.106.654715

13. Jost D, Degrange H, Verret C, Hersan O, Banville IL, Chapman FW, Lank P, Petit JL, Fuilla C, Migliani R, et al; DEFI 2005 Work Group. DEFI 2005: a randomized controlled trial of the effect of automated external defibrillator cardiopulmonary resuscitation protocol on outcome from out-of-hospital cardiac arrest. *Circulation*. 2010;121:1614–1622. doi: 10.1161/CIRCULATIONAHA.109.878389

14. Bobrow BJ, Clark LL, Ewy GA, Chikani V, Sanders AB, Berg RA, Richman PB, Kern KB. Minimally interrupted cardiac resuscitation by emergency medical services for out-of-hospital cardiac arrest. *JAMA*. 2008;299:1158–1165. doi: 10.1001/jama.299.10.1158

15. Cao D, Arens AM, Chow SL, Easter SR, Hoffman RS, Lagina AT, Lavonas EJ, Patil KD, Sutherland LD, Tijssen JA, et al. Part 10: adult and pediatric special circumstances of resuscitation: 2025 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2025;152(suppl 2):S578–S672. doi: 10.1161/CIR.0000000000001378

16. Hess EP, Russell JK, Liu PY, White RD. A high peak current 150-J fixed-energy defibrillation protocol treats recurrent ventricular fibrillation (VF) as effectively as initial VF. *Resuscitation*. 2008;79:28–33. doi: 10.1016/j.resuscitation.2008.04.028

17. Stiell IG, Walker RG, Nesbitt LP, Chapman FW, Cousineau D, Christenson J, Bradford P, Sookram S, Berringer R, Lank P, et al. BIPHASIC Trial: a randomized comparison of fixed lower versus escalating higher energy levels for defibrillation in out-of-hospital cardiac arrest. *Circulation*. 2007;115:1511–1517. doi: 10.1161/CIRCULATIONAHA.106.648204

18. Tang H, Wu R, Lin L, Hao W, Shi J, Zhu H, Shengyong X, Xu J. Escalating vs fixed energy defibrillation in out-of-hospital cardiac arrest ventricular fibrillation. *JAMA Netw Open*. 2025;8:E257411. doi: 10.1001/jamanetworkopen.2025.7411

19. Deakin CD, Sado DM, Petley GW, Clewlow F. Determining the optimal paddle force for external defibrillation. *Am J Cardiol*. 2002;90:812–813. doi: 10.1016/s0002-9149(02)02623-1

20. Morrison LJ, Henry RM, Ku V, Nolan JP, Morley P, Deakin CD. Single-shock defibrillation success in adult cardiac arrest: a systematic review. *Resuscitation*. 2013;84:1480–1486. doi: 10.1016/j.resuscitation.2013.07.008

ANCILLARY WAVEFORM TECHNOLOGY

Recommendations for Ancillary Waveform Technology		
COR	LOE	Recommendations
2b	C-LD	1. The effectiveness of artifact-filtering algorithms for analysis of electrocardiogram rhythms during chest compressions for adults in cardiac arrest has not been established.
2b	C-LD	2. The effectiveness of VF waveform analysis to guide the acute management of adults with cardiac arrest has not been established.

Synopsis

CPR obscures interpretation of the underlying rhythm because of the artifact created by chest compressions on the electrocardiogram. This makes it difficult to plan the next step of care and can potentially delay or even misdirect drug therapies if given empirically (blindly) on the basis of the patient's presumed, but not actual, underlying rhythm. Time taken for rhythm analysis also disrupts CPR. Artifact-filtering and other innovative techniques to identify the underlying rhythm beneath ongoing CPR can surmount these challenges and minimize interruptions in chest compressions while offering a diagnostic advantage to better direct therapies.^{1–6} These still require further testing and validation before routine use.

Recommendation-Specific Supportive Text

1. Despite the theoretical advantages, no study has evaluated artifact-filtering technologies in a real-time clinical setting or validated their clinical effectiveness compared to current resuscitation strategies. At present, filtering algorithms are strictly used for visual (manual) rhythm interpretation and not for automated VF/VT rhythm detection in automated external defibrillators during ongoing CPR. Further investigation and clinical validation are necessary before these technologies are adopted into routine clinical practice.⁷
2. The electric characteristics of the VF waveform are known to change over time.⁸ VF waveform analysis may be of value in predicting the success of defibrillation or other therapies during the course of resuscitation.^{9–11} The prospect of basing therapies on a prognostic analysis of the VF waveform in real time is an exciting and developing avenue of new research. However, the validity, reliability, and clinical effectiveness of an approach that prompts or withholds shock or other therapies on the basis of predictive analyses is currently uncertain. The only prospective clinical trial comparing a standard shock-first protocol with a waveform analysis-guided shock algorithm was underpowered, however, observed no difference in ROSC.¹² The consensus of the writing group is that there is currently insufficient evidence to support the routine use of waveform analysis to guide resuscitation care, but it is an area in which further research with clinical validation is needed and encouraged.

REFERENCES

1. Tan Q, Freeman GA, Geheb F, Bisera J. Electrocardiographic analysis during uninterrupted cardiopulmonary resuscitation. *Crit Care Med*. 2008;36:S409–S412. doi: 10.1097/ccm.0b013e31818a77bf

2. Li Y, Bisera J, Weil MH, Tang W. An algorithm used for ventricular fibrillation detection without interrupting chest compression. *IEEE Trans Biomed Eng*. 2012;59:78–86. doi: 10.1109/TBME.2011.2118755

3. Li Y, Bisera J, Tang W, Weil MH. Automated detection of ventricular fibrillation to guide cardiopulmonary resuscitation. *Crit Pathw Cardiol*. 2007;6:131–134. doi: 10.1097/HPC.0b013e31813429b0

4. Hu Y, Tang H, Liu C, Jing D, Zhu H, Zhang Y, Yu X, Zhang G, Xu J. The performance of a new shock advisory algorithm to reduce interruptions during CPR. *Resuscitation*. 2019;143:1–9. doi: 10.1016/j.resuscitation.2019.07.026

5. Fumagalli F, Silver AE, Tan Q, Zaidi N, Ristagno G. Cardiac rhythm analysis during ongoing cardiopulmonary resuscitation using the Analysis During Compressions with Fast Reconfirmation technology. *Heart Rhythm*. 2018;15:248–255. doi: 10.1016/j.hrthm.2017.09.003

6. Babaeizadeh S, Firoozabadi R, Han C, Helffenbein ED. Analyzing cardiac rhythm in the presence of chest compression artifact for automated shock advisory. *J Electrocardiol*. 2014;47:798–803. doi: 10.1016/j.jelectrocard.2014.07.021

7. Berg KM, Soar J, Andersen LW, Bottiger BW, Cacciola S, Callaway CW, Couper K, Cronberg T, D'Arrigo S, Deakin CD, et al; Adult Advanced Life Support Collaborators. Adult Advanced Life Support: 2020 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2020;142:S92–S139. doi: 10.1161/CIR.0000000000000893

8. Asano Y, Davidenko JM, Baxter WT, Gray RA, Jalife J. Optical mapping of drug-induced polymorphic arrhythmias and torsade de pointes

in the isolated rabbit heart. *J Am Coll Cardiol*. 1997;29:831–842. doi: 10.1016/s0735-1097(96)00588-8

9. Coult J, Kwok H, Sherman L, Blackwood J, Kudenchuk PJ, Rea TD. Ventricular fibrillation waveform measures combined with prior shock outcome predict defibrillation success during cardiopulmonary resuscitation. *J Electrocardiol*. 2018;51:99–106. doi: 10.1016/j.jelectrocard.2017.07.016

10. Coult J, Blackwood J, Sherman L, Rea TD, Kudenchuk PJ, Kwok H. Ventricular Fibrillation Waveform Analysis During Chest Compressions to Predict Survival From Cardiac Arrest. *Circ Arrhythm Electrophysiol*. 2019;12:e006924. doi: 10.1161/CIRCEP.118.006924

11. Callaway CW, Sherman LD, Mosesso VN Jr, Dietrich TJ, Holt E, Clarkson MC. Scaling exponent predicts defibrillation success for out-of-hospital ventricular fibrillation cardiac arrest. *Circulation*. 2001;103:1656–1661. doi: 10.1161/01.cir.103.12.1656

12. Ruggeri L, Fumagalli F, Bernasconi F, Semeraro F, Meessen J, Blanda A, Migliari M, Magliocca A, Gordini G, Fumagalli R, et al. Amplitude Spectrum Area of ventricular fibrillation to guide defibrillation: a small open-label, pseudo-randomized controlled multicenter trial. *EBioMedicine*. 2023;90:104544. doi: 10.1016/j.ebiom.2023.104544

VECTOR CHANGE AND DOUBLE SEQUENTIAL DEFIBRILLATION

Recommendations for Vector Change and Double Sequential Defibrillation		
COR	LOE	Recommendations
2b	B-R	1. The usefulness of vector change for adults in cardiac arrest with persisting VF/pVT after ≥3 consecutive shocks has not been established.
2b	B-R	2. The usefulness of double sequential defibrillation for adults in cardiac arrest with persisting VF/pVT after ≥3 consecutive shocks has not been established.

Synopsis

Combined with CPR, successful defibrillation is essential to survival from cardiac arrest caused by VF/VT.^{1,2} Defibrillation delivers shock energy (joules), creating a current across the thorax from pad-to-pad.³ Its success is influenced by a variety of factors independent of energy setting, including the electrical resistance of tissues encountered (typically highest at the skin-pad interface) that can dramatically reduce current delivery,^{4–7} pad positions (and resulting shock “vector”) that encompass the heart anatomically^{8–11} and concomitant high quality CPR. When these are all properly managed, biphasic shock can be highly effective (>75% success) in terminating VF/pVT.^{12,13}

“Shock-resistant” and “shock-refractory” are commonly used to describe VF/pVT that continues to be seen on rhythm checks after repeated shocks. In some cases this is due to true failure of shock to terminate VF/pVT, resulting in ongoing, incessant VF/pVT. Causes of such shock failure can be related to correctable deficiencies in defibrillation technique that are best addressed before employing alternate shock strategies such as pad repositioning (vector change) or double sequential defibrillation (DSD). Conversely, evidence indicates a far more common mechanism for VF/pVT that continues to be seen on post-shock rhythm checks (≈80% of instances) is not due to shock failure, but rather to VF/pVT’s recurrence

following successful termination by shock, an event that is obscured by the ensuing period of CPR. When CPR is later paused for rhythm/pulse checks, VF/pVT then seen is attributed to shock failure rather than to its recurrence after successful termination, which is more often the actual case.^{12,14,15} This distinction is important since recurrent VF/pVT may be better remedied by post-shock rhythm stabilization therapy rather than altering an already successful defibrillation technique. Thus labels like “shock-refractory” can be misleading when they conflate different mechanisms for post-shock VF/pVT, for which different treatment strategies may be required. Accordingly, we propose using “persisting VF/pVT” in these guidelines for patients who remain in VF/pVT arrest after ≥3 consecutive shocks, when its actual mechanism (incessant VF/pVT due to true shock failure vs recurrent VF/pVT following successful termination by shock) is not known with certainty.

Recommendation-Specific Supportive Text

1 and 2. Prior systematic reviews of low-quality evidence have not reported benefit from DSD for persisting VF.^{16–20} Defibrillator pad relocation, called vector change (VC), an inherent feature of DSD, was evaluated in only one trial as a stand-alone measure. That trial (DOSE-VF), compared standard defibrillation, VC, and DSD in patients with cardiac arrest in whom VF continued to be seen on rhythm checks after 3 standard shocks. It found significant improvement in survival at hospital discharge with VC and DSD compared to standard defibrillation by intention-to-treat, but notably not when trial findings were analyzed by the treatment strategy patients actually received.²¹ Furthermore, in a secondary exploratory analysis a significant survival benefit from DSD was only observed in the 17% of study patients in whom VF was incessant, and not in the vast majority (83%) of patients in whom VF recurred after a successful shock.²² The interval between each sequential “double” shock required for successfully terminating VF has also been shown experimentally^{23,24} and demonstrated in DOSE-VF itself²⁵ to require a level of precision (separated by milliseconds) unlikely to be consistently achievable by manual activation of two defibrillators. Based on its review, ILCOR’s 2023 International Consensus on CPR and Emergency Cardiovascular Care Science With Treatment Recommendations (CoSTR) judged the overall supportive evidence as relatively weak when issuing “may be considered” recommendations for VC and DSD.²⁶ The adoption of VC or DSD into routine clinical practice for persisting VF/pVT (by whatever mechanism) thus requires further investigation, given its diagnostic and technological requirements.²⁷ These include technologies that can reliably distinguish recurrent from incessant

Downloaded from http://ahajournals.org by on July 20, 2026

post-shock VF during ongoing CPR,²⁸ provide the precise timing interval needed between shocks, and best direct if, how, when, and in whom such a strategy may be applicable.

REFERENCES

1. Swor RA, Jackson RE, Cynar M, Sadler E, Basse E, Boji B, Rivera-Rivera EJ, Maher A, Grubb W, Jacobson R. Bystander CPR, ventricular fibrillation, and survival in witnessed, unmonitored out-of-hospital cardiac arrest. *Ann Emerg Med.* 1995;25:780–784. doi: 10.1016/s0196-0644(95)70207-5

2. Larsen MP, Eisenberg MS, Cummins RO, Hallstrom AP. Predicting survival from out-of-hospital cardiac arrest: a graphic model. *Ann Emerg Med.* 1993;22:1652–1658. doi: 10.1016/s0196-0644(05)81302-2

3. Kerber RE, Martins JB, Kienle MG, Constantin L, Olshansky B, Hopson R, Charbonnier F. Energy, current, and success in defibrillation and cardioversion: clinical studies using an automated impedance-based method of energy adjustment. *Circulation.* 1988;77:1038–1046. doi: 10.1161/01.cir.77.5.1038

4. Kerber RE, Kouba C, Martins J, Kelly K, Low R, Hoyt R, Ferguson D, Bailey L, Bennett P, Charbonnier F. Advance prediction of transthoracic impedance in human defibrillation and cardioversion: importance of impedance in determining the success of low-energy shocks. *Circulation.* 1984;70:303–308. doi: 10.1161/01.cir.70.2.303

5. Li W, Li J, Wei L, Wang J, Peng L, Wang J, Yin C, Li Y. A framework of current based defibrillation improves defibrillation efficacy of biphasic truncated exponential waveform in rabbits. *Sci Rep.* 2021;11:1586. doi: 10.1038/s41598-020-80521-9

6. Squara F, Elbaum C, Garret G, Liprandi L, Scarlatti D, Bun SS, Mossaz B, Rocher M, Bateau J, Moceri P, et al. Active compression versus standard anterior-posterior defibrillation for external cardioversion of atrial fibrillation: A prospective randomized study. *Heart Rhythm.* 2021;18:360–365. doi: 10.1016/j.hrthm.2020.11.005

7. Sirna SJ, Ferguson DW, Charbonnier F, Kerber RE. Factors affecting transthoracic impedance during electrical cardioversion. *Am J Cardiol.* 1988;62:1048–1052. doi: 10.1016/0002-9149(88)90546-2

8. Guidelines 2000 for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. Part 6: advanced cardiovascular life support: section 2: defibrillation. The American Heart Association in collaboration with the International Liaison Committee on Resuscitation. *Circulation.* 2000;102:190–194.

9. Koster RW, Walker RG, Chapman FW. Recurrent ventricular fibrillation during advanced life support care of patients with pre-hospital cardiac arrest. *Resuscitation.* 2008;78:252–257. doi: 10.1016/j.resuscitation.2008.03.231

10. Esibov A, Chapman FW, Melnick SB, Sullivan JL, Walcott GP. Minor Variations in Electrode Pad Placement Impact Defibrillation Success. *Prehosp Emerg Care.* 2016;20:292–298. doi: 10.3109/10903127.2015.1076095

11. Yeh KH, Chang HC, Lin JF. Paddle position and contact force: an important step to check when troubleshooting for refractory ventricular fibrillation. *Hong Kong J Emerg Med.* 2016;23:289–292. doi: 10.1177/102490791602300506

12. Spies DM, Kiekenap J, Rupp D, Betz S, Kill C, Sassen MC. Time to change the times? Time of recurrence of ventricular fibrillation during OHCA. *Resuscitation.* 2020;157:219–224. doi: 10.1016/j.resuscitation.2020.09.029

13. Verkaik BJ, Walker RG, Taylor TG, Ekkel MM, Marx R, Stieglis R, van Eeden VGM, Doleman LC, Hulleman M, Chapman FW, et al. Defibrillation and refractory ventricular fibrillation. *Eur Heart J.* 2024;46:582–584. doi: 10.1093/eurheartj/ehae767

14. van Alem AP, Post J, Koster RW. VF recurrence: characteristics and patient outcome in out-of-hospital cardiac arrest. *Resuscitation.* 2003;59:181–188. doi: 10.1016/s0300-9572(03)00208-9

15. Bhandari S, Doan J, Blackwood J, Coult J, Kudenchuk P, Sherman L, Rea T, Kwok H. Rhythm profiles and survival after out-of-hospital ventricular fibrillation cardiac arrest. *Resuscitation.* 2018;125:22–27. doi: 10.1016/j.resuscitation.2018.01.037

16. Miraglia D, Ramzy M. Double external defibrillation for shock-refractory ventricular fibrillation cardiac arrest: A step towards standardization. *Am J Emerg Med.* 2021;41:73–79. doi: 10.1016/j.ajem.2020.12.031

17. Li Y, He X, Li Z, Li D, Yuan X, Yang J. Double sequential external defibrillation versus standard defibrillation in refractory ventricular fibrillation: A systematic review and meta-analysis. *Front Cardiovasc Med.* 2022;9:1017935. doi: 10.3389/fcvm.2022.1017935

18. Delorenzo A, Nehme Z, Yates J, Bernard S, Smith K. Double sequential externa defibrillation for refractory ventricular fibrillation out-of-hospital cardiac arrest: A systematic review and meta-analysis. *Resuscitation.* 2019;135:124–129. doi: 10.1016/j.resuscitation.2018.10.025

19. Deakin CD, Morley P, Soar J, Drennan IR. Double (dual) sequential defibrillation for refractory ventricular fibrillation cardiac arrest: A systematic review. *Resuscitation.* 2020;155:24–31. doi: 10.1016/j.resuscitation.2020.06.008

20. Abuelazm MT, Ghanem A, Katamesh BE, Hassan AR, Abdalshafy H, Seri AR, Awad AK, Abdelnabi M, Abdelazeem B. Defibrillation strategies for refractory ventricular fibrillation out- of-hospital cardiac arrest: A systematic review and network meta-analysis. *Ann Noninvasive Electrolcardiol.* 2023;28:e13075. doi: 10.1111/anec.13075

21. Cheskes S, Verbeek PR, Drennan IR, McLeod SL, Turner L, Pinto R, Feldman M, Davis M, Vaillancourt C, Morrison LJ, et al. Defibrillation Strategies for Refractory Ventricular Fibrillation. *N Engl J Med.* 2022;387:1947–1956. doi: 10.1056/NEJMoa2207304

22. Cheskes S, Drennan IR, Turner L, Pandit SV, Dorian P. The impact of alternate defibrillation strategies on shock-refractory and recurrent ventricular fibrillation: A secondary analysis of the DOSE VF cluster randomized controlled trial. *Resuscitation.* 2024;198:110186. doi: 10.1016/j.resuscitation.2024.110186

23. Taylor TG, Melnick SB, Chapman FW, Walcott GP. An investigation of inter-shock timing and electrode placement for double-sequential defibrillation. *Resuscitation.* 2019;140:194–200. doi: 10.1016/j.resuscitation.2019.04.042

24. Johnson EE, Alferness CA, Wolf PD, Smith WM, Ideker RE. Effect of pulse separation between two sequential biphasic shocks given over different lead configurations on ventricular defibrillation efficacy. *Circulation.* 1992;85:2267–2274. doi: 10.1161/01.cir.85.6.2267

25. Rahimi M, Drennan IR, Turner L, Dorian P, Cheskes S. The impact of double sequential shock timing on outcomes during refractory out-of-hospital cardiac arrest. *Resuscitation.* 2024;194:110082. doi: 10.1016/j.resuscitation.2023.110082

26. Berg KM, Bray JE, Ng KC, Liley HG, Greif R, Carlson JN, Morley PT, Drennan IR, Smyth M, Scholefield BR, et al; Collaborators. 2023 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations: Summary From the Basic Life Support; Advanced Life Support; Pediatric Life Support; Neonatal Life Support; Education, Implementation, and Teams; and First Aid Task Forces. *Circulation.* 2023;148:e187–e280. doi: 10.1161/CIR.0000000000001179

27. Kudenchuk PJ. Shocking insights on double defibrillation: How, when and why not? *Resuscitation.* 2019;140:209–210. doi: 10.1016/j.resuscitation.2019.05.022

28. Kwok H, Coult J, Blackwood J, Sotoodehnia N, Kudenchuk P, Rea T. A method for continuous rhythm classification and early detection of ventricular fibrillation during CPR. *Resuscitation.* 2022;176:90–97. doi: 10.1016/j.resuscitation.2022.05.019

NONDEFIBRILLATION ELECTRICAL THERAPIES FOR CARDIAC ARREST

Nondefibrillation Electrical Therapies for Cardiac Arrest		
COR	LOE	Recommendation
3: No Benefit	B-R	1. Routine use of electrical pacing is not recommended during the resuscitation of an established adult cardiac arrest.

Synopsis

In addition to defibrillation, alternative electrical therapies have been explored as possible treatment options during cardiac arrest. Transcutaneous pacing has been studied during cardiac arrest with bradysystolic cardiac rhythm. In theory, the heart will respond to electrical stimuli by producing myocardial contraction and generating forward movement of blood, but clinical trials have not shown pacing to improve patient outcomes.

Other pseudo-electrical therapies, such as cough CPR, fist or percussion pacing, and precordial thump have all been described as temporizing measures in select patients who are either peri-arrest or in the initial seconds of witnessed cardiac arrest (before losing consciousness in the case of cough CPR) when definitive therapy is not readily available.¹ These therapies are described elsewhere (see Part 10. Special Circumstances² for more on precordial thump, fist pacing, and cough CPR).

Recommendation-Specific Supportive Text

- 1. Existing evidence, including observational and quasi-randomized controlled trial (RCT) data, suggests that pacing by a transcutaneous, transvenous, or transmyocardial approach during cardiac arrest does not improve the likelihood of ROSC or survival, regardless of the timing of pacing administration in established asystole, location of arrest (in-hospital or out-of-hospital), or primary cardiac rhythm (asystole, pulseless electrical activity).³⁻⁷ Protracted interruptions in chest compressions while the success of pacing is assessed can be detrimental to survival. Specifically, attempts at electrical pacing may delay evidence-based resuscitative measures, such as CPR. It is not known whether the timing of pacing initiation may influence pacing success such that pacing may be useful in the initial minute of select cases of witnessed, monitored cardiac arrest (see Part 10. Special Circumstances, Cardiac Arrest After Cardiac Surgery²). If pacing is attempted during cardiac arrest related to the special circumstances described above, professionals are cautioned that its performance could be at the expense of high-quality CPR. Of note, this recommendation addressing electrical pacing is specifically focused upon patients in cardiac arrest and does not address prevention of cardiac arrest nor does it address the utility of overdrive pacing for arrhythmia.

REFERENCES

1. Tucker KJ, Shaburhivi TS, Gedevarishvili AT. Manual external (fist) pacing during high-degree atrioventricular block: a lifesaving intervention. *Am J Emerg Med.* 1995;13:53-54. doi: 10.1016/0735-6757(95)90243-0

2. Cao D, Arens AM, Chow SL, Easter SR, Hoffman RS, Lagina AT, Lavonas EJ, Patil KD, Sutherland LD, Tijssen JA, et al. Part 10: adult and pediatric special circumstances of resuscitation: 2025 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation.* 2025;152(suppl 2):S578-S672. doi: 10.1161/CIR.0000000000001378

3. White JD, Brown CG. Immediate transthoracic pacing for cardiac asystole in an emergency department setting. *Am J Emerg Med.* 1985;3:125-128. doi: 10.1016/0735-6757(85)90034-8

4. Sherbino J, Verbeek PR, MacDonald RD, Sawadsky BV, McDonald AC, Morrison LJ. Prehospital transcutaneous cardiac pacing for symptomatic bradycardia or bradyasystolic cardiac arrest: a systematic review. *Resuscitation.* 2006;70:193-200. doi: 10.1016/j.resuscitation.2005.11.019

5. Hedges JR, Syverud SA, Dalsey WC, Feero S, Easter R, Shultz B. Pre-hospital trial of emergency transcutaneous cardiac pacing. *Circulation.* 1987;76:1337-1343. doi: 10.1161/01.cir.76.6.1337

6. Cummins RO, Graves JR, Larsen MP, Hallstrom AP, Hearne TR, Ciliberti J, Nicola RM, Horan S. Out-of-hospital transcutaneous pacing by emergency medical technicians in patients with asystolic cardiac arrest. *N Engl J Med.* 1993;328:1377-1382. doi: 10.1056/NEJM199305133281903

7. Barthell E, Troiano P, Olson D, Stueven HA, Hendley G. Prehospital external cardiac pacing: a prospective, controlled clinical trial. *Ann Emerg Med.* 1988;17:1221-1226. doi: 10.1016/s0196-0644(88)80074-x

VASCULAR ACCESS IN CARDIAC ARREST MANAGEMENT

Recommendations for Vascular Access in Cardiac Arrest Management		
COR	LOE	Recommendations
1	A	1. It is recommended that professionals first attempt establishing intravenous access for drug administration in adult patients in cardiac arrest.
2a	A	2. Intraosseous access is reasonable if initial attempts at intravenous access are unsuccessful or not feasible for adult patients in cardiac arrest.
2b	C-LD	3. For appropriately trained professionals, central venous access may be considered for adult patients in cardiac arrest if attempts to establish intravenous and intraosseous access are unsuccessful or not feasible.

Synopsis

Traditionally, peripheral IV access has been used to administer medication during cardiac arrest. However, obtaining IV access under emergent conditions can be challenging because of patient characteristics and operator experience leading to delay in pharmacological treatments. Alternatives to IV access for drug administration include IO, central venous, intracardiac, and endotracheal routes. Intracardiac drug administration was discouraged in the “2000 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care” due to its specialized skill set requirements, potential for morbidity, and the availability of other access options.¹ Similarly, drug administration via secured endotracheal tube results in low blood concentrations and unpredictable pharmacological effects and has also been removed from the guidelines.² Central venous access is primarily used in a hospital setting because it requires specialized training to acquire and maintain proficiency.

Recommendation-Specific Supportive Text

- 1 and 2. The peripheral IV route for vascular access has traditionally been preferred for emergency drug and fluid administration during adult resuscitation. The pharmacokinetic properties, acute effects, and clinical efficacy of emergency drugs have primarily been described for IV administration.³⁻⁶ However, observational studies have noted a significant increase in the use of IO access in adult OHCA, despite the absence of high-quality evidence.^{7,8} Three recent large RCTs evaluated the clinical effectiveness of initial IO access compared with

Downloaded from http://ahajournals.org by on July 20, 2026

initial IV access in adult OHCA and found no differences in clinical outcomes.^{9–11} Each RCT used a superiority design; therefore, the absence of outcome differences between the two groups does not indicate equivalence. An ILCOR systematic review, including data from these RCTs, found that the use of IO access compared with IV access did not result in a statistically significant improvement in outcomes, including survival to discharge, survival with favorable neurological outcome, or health-related quality of life.¹² This systematic review noted lower odds of achieving sustained ROSC for the IO route compared with the IV route (odds ratio [OR] 0.89; 95% CI, 0.80–0.99).¹² Patient, EMS professional, or circumstantial characteristics may limit successful IV access or make IV access infeasible. In these cases, the available evidence supports IO access as an alternative. The optimal anatomical location for IO access (ie, tibial or humeral) remains a knowledge gap.

3. Drug administration through central venous access can achieve faster circulation times and higher plasma concentrations in adults in cardiac arrest when compared with peripheral IV administration.^{13–15} However, data comparing clinical outcomes from cardiac arrest based on different access routes, including central venous access, remain limited. A small, single-center study reported higher rates of ROSC among participants who were randomized to femoral or internal jugular vein access compared to those with peripheral IV access, although with a high risk of bias.¹⁶ If central venous access is attempted by appropriately trained professionals, special attention is required to avoid delays in chest compressions or defibrillation.

REFERENCES

1. Guidelines 2000 for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. Part 6: advanced cardiovascular life support: section 2: defibrillation. The American Heart Association in collaboration with the International Liaison Committee on Resuscitation. *Circulation*. 2000;102:I90–I94.

2. Panchal AR, Bartos JA, Cabanas JG, Donnino MW, Drennan IR, Hirsch KG, Kudenchuk PJ, Kurz MC, Lavonas EJ, Morley PT, et al; Adult Basic and Advanced Life Support Writing Group. Part 3: Adult Basic and Advanced Life Support: 2020 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2020;142:S366–S468. doi: 10.1161/CIR.0000000000000916

3. Riva E, Gerna M, Latini R, Giani P, Volpi A, Maggioni A. Pharmacokinetics of amiodaron in man. *J Cardiovasc Pharmacol*. 1982;4:264–269. doi: 10.1097/00005344-198203000-00015

4. Orlowski JP, Porembka DT, Gallagher JM, Lockrem JD, VanLente F. Comparison study of intraosseous, central intravenous, and peripheral intravenous infusions of emergency drugs. *Am J Dis Child*. 1990;144:112–117. doi: 10.1001/archpedi.1990.02150250124049

5. Greenblatt DJ, Bolognini V, Koch-Weser J, Harmatz JS. Pharmacokinetic approach to the clinical use of lidocaine intravenously. *JAMA*. 1976;236:273–277.

6. Collinsworth KA, Kalman SM, Harrison DC. The clinical pharmacology of lidocaine as an antiarrhythmic drug. *Circulation*. 1974;50:1217–1230. doi: 10.1161/01.cir.50.6.1217

7. Vadeyar S, Buckle A, Hooper A, Booth S, Deakin CD, Fothergill RJ, Ji C, Nolan JP, Brown M, Cowley A, et al. Trends in use of intraosseous and intravenous access in out-of-hospital cardiac arrest across English ambulance services: A registry-based, cohort study. *Resuscitation*. 2023;191:109951. doi: 10.1016/j.resuscitation.2023.109951

8. Agostinucci JM, Alheritiere A, Metzger J, Nadiras P, Martineau L, Bertrand P, Gentilhomme A, Petrovic T, Adnet F, Lapostolle F. Evolution of the use of intraosseous vascular access in prehospital advanced cardiopulmonary resuscitation: The IOVA-CPR study. *Int J Nurs Pract*. 2024;30:e13244. doi: 10.1111/ijn.13244

9. Vallentin MF, Granfeldt A, Klitgaard TL, Mikkelsen S, Folke F, Christensen HC, Povlsen AL, Petersen AH, Winther S, Frilund LW, et al. Intraosseous or Intravenous Vascular Access for Out-of-Hospital Cardiac Arrest. *N Engl J Med*. 2025;392:349–360. doi: 10.1056/NEJMoa2407616

10. Ko YC, Lin HY, Huang EP, Lee AF, Hsieh MJ, Yang CW, Lee BC, Wang YC, Yang WS, Chien YC, et al. Intraosseous versus intravenous vascular access in upper extremity among adults with out-of-hospital cardiac arrest: cluster randomised clinical trial (VICTOR trial). *BMJ*. 2024;386:e079878. doi: 10.1136/bmj-2024-079878

11. Couper K, Ji C, Deakin CD, Fothergill RT, Nolan JP, Long JB, Mason JM, Michelet F, Norman C, Nwankwo H, et al. A randomized trial of drug route in out-of-hospital cardiac arrest. *N Engl J Med*. 2024;392:336–348. doi: 10.1056/NEJMoa2407780

12. Couper K, Drennan IR, Garrett JS, Grunau BE, Vallentin MF. Intraosseous and intravenous vascular access during adult cardiac arrest: a systematic review update. *Resuscitation*. 2025;209:110564. doi: 10.1016/j.resuscitation.2025.110564

13. Kuhn GJ, White BC, Swetnam RE, Mumey JF, Rydesky MF, Tintinalli JE, Krome RL, Hoehner PJ. Peripheral vs central circulation times during CPR: a pilot study. *Ann Emerg Med*. 1981;10:417–419. doi: 10.1016/s0196-0644(81)80308-3

14. Emerman CL, Pinchak AC, Hancock D, Hagen JF. Effect of injection site on circulation times during cardiac arrest. *Crit Care Med*. 1988;16:1138–1141. doi: 10.1097/00003246-198811000-00011

15. Barsan WG, Levy RC, Weir H. Lidocaine levels during CPR: differences after peripheral venous, central venous, and intracardiac injections. *Ann Emerg Med*. 1981;10:73–78. doi: 10.1016/s0196-0644(81)80339-3

16. Qin H, Wang L, Yu B, Xing D, Su J, Bai Z. Effect of other venous access on cardiopulmonary resuscitation quality: A prospective, randomized, controlled trial. *Biotechnol Genet Eng Rev*. 2024;40:2262–2272. doi: 10.1080/02648725.2023.2199239

RECOMMENDATIONS FOR VASOPRESSOR MEDICATIONS IN ADULT CARDIAC ARREST

Recommendations for Vasopressor Management in Cardiac Arrest		
COR	LOE	Recommendations
1	B-R	1. Administration of epinephrine is recommended for adult patients in cardiac arrest.
2a	B-R	2. It is reasonable to administer epinephrine (1 mg) every 3 to 5 min for adult patients in cardiac arrest.
2a	B-NR	3. With respect to timing, for adult patients in cardiac arrest with a nonshockable rhythm, it is reasonable to administer epinephrine as soon as feasible.
2a	B-NR	4. With respect to timing, for adult patients in cardiac arrest with a shockable rhythm, it is reasonable to administer epinephrine after initial defibrillation attempts have failed.
3: No Benefit	B-R	5. Vasopressin alone or vasopressin in combination with epinephrine offers no advantage as a substitute for epinephrine for adult patients in cardiac arrest.
3: No Benefit	B-R	6. High-dose epinephrine is not recommended for routine use for adult patients in cardiac arrest.

Synopsis

The potent vasopressor effects of epinephrine have been recognized for nearly 150 years. Vasopressors increase coronary perfusion pressure and increase the likelihood of ROSC. Epinephrine has been shown to increase ROSC and survival to hospital admission in placebo-controlled RCTs, cohort studies, and registry studies, but there is limited evidence to support improvement in survival to hospital discharge or functional neurologic outcome. Studies comparing different dosing intervals, high-dose epinephrine, and low-dose epinephrine against standard-dose epinephrine (1 mg every 3–5 minutes) have not demonstrated an advantage in survival outcomes. Defining the optimal number of doses or maximum dose of epinephrine is a critical knowledge gap and more research is needed.

Vasopressin, a naturally occurring antidiuretic hormone, has been studied as an alternative to epinephrine during cardiac resuscitation. In high doses, vasopressin acts as a potent vasoconstrictor and, like epinephrine, increases systemic vascular resistance and raises coronary perfusion pressure. However, there have been no clinical trials demonstrating an advantage to using vasopressin alone or vasopressin in addition epinephrine over standard-dose epinephrine alone.

Recommendation-Specific Supportive Text

1. There have been no new RCTs comparing epinephrine to placebo since the publication of the 2020 ALS guidelines. There is consistent and compelling evidence from previous RCTs, cohort, and registry studies to support a potent effect of epinephrine on ROSC, survival to hospital admission, and survival to hospital discharge.^{1,2} However, studies have failed to show an increase in survival with functional neurologic outcome.^{2–5} A secondary analysis of the PARAMEDIC2 (Prehospital Assessment of the Role of Adrenaline: Measuring the Effectiveness of Drug Administration in Cardiac Arrest) trial found 12-month survival favored epinephrine, whereas there was no significance difference in 6-month survival with functional neurologic outcome between arms.⁶ While there is no evidence supporting improvement in neurologic outcome, epinephrine does improve short-term survival, a prerequisite to meaningful recovery. In the absence of a feasible, intra-arrest method of determining the likelihood of favorable neurological outcome, epinephrine remains standard therapy to treat cardiac arrest.
2. Existing clinical trials have used a protocol of 1 mg of epinephrine administered every 3 to 5 minutes.^{3–6} Operationally, administering epinephrine every second cycle of CPR, after the initial dose, meets this recommendation.
3. Multiple observational studies have demonstrated an association between earlier epinephrine administration and ROSC.² A post-hoc analysis of the PARAMEDIC2 trial found that the effectiveness of epinephrine, compared to placebo, converges at approximately 20 minutes of pulselessness where there is no difference in survival to hospital discharge, 30-day survival, or functional neurologic outcome with longer times to initial epinephrine administration.⁷
4. Systematic reviews and meta-analyses of RCTs and cohort studies have found that epinephrine is associated with improved ROSC in all rhythms, but the benefit in shockable rhythms is less than in nonshockable rhythms.^{2–4,8} One observational study of IHCA demonstrated 10% lower risk-adjusted survival in hospitals with the highest rates of epinephrine administration prior to initial defibrillation when compared with those hospitals with the lowest rates.⁹ The optimal timing for epinephrine in relation to defibrillation is unknown. In patients with shockable rhythms, this literature supports prioritizing rapid defibrillation and administering epinephrine after initial attempts with CPR and defibrillation are not successful.
5. When compared to placebo, administration of vasopressin improves ROSC regardless of presenting rhythms and increases survival to hospital admission and discharge in nonshockable rhythms.¹⁰ However, multiple systematic reviews and meta-analyses of RCTs and observational studies have found no difference in survival outcomes when comparing vasopressin alone or vasopressin combined with epinephrine versus epinephrine alone.^{2,10–12}
6. Multiple RCTs have compared administration of high-dose epinephrine with standard-dose, but the definition of high-dose epinephrine varies widely by study. There were no new RCTs published since the 2020 Guidelines comparing standard-dose epinephrine to any high-dose epinephrine. Systematic reviews and meta-analysis of RCTs have yielded mixed results.^{10,12} One registry-based study found dosing epinephrine more frequently than the standard interval may be potentially harmful.¹³ High-dose intramuscular (IM) epinephrine for the treatment of OHCA, with a different pharmacokinetic profile than traditional IV administration, has also been evaluated. One single-center, before-after study of 1405 OHCA patients receiving an initial IM dose of epinephrine followed by standard IV/IO administration shortened time to epinephrine administration and improved survival to hospital admission, hospital survival, and favorable neurologic status at hospital discharge. However, the study design could not account for multiple

confounders, such as temporal trends, misclassification, and resuscitation characteristics.¹⁴ Further study is needed to evaluate the potential benefit of high-dose IM epinephrine in cardiac arrest.

REFERENCES

1. Perkins GD, Ji C, Deakin CD, Quinn T, Nolan JP, Scopin C, Regan S, Long J, Slowther A, Pocock H, et al; PARAMEDIC2 Collaborators. PARAMEDIC2 Collaborators. A Randomized Trial of Epinephrine in Out-of-Hospital Cardiac Arrest. *N Engl J Med*. 2018;379:711–721. doi: 10.1056/NEJMoa1806842

2. Holmberg MJ, Fernando S, Elshaer A, Leong C, Drennan I; International Liaison Committee on Resuscitation Advanced Life Support Task. Vasopressors in adult cardiac arrest. Updated November 3, 2024. Accessed June 20, 2025. <https://costr.ilcor.org/document/vasopressors-in-adult-cardiac-arrest-als-3208-tf-sr>

3. Zhong H, Yin Z, Kou B, Shen P, He G, Huang T, Liang J, Huang S, Huang J, Zhou M, et al. Therapeutic and adverse effects of adrenaline on patients who suffer out-of-hospital cardiac arrest: a systematic review and meta-analysis. *Eur J Med Res*. 2023;28:24. doi: 10.1186/s40001-022-00974-8

4. Ludwin K, Safiejko K, Smereka J, Nadolny K, Cyran M, Yakubtsevich R, Jaguszewski MJ, Filipiak KJ, Szarpak L, Rodríguez-Núñez A. Systematic review and meta-analysis appraising efficacy and safety of adrenaline for adult cardiopulmonary resuscitation. *Cardiol J*. 2021;28:279–292. doi: 10.5603/CJ.a2020.0133

5. Fernando SM, Mathew R, Sadeghirad B, Rochwerf B, Hibbert B, Munshi L, Fan E, Brodie D, Di Santo P, Tran A, et al. Epinephrine in Out-of-Hospital Cardiac Arrest: A Network Meta-analysis and Subgroup Analyses of Shockable and Nonshockable Rhythms. *Chest*. 2023;164:381–393. doi: 10.1016/j.chest.2023.01.033

6. Haywood KL, Ji C, Quinn T, Nolan JP, Deakin CD, Scopin C, Lall R, Gates S, Long J, Regan S, et al. Long term outcomes of participants in the PARAMEDIC2 randomised trial of adrenaline in out-of-hospital cardiac arrest. *Resuscitation*. 2021;160:84–93. doi: 10.1016/j.resuscitation.2021.01.019

7. Perkins GD, Kenna C, Ji C, Deakin CD, Nolan JP, Quinn T, Scopin C, Fothergill R, Gunson I, Pocock H, et al. The influence of time to adrenaline administration in the Paramedic 2 randomised controlled trial. *Intensive Care Med*. 2020;46:426–436. doi: 10.1007/s00134-019-05836-2

8. Perkins GD, Kenna C, Ji C, Deakin CD, Nolan JP, Quinn T, Fothergill R, Gunson I, Pocock H, Rees N, et al. The effects of adrenaline in out of hospital cardiac arrest with shockable and non-shockable rhythms: Findings from the PACA and PARAMEDIC-2 randomised controlled trials. *Resuscitation*. 2019;140:55–63. doi: 10.1016/j.resuscitation.2019.05.007

9. Stewart C, Chan PS, Kennedy K, Swanson MB, Girotra S; American Heart Association's Get With The Guidelines (GWTG) Resuscitation Investigators. American Heart Association's Get With The Guidelines (GWTG) Resuscitation Investigators. Hospital Variation in Epinephrine Administration Before Defibrillation for Cardiac Arrest Due to Shockable Rhythm. *Crit Care Med*. 2024;52:878–886. doi: 10.1097/CCM.00000000000006203

10. Fernando SM, Mathew R, Sadeghirad B, Rochwerf B, Hibbert B, Munshi L, Fan E, Brodie D, Di Santo P, Tran A, et al. Epinephrine in Out-of-Hospital Cardiac Arrest: A Network Meta-analysis and Subgroup Analyses of Shockable and Nonshockable Rhythms. *Chest*. 2023;164:381–393. doi: 10.1016/j.chest.2023.01.033

11. Elbadawi A, Tan BE, Assaf Y, Elzeneini M, Baig B, Hamed M, Elgendy IY, Mamas M. Meta-Analysis of Efficacy of Vasopressin During Cardiopulmonary Resuscitation. *Am J Cardiol*. 2022;181:122–129. doi: 10.1016/j.amjcard.2022.06.042

12. An HR, Han YR, Wang TH, Chi F, Meng Y, Zhang CY, Liang JQ, Li XL. Meta-Analysis of the Factors Influencing the Restoration of Spontaneous Circulation After Cardiopulmonary Resuscitation. *Front Physiol*. 2022;13:834352. doi: 10.3389/fphys.2022.834352

13. Jaeger D, Baert V, Javaudin F, Debaty G, Duhem H, Koger J, Gueugniaud PY, Tazarourte K, El Khoury C, Hubert H, et al; GR-RéAC. Effect of adrenaline dose on neurological outcome in out-of-hospital cardiac arrest: a nationwide propensity score analysis. *Eur J Emerg Med*. 2022;29:63–69. doi: 10.1097/MEJ.0000000000000891

14. Palatinus HN, Johnson MA, Wang HE, Hoareau GL, Youngquist ST. Early intramuscular adrenaline administration is associated with improved survival from out-of-hospital cardiac arrest. *Resuscitation*. 2024;201:110266. doi: 10.1016/j.resuscitation.2024.110266

NONVASOPRESSOR MEDICATIONS DURING CARDIAC ARREST

Recommendations for Nonvasopressor Medications in Adult Cardiac Arrest		
COR	LOE	Recommendations
2b	B-R	1. For adults in cardiac arrest amiodarone or lidocaine may be considered for ventricular fibrillation (VF) or pulseless ventricular tachycardia (pVT) that is unresponsive to defibrillation.
2b	C-LD	2. For adults in cardiac arrest, the use of β -blockers, bretylium, procainamide, or sotalol for VF/pVT unresponsive to defibrillation is of uncertain benefit.
2b	C-LD	3. For adults in cardiac arrest, the use of steroids is of uncertain benefit.
3: No Benefit	B-R	4. For adults in cardiac arrest, routine administration of calcium is not recommended.
3: No Benefit	B-R	5. For adults in cardiac arrest, routine administration of sodium bicarbonate is not recommended.
3: No Benefit	B-R	6. For adults in cardiac arrest, routine administration of magnesium is not recommended.

Synopsis

Pharmacological treatment of cardiac arrest is provided (if indicated) when ROSC is not achieved by CPR and defibrillation.¹ Treatments include vasopressor agents such as epinephrine (discussed in Recommendations for Vasopressor Medications During Cardiac Arrest) as well as drugs without direct hemodynamic effects (nonvasopressors) and their administration is delineated in Figure 2. The latter group includes antiarrhythmic medications (which include β -blockers), magnesium, sodium bicarbonate, calcium, and steroids, many of which are commonly administered by health care professionals, including EMS, during cardiac arrest. Although theoretically attractive and of modest benefit in animal studies, none of these therapies have definitively demonstrated improvement in survival after cardiac arrest. Some, however, offer potential benefit in selected populations and circumstances.

Guidance for the treatment of cardiac arrest due to hyperkalemia, including the use of calcium and sodium bicarbonate for this indication, is presented in Electrolyte Abnormalities within Part 10. Special Circumstances.² In addition, Part 10 offers guidance for the management of cardiac arrest due to toxic ingestion that discusses the administration of these medications in these specific circumstances. Guidance for management of torsades de pointes is presented in Recommendations for Polymorphic VT later in these guidelines.

Recommendation-Specific Supportive Text

1. The 2023 AHA focused update on adult advanced cardiovascular life support³ and ILCOR's 2024 CoSTR summary⁴ addressed administration of parenteral antiarrhythmic medications to patients with cardiac arrest unresponsive to defibrillation.




Figure 2. Adult Cardiac Arrest Algorithm. CPR indicates cardiopulmonary resuscitation; ET, endotracheal; IO, intraosseous; IV, intravenous; PEA, pulseless electrical activity; pVT, pulseless ventricular tachycardia; and VF, ventricular fibrillation.

A large, randomized, placebo-controlled prehospital trial found amiodarone and lidocaine each improved survival to hospital admission over placebo, however, there was no difference in survival to hospital discharge.⁵ Secondary analyses of both drugs found improved survival to hospital

discharge in bystander-witnessed arrest, with a significant interaction between active drug effect and witnessed arrest status, as did amiodarone in EMS-witnessed arrest. When administered within 8 minutes of ALS-capable EMS arrival, amiodarone also improved survival to hospital admission, discharge, and functional survival at hospital discharge.⁶ This suggests the potential for a small, time-dependent therapeutic window in patients with rapidly recognized and treated cardiac arrest. Data are insufficient to definitively distinguish between the effectiveness of lidocaine and amiodarone, nor their benefit when given in combination.

2. No new evidence emerged from a 2025 ILCOR evidence update⁷ regarding the use of other parenteral antiarrhythmic agents in cardiac arrest. These include bretylium tosylate (which was recently reintroduced in the United States market with no new evidence on its effectiveness or safety); sotalol (which requires a slow infusion and has unknown benefit when given as a bolus in cardiac arrest⁸), procainamide (also requiring slow infusion, and of uncertain benefit when given by rapid infusion as a second-line agent in cardiac arrest⁹), and beta blockers (for which the ILCOR update found insufficient evidence to recommend for or against use).¹⁰ The effectiveness of these drugs administered in combination for cardiac arrest has not been systematically addressed and remains a knowledge gap. Nifekalant is currently unavailable in the United States and is therefore excluded from this recommendation.
3. Both the 2023 AHA focused update and ILCOR's 2023 CoSTR summary identified no new compelling data to alter previous recommendations regarding the use of steroids bundled with a vasopressor agent in cardiac arrest.^{3,11} A large multicenter, blinded, placebo-controlled randomized trial of IHCA deploying this drug combination found an improved rate of ROSC but not survival to hospital discharge or neurological outcome, failing to confirm results of 2 previous single center trials.¹² Observational studies combining intra-arrest corticosteroids with standard resuscitation have demonstrated mixed outcomes.^{13–15} An ongoing registered clinical trial (NCT06203847) may provide further clarity.
4. The 2023 AHA focused update,³ ILCOR's 2023 CoSTR summary,¹¹ and a systematic review including 3 RCTs,¹⁶ found routine calcium administration during cardiac arrest, including cases of refractory asystole and pulseless electrical activity, did not improve survival to hospital discharge or neurological outcome, with a cautionary trend toward potential harm. A Get With the Guidelines registry report also found no evidence of benefit from

calcium administration, independent of rhythm presentation, during IHCA.¹⁷ Although the findings of registry data are confounded by temporal bias (ie, calcium administered as a last resort during prolonged resuscitations, where outcomes are already likely poor), the randomized trial data, designed to mitigate that bias, found no evidence of benefit when calcium was administered immediately after the first dose of epinephrine.¹⁸ Administration of calcium in special circumstances such as hyperkalemia and toxic ingestion is discussed elsewhere as noted above.

5. The 2023 AHA focused update found routine administration of sodium bicarbonate in cardiac arrest was of no benefit.³ However, recent observational data examining prehospital administration of sodium bicarbonate in pulseless electrical activity and asystole suggests a potential association with improved survival.¹⁹ As observed with calcium, issues of resuscitation time bias may also apply to bicarbonate, resulting in difficulty interpreting the findings.^{20,21} Although contemporary evidence is insufficient to change the recommendation, there may be clinical equipoise to warrant further clinical trials to assess its utility in nonshockable presenting rhythms. Use of sodium bicarbonate in special circumstances such as hyperkalemia is discussed elsewhere as noted above.
6. The 2024 ILCOR CoSTR found intra-arrest magnesium administration did not improve ROSC, survival, or neurological outcome regardless of the presenting cardiac arrest rhythm,^{22–25} nor was it useful for monomorphic VT.²⁶ Magnesium's efficacy in the treatment of torsade de pointes is addressed later in this Part (see Treatment of Adults With Polymorphic Ventricular Tachycardia).

REFERENCES

1. Panchal AR, Bartos JA, Cabanas JG, Donnino MW, Drennan IR, Hirsch KG, Kudenchuk PJ, Kurz MC, Lavonas EJ, Morley PT, et al; Adult Basic and Advanced Life Support Writing Group. Part 3: Adult Basic and Advanced Life Support: 2020 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2020;142:S366–S468. doi: 10.1161/CIR.0000000000000916
2. Cao D, Arens AM, Chow SL, Easter SR, Hoffman RS, Lagina AT, Lavonas EJ, Patil KD, Sutherland LD, Tijssen JA, et al. Part 10: Adult and pediatric special circumstances of resuscitation: 2025 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2025;152(suppl 2):S578–S672. doi: 10.1161/CIR.0000000000001378
3. Perman SM, Elmer J, Maciel CB, Uzendu A, May T, Mumma BE, Bartos JA, Rodriguez AJ, Kurz MC, Panchal AR, et al; American Heart Association. 2023 American Heart Association Focused Update on Adult Advanced Cardiovascular Life Support: An Update to the American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2024;149:e254–e273. doi: 10.1161/CIR.0000000000001194
4. Greif R, Bray JE, Djarv T, Drennan IR, Liley HG, Ng KC, Cheng A, Douma MJ, Scholfield BR, Smyth M, et al. 2024 International Consensus on

Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations: Summary From the Basic Life Support; Advanced Life Support; Pediatric Life Support; Neonatal Life Support; Education, Implementation, and Teams; and First Aid Task Forces. *Circulation*. 2024;150:e580–e687. doi: 10.1161/CIR.0000000000001288

5. Kudenchuk PJ, Brown SP, Daya M, Rea T, Nichol G, Morrison LJ, Leroux B, Vaillancourt C, Wittwer L, Callaway CW, et al; Resuscitation Outcomes Consortium Investigators. Amiodarone, Lidocaine, or Placebo in Out-of-Hospital Cardiac Arrest. *N Engl J Med*. 2016;374:1711–1722. doi: 10.1056/NEJMoa1514204

6. Lupton JR, Neth MR, Sahni R, Rui J, Wittwer L, Newgard CD, Daya MR. Survival by time-to-administration of amiodarone, lidocaine, or placebo in shock-refractory out-of-hospital cardiac arrest. *Acad Emerg Med*. 2023;30:906–917. doi: 10.1111/acem.14716

7. Drennan IR, Berg KM, Böttiger BW, Chia YW, Couper K, Crowley C, D'Arrigo S, Deakin CD, Fernando SM, Garg R, et al. 2025 International Liaison Committee on Resuscitation Consensus on Science With Treatment Recommendations: Advanced Life Support. *Circulation*. 2025;152(suppl 1):S72–S115. doi: 10.1161/CIR.0000000000001360

8. Kovoor P, Love A, Hall J, Krut R, Sadick N, Ho D, Adelstein BA, Ross DL. Randomized double-blind trial of sotalol versus lignocaine in out-of-hospital refractory cardiac arrest due to ventricular tachyarrhythmia. *Intern Med J*. 2005;35:518–525. doi: 10.1111/j.1445-5994.2005.00896.x

9. Markel DT, Gold LS, Allen J, Fahrenbruch CE, Rea TD, Eisenberg MS, Kudenchuk PJ. Procainamide and survival in ventricular fibrillation out-of-hospital cardiac arrest. *Acad Emerg Med*. 2010;17:617–623. doi: 10.1111/j.1553-2712.2010.00763.x

10. Gottlieb M, Dyer S, Peksa GD. Beta-blockade for the treatment of cardiac arrest due to ventricular fibrillation or pulseless ventricular tachycardia: A systematic review and meta-analysis. *Resuscitation*. 2020;146:118–125. doi: 10.1016/j.resuscitation.2019.11.019

11. Berg KM, Bray JE, Ng KC, Liley HG, Greif R, Carlson JN, Morley PT, Drennan IR, Smyth M, Scholefield BR, et al; Collaborators. 2023 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations: Summary From the Basic Life Support; Advanced Life Support; Pediatric Life Support; Neonatal Life Support; Education, Implementation, and Teams; and First Aid Task Forces. *Circulation*. 2023;148:e187–e280. doi: 10.1161/CIR.0000000000001179

12. Andersen LW, Isbye D, Kjaergaard J, Kristensen CM, Darling S, Zwisler ST, Fisker S, Schmidt JC, Kirkegaard H, Grejs AM, et al. Effect of Vasopressin and Methylprednisolone vs Placebo on Return of Spontaneous Circulation in Patients With In-Hospital Cardiac Arrest: A Randomized Clinical Trial. *JAMA*. 2021;326:1586–1594. doi: 10.1001/jama.2021.16628

13. Tsai MS, Huang CH, Chang WT, Chen WJ, Hsu CY, Hsieh CC, Yang CW, Chiang WC, Ma MH, Chen SC. The effect of hydrocortisone on the outcome of out-of-hospital cardiac arrest patients: a pilot study. *Am J Emerg Med*. 2007;25:318–325. doi: 10.1016/j.ajem.2006.12.007

14. Tsai MS, Chuang PY, Yu PH, Huang CH, Tang CH, Chang WT, Chen WJ. Glucocorticoid use during cardiopulmonary resuscitation may be beneficial for cardiac arrest. *Int J Cardiol*. 2016;222:629–635. doi: 10.1016/j.ijcard.2016.08.017

15. Andersen LW, Holmberg MJ, Hoybye M, Isbye D, Kjaergaard J, Darling S, Zwisler ST, Larsen JM, Rasmussen BS, Iversen K, et al. Vasopressin and methylprednisolone and hemodynamics after in-hospital cardiac arrest - A post hoc analysis of the VAM-IHCA trial. *Resuscitation*. 2023;191:109922. doi: 10.1016/j.resuscitation.2023.109922

16. Hsu CH, Couper K, Nix T, Drennan I, Reynolds J, Kleinman M, Berg KM; Advanced Life Support and Paediatric Life Support Task Forces at the International Liaison Committee on Resuscitation (ILCOR) Paediatric Life Support Task Forces at the International Liaison Committee on R. Calcium during cardiac arrest: A systematic review. *Resusc Plus*. 2023;14:100379. doi: 10.1016/j.resplu.2023.100379

17. Holmberg MJ, Granfeldt A, Andersen LW. Bicarbonate, calcium, and magnesium for in-hospital cardiac arrest - An instrumental variable analysis. *Resuscitation*. 2023;191:109958. doi: 10.1016/j.resuscitation.2023.109958

18. Vallentin MF, Granfeldt A, Meilandt C, Povlsen AL, Sindberg B, Holmberg MJ, Iversen BN, Maerkedahl R, Mortensen LR, Nyboe R, et al. Effect of Intravenous or Introsseous Calcium vs Saline on Return of Spontaneous Circulation in Adults With Out-of-Hospital Cardiac Arrest: A Randomized Clinical Trial. *JAMA*. 2021;326:2268–2276. doi: 10.1001/jama.2021.20929

19. Niederberger SM, Crowe RP, Salcido DD, Menegazzi JJ. Sodium bicarbonate administration is associated with improved survival in asystolic and

PEA Out-of-Hospital cardiac arrest. *Resuscitation*. 2023;182:109641. doi: 10.1016/j.resuscitation.2022.11.007

20. Kawano T, Grunau B, Scheuermeyer FX, Gibo K, Dick W, Fordyce CB, Dorian P, Stenstrom R, Straight R, Christenson J. Prehospital sodium bicarbonate use could worsen long term survival with favorable neurological recovery among patients with out-of-hospital cardiac arrest. *Resuscitation*. 2017;119:63–69. doi: 10.1016/j.resuscitation.2017.08.008

21. Ahn S, Kim YJ, Sohn CH, Seo DW, Lim KS, Donnino MW, Kim WY. Sodium bicarbonate on severe metabolic acidosis during prolonged cardiopulmonary resuscitation: a double-blind, randomized, placebo-controlled pilot study. *J Thorac Dis*. 2018;10:2295–2302. doi: 10.21037/jtd.2018.03.124

22. Thel MC, Armstrong AL, McNulty SE, Califf RM, O'Connor CM. Randomised trial of magnesium in in-hospital cardiac arrest. *Duke Internal Medicine Housestaff*. *Lancet*. 1997;350:1272–1276. doi: 10.1016/s0140-6736(97)05048-4

23. Hassan TB, Jagger C, Barnett DB. A randomised trial to investigate the efficacy of magnesium sulphate for refractory ventricular fibrillation. *Emerg Med J*. 2002;19:57–62. doi: 10.1136/emj.19.1.57

24. Fatovich DM, Prentice DA, Dobb GJ. Magnesium in cardiac arrest (the magic trial). *Resuscitation*. 1997;35:237–241. doi: 10.1016/s0300-9572(97)00062-2

25. Allegra J, Lavery R, Cody R, Birnbaum G, Brennan J, Hartman A, Horowitz M, Nashed A, Yablonski M. Magnesium sulfate in the treatment of refractory ventricular fibrillation in the prehospital setting. *Resuscitation*. 2001;49:245–249. doi: 10.1016/s0300-9572(00)00375-0

26. Manz M, Jung W, Luderitz B. [Effect of magnesium on sustained ventricular tachycardia]. *Herz*. 1997;22:51–55. doi: 10.1007/BF03042655

ADJUNCTS FOR PERFORMANCE OF CPR

Recommendations for Adjuncts to CPR		
COR	LOE	Recommendations
2b	B-NR	1. An abrupt increase in end-tidal CO ₂ may be used to detect ROSC during chest compressions in adults in cardiac arrest or when a rhythm check reveals an organized rhythm.
2b	C-LD	2. Use of point-of-care ultrasonography during adult cardiac arrest resuscitation to diagnose reversible causes is not well established.
2b	C-LD	3. Use of point-of-care echocardiography to assess cardiac function for prognostication during adult cardiac arrest resuscitation is not well established.
2b	C-LD	4. When supplemental oxygen is available, it may be reasonable to use the maximal feasible inspired oxygen concentration during CPR for adults with cardiac arrest.
2b	C-LD	5. Measurement of arterial blood gases during CPR has uncertain benefit in adults with cardiac arrest.
2b	C-LD	6. In adult patients with cardiac arrest, it may be reasonable to use physiological parameters such as arterial blood pressure or end-tidal CO ₂ when feasible to monitor and optimize CPR quality.
2b	C-EO	7. Arterial pressure monitoring by arterial line may be used to detect ROSC during chest compressions or when a rhythm check reveals an organized rhythm in adults with cardiac arrest.
3: No Benefit	C-LD	8. Head-up cardiopulmonary resuscitation in adults with cardiac arrest is not recommended except in setting of clinical trials.

Synopsis

The foundation of cardiac arrest resuscitation focuses on high-quality CPR, early defibrillation when indicated, and guideline-directed pharmacologic therapies and airway

management strategies. With appropriate expertise and resources, adjuncts to standard cardiac arrest resuscitation provide additional data to support and enhance care delivery. Adjuncts—such as POCUS, ETCO_2 monitoring, blood gas analysis, or arterial lines—may enable the incorporation of precise physiologic parameters to optimize individual resuscitation.

Head-up CPR seeks to augment conventional supine CPR to increase cerebral perfusion pressure with the goal of improving neurologically favorable outcome. Head-up CPR has been studied as a bundle of care combining mechanical chest compressions, an impedance threshold device, and application of an automated device to control sequential elevation of the head and thorax during compressions.

While emerging but limited data exist for many of these CPR adjuncts, more research is needed to inform future guidelines.

Recommendation-Specific Supportive Text

1. ETCO_2 values during CPR serve as a surrogate for cardiac output.^{1,2} A secondary analysis of the Pragmatic Airway Resuscitation Trial found that temporal increases in ETCO_2 were associated with ROSC.³ A systematic review of 8 observational studies found that an abrupt increase in ETCO_2 was a predictor of ROSC; however, no absolute cutoff value was identified.⁴ Data suggests that a sudden increase greater than 10 mm Hg may indicate ROSC,^{5,6} although ROSC can still occur with ETCO_2 increases of less than 10 mm Hg.⁷ These findings may be influenced by other factors known to impact ETCO_2 , including minute ventilation, CPR quality, the administration of epinephrine or sodium bicarbonate, airway management strategies, and the etiology of cardiac arrest.^{8,9}
2. Multiple observational studies report the use of POCUS during cardiac arrest to aid in the identification of potentially reversible causes of cardiac arrest, such as pulmonary embolism, cardiac tamponade, effusion, myocardial infarction, aortic dissection, and hypovolemia. However, evidence for diagnosing these target conditions is limited by inconsistent clinical protocols, sonographic measures, and operator experience, resulting in a high degree of heterogeneity among the studies with a high concern for bias.¹⁰ Furthermore, the use of POCUS has been associated with longer interruptions in chest compressions, longer duration of resuscitation, and higher rates of interventions.^{11,12} Therefore, even when a user with adequate skill and expertise uses POCUS during cardiac arrest, attention to minimizing interruptions in CPR is a priority.^{10,13–16}

3. POCUS has been used to assess cardiac activity and, in the absence of cardiac motion in conjunction with other modalities, to terminate resuscitative efforts.^{14,17–20} However, only a small number of observational studies, with significant limitations, including variability in the definitions of sonographic findings, describe its use in this setting.^{21–23} A single small RCT, multiple systematic reviews, and a meta-analysis of observational studies have consistently found no improvement in outcomes with the use of POCUS during cardiac arrest.^{23–25} Future research that standardizes image acquisition and other study methodology to examine the utility of POCUS for this indication are required. An alternative modality, transesophageal echocardiography, has the potential to provide high-quality imaging of cardiac structure and function without interrupting CPR. However, the utility of transesophageal echocardiography during CPR remains a knowledge gap.
4. No adult human trials directly compare levels of inspired oxygen concentration during CPR. Retrospective studies have found that higher intra-arrest partial pressure of oxygen in the alveoli is associated with survival to hospital admission,²⁶ survival to hospital discharge,²⁷ and survival with favorable neurological outcome.²⁸ However, these findings may be influenced by patient selection, airway management strategies, and the quality of resuscitation.
5. Small prospective studies with significant limitations have found arterial blood gas parameters, such as partial pressure of oxygen and partial pressure of carbon dioxide, to be predictive of ROSC.^{29,30} However, both partial pressure of oxygen and partial pressure of carbon dioxide are dependent on cardiac output and can be influenced by patient factors and the quality of CPR. Additionally, numerous retrospective studies have assessed intra-arrest blood gas analysis as a predictive tool for outcomes but suffer from similar limitations, compounded by their retrospective design.^{26–28,31–38}
6. A retrospective study from the AHA's Get With the Guidelines-Resuscitation found the use of ETCO_2 or diastolic blood pressure monitoring during CPR was associated with increased rate of ROSC.³⁹ ETCO_2 values reflect pulmonary circulation and cardiac output^{1,2} and are positively correlated with increased compression depth^{40–42} and release velocity.^{40,41} Multiple systematic reviews have found numeric ETCO_2 measures to be a predictor of ROSC.^{4,43} An ETCO_2 less than 10 mm Hg is generally associated with poor outcomes, whereas values above 10 mm Hg, and

ideally above 20 mm Hg, are associated with increased rates of ROSC.⁴³ The combination of the association of higher ETCO_2 with ROSC and the findings that CPR quality can increase ETCO_2 suggests that targeting compressions to a value of at least 10 mm Hg, and ideally 20 mm Hg or greater, may indicate mechanically adequate technique. Other factors known to impact ETCO_2 values need to be considered, including minute ventilation, drug administration, airway management strategies, and cardiac arrest etiology. ETCO_2 monitoring is most reliable when measured through an endotracheal tube, but it can also be used with a supraglottic airway or bag-mask ventilation with unclear utility. When an invasive arterial line is in place, arterial diastolic pressure can approximate coronary perfusion pressure and myocardial perfusion during CPR.⁴⁴ The use of diastolic blood pressure monitoring during cardiac arrest is associated with higher ROSC, but there are inadequate adult data to suggest any specific pressure in adults.

7. If an arterial line is in place, the development of an arterial waveform or a sudden increase in diastolic pressure during a rhythm check showing an organized rhythm may indicate ROSC.⁴⁵ When a waveform is present, it is important to ensure it correlates with a palpable pulse to verify ROSC. The placement of an arterial line requires appropriate expertise and resources; however, placement during cardiac arrest may detract from the provision of high-quality CPR.
8. Original work evaluating head-up CPR was done in porcine models, which demonstrated conflicting evidence of efficacy.^{46,47} A recent ILCOR systematic review identified no randomized controlled trials and only 3 observational studies, each with significant methodological limitations.^{48–50} With these and other challenges the systematic review identified, for the outcome of survival to discharge and survival to discharge with favorable neurological outcome, evidence on Head up CPR was considered very low—certainty of evidence downgraded for serious risk of bias. Taken together, current evidence regarding head-up CPR is limited in the absence of RCTs or appropriate comparisons. Furthermore, implementation of this approach requires specialized equipment (automated positioning device, mechanical CPR device, an impedance threshold device) and significant training. Based on this, there is currently insufficient evidence for its use outside of well-designed clinical trials, although future work is needed to evaluate this adjunct.

REFERENCES

1. Weil MH, Bisera J, Trevino RP, Rackow EC. Cardiac output and end-tidal carbon dioxide. *Crit Care Med*. 1985;13:907–909. doi: 10.1097/00003246-198511000-00011
2. Ornato JP, Garnett AR, Glauser FL. Relationship between cardiac output and the end-tidal carbon dioxide tension. *Ann Emerg Med*. 1990;19:1104–1106. doi: 10.1016/s0196-0644(05)81512-4
3. Nassal MMJ, Elola A, Aramendi E, Jaureguibertia X, Powell JR, Idris A, Raya Krishnamoorthy BP, Daya MR, Aufderheide TP, Carlson JN, et al. Temporal Trends in End-Tidal Capnography and Outcomes in Out-of-Hospital Cardiac Arrest: A Secondary Analysis of a Randomized Clinical Trial. *JAMA Netw Open*. 2024;7:e2419274. doi: 10.1001/jamanetworkopen.2024.19274
4. Caro-Alonso PA, Rodriguez-Martin B. [The end-tidal carbon dioxide as an early sign and predictor of the return of spontaneous circulation during out-of-hospital cardiac arrest. A systematic review]. *Rev Esp Salud Publica*. 2021;95:e202104068.
5. Pokorna M, Necas E, Kratochvil J, Skripsky R, Andrik M, Franek O. A sudden increase in partial pressure end-tidal carbon dioxide (P(ET)CO(2)) at the moment of return of spontaneous circulation. *J Emerg Med*. 2010;38:614–621. doi: 10.1016/j.jemermed.2009.04.064
6. Lui CT, Poon KM, Tsui KL. Abrupt rise of end tidal carbon dioxide level was a specific but non-sensitive marker of return of spontaneous circulation in patient with out-of-hospital cardiac arrest. *Resuscitation*. 2016;104:53–58. doi: 10.1016/j.resuscitation.2016.04.018
7. Grabman B, Bulger NE, Harrington BM, Walker RG, Latimer AJ, Snyder BD, Sayre MR, Maynard C, Johnson NJ, Van Dyke M, et al. Increase in end-tidal carbon dioxide after defibrillation predicts sustained return of spontaneous circulation during out-of-hospital cardiac arrest. *Resuscitation*. 2022;181:48–54. doi: 10.1016/j.resuscitation.2022.10.001
8. Sandroni C, De Santis P, D'Arigo S. Capnography during cardiac arrest. *Resuscitation*. 2018;132:73–77. doi: 10.1016/j.resuscitation.2018.08.018
9. Pantazopoulos C, Xanthos T, Pantazopoulos I, Papalois A, Kouskouni E, Iacovidou N. A Review of Carbon Dioxide Monitoring During Adult Cardiopulmonary Resuscitation. *Heart Lung Circ*. 2015;24:1053–1061. doi: 10.1016/j.hlc.2015.05.013
10. Reynolds JC, Nicholson T, O'Neil B, Drennan IR, Issa M, Welsford M; Advanced Life Support Task Force at the International Liaison Committee on Resuscitation ILCOR. Diagnostic test accuracy of point-of-care ultrasound during cardiopulmonary resuscitation to indicate the etiology of cardiac arrest: A systematic review. *Resuscitation*. 2022;172:54–63. doi: 10.1016/j.resuscitation.2022.01.006
11. Atkinson PR, Beckett N, French J, Banerjee A, Fraser J, Lewis D. Does Point-of-care Ultrasound Use Impact Resuscitation Length, Rates of Intervention, and Clinical Outcomes During Cardiac Arrest? A Study from the Sonography in Hypotension and Cardiac Arrest in the Emergency Department (SHoC-ED) Investigators. *Cureus*. 2019;11:e4456. doi: 10.7759/cureus.4456
12. Clattenburg EJ, Wroe P, Brown S, Gardner K, Losonczy L, Singh A, Nagdev A. Point-of-care ultrasound use in patients with cardiac arrest is associated prolonged cardiopulmonary resuscitation pauses: A prospective cohort study. *Resuscitation*. 2018;122:65–68. doi: 10.1016/j.resuscitation.2017.11.056
13. Zanatta M, Lorenzi C, Scorpiniti M, Cianci V, Pasini R, Barchitta A. Ultrasound-Guided Chest Compressions in Out-of-Hospital Cardiac Arrests. *J Emerg Med*. 2020;59:e225–e233. doi: 10.1016/j.jemermed.2020.07.005
14. Zaki HA, Ifthikhar H, Shaban EE, Najam M, Alkahlout BH, Shallik N, Elnabawy W, Basharat K, Azad AM. The role of point-of-care ultrasound (POCUS) imaging in clinical outcomes during cardiac arrest: a systematic review. *Ultrasound J*. 2024;16:4. doi: 10.1186/s13089-023-00346-1
15. Gaspari R, Weekes A, Adhikari S, Noble VE, Nomura JT, Theodoro D, Woo M, Atkinson P, Blehar D, Brown SM, et al. Emergency department point-of-care ultrasound in out-of-hospital and in-ED cardiac arrest. *Resuscitation*. 2016;109:33–39. doi: 10.1016/j.resuscitation.2016.09.018
16. Breitkreutz R, Price S, Steiger HV, Seeger FH, Ilper H, Ackermann H, Rudolph M, Uddin S, Weigand MA, Muller E, et al; Emergency Ultrasound Working Group of the Johann Wolfgang Goethe-University Hospital, Frankfurt am Main. Focused echocardiographic evaluation in life support and peri-resuscitation of emergency patients: a prospective trial. *Resuscitation*. 2010;81:1527–1533. doi: 10.1016/j.resuscitation.2010.07.013
17. Wu C, Zheng Z, Jiang L, Gao Y, Xu J, Jin X, Chen Q, Zhang M. The predictive value of bedside ultrasound to restore spontaneous circulation in patients

- with pulseless electrical activity: A systematic review and meta-analysis. *PLoS One*. 2018;13:e0191636. doi: 10.1371/journal.pone.0191636
18. Kedan I, Ciozda W, Palatinus JA, Palatinus HN, Kimchi A. Prognostic value of point-of-care ultrasound during cardiac arrest: a systematic review. *Cardiovasc Ultrasound*. 2020;18:1. doi: 10.1186/s12947-020-0185-8
 19. Albaroudi O, Albaroudi B, Haddad M, Abdle-Rahman ME, Kumar TSS, Jarman RD, Harris T. Can absence of cardiac activity on point-of-care echocardiography predict death in out-of-hospital cardiac arrest? A systematic review and meta-analysis. *Ultrasound J*. 2024;16:10. doi: 10.1186/s13089-024-00360-x
 20. Tsou PY, Kurbedin J, Chen YS, Chou EH, Lee MG, Lee MC, Ma MH, Chen SC, Lee CC. Accuracy of point-of-care focused echocardiography in predicting outcome of resuscitation in cardiac arrest patients: A systematic review and meta-analysis. *Resuscitation*. 2017;114:92–99. doi: 10.1016/j.resuscitation.2017.02.021
 21. Reynolds JC, Issa MS, T CN, Drennan IR, Berg KM, O'Neil BJ, Welsford M; Advanced Life Support Task Force of the International Liaison Committee on R. Prognostication with point-of-care echocardiography during cardiac arrest: A systematic review. *Resuscitation*. 2020;152:56–68. doi: 10.1016/j.resuscitation.2020.05.004
 22. Lalande E, Burwash-Brennan T, Burns K, Atkinson P, Lambert M, Jarman B, Lamprecht H, Banerjee A, Woo MY; Investigators SH. Is point-of-care ultrasound a reliable predictor of outcome during atraumatic, non-shockable cardiac arrest? A systematic review and meta-analysis from the SHoC investigators. *Resuscitation*. 2019;139:159–166. doi: 10.1016/j.resuscitation.2019.03.027
 23. Dudek M, Szarpak L, Peacock FW, Gasecka A, Michalski T, Wroblewski P, Kaminska H, Borkowska G, Skrzypek E, Smereka A, et al. Diagnostic performance of point-of-use ultrasound of resuscitation outcomes: A systematic review and meta-analysis of 3265 patients. *Cardiol J*. 2023;30:237–246. doi: 10.5603/CJ.a2021.0044
 24. Chardoli M, Heidari F, Rabiee H, Sharif-Alhoseini M, Shokoohi H, Rahimi-Movaghar V. Echocardiography integrated ACLS protocol versus conventional cardiopulmonary resuscitation in patients with pulseless electrical activity cardiac arrest. *Chin J Traumatol*. 2012;15:284–287.
 25. Thandar S, Sahu AK, Sinha TP, Bhoi S. Role of initial cardiac activity assessed by point-of-care ultrasonography in predicting cardiac arrest outcomes: A prospective cohort study. *Turk J Emerg Med*. 2023;23:24–29. doi: 10.4103/2452-2473.366482
 26. Spindelboeck W, Schindler O, Moser A, Hausler F, Wallner S, Strasser C, Haas J, Gemes G, Prause A. Increasing arterial oxygen partial pressure during cardiopulmonary resuscitation is associated with improved rates of hospital admission. *Resuscitation*. 2013;84:770–775. doi: 10.1016/j.resuscitation.2013.01.012
 27. Patel JK, Schoenfeld E, Parikh PB, Parnia S. Association of Arterial Oxygen Tension During In-Hospital Cardiac Arrest With Return of Spontaneous Circulation and Survival. *J Intensive Care Med*. 2018;33:407–414. doi: 10.1177/0885066616658420
 28. Izawa J, Komukai S, Nishioka N, Kiguchi T, Kitamura T, Iwami T. Outcomes associated with intra-arrest hyperoxaemia in out-of-hospital cardiac arrest: A registry-based cohort study. *Resuscitation*. 2022;181:173–181. doi: 10.1016/j.resuscitation.2022.11.008
 29. Calbay A, Cakir Z, Bayramoglu A. Prognostic value of blood gas parameters and end-tidal carbon dioxide values in out-of-hospital cardiopulmonary arrest patients. *Turk J Med Sci*. 2019;49:1298–1302. doi: 10.3906/sag-1812-156
 30. Kim YJ, Lee YJ, Ryoo SM, Sohn CH, Ahn S, Seo DW, Lim KS, Kim WY. Role of blood gas analysis during cardiopulmonary resuscitation in out-of-hospital cardiac arrest patients. *Medicine (Baltimore)*. 2016;95:e3960. doi: 10.1097/MD.00000000000003960
 31. Wang CH, Wu MC, Wu CY, Huang CH, Tsai MS, Lu TC, Chou E, Wu YW, Chang WT, Chen WJ. Blood gas phenotyping and tracheal intubation timing in adult in-hospital cardiac arrest: a retrospective cohort study. *Sci Rep*. 2021;11:10480. doi: 10.1038/s41598-021-89920-y
 32. Wang CH, Chang WT, Huang CH, Tsai MS, Yu PH, Wu YW, Chen WJ. Associations between early intra-arrest blood acidemia and outcomes of adult in-hospital cardiac arrest: A retrospective cohort study. *J Formos Med Assoc*. 2020;119:644–651. doi: 10.1016/j.jfma.2019.08.020
 33. Shin J, Lim YS, Kim K, Lee HJ, Lee SJ, Jung E, You KM, Yang HJ, Kim JJ, Kim J, et al. Initial blood pH during cardiopulmonary resuscitation in out-of-hospital cardiac arrest patients: a multicenter observational registry-based study. *Crit Care*. 2017;21:322. doi: 10.1186/s13054-017-1893-9
 34. Nalbant E, A M, Çelik A, Yavaş O, Ersunan G, Bilir O, Altuntaş G. Early Predictors of Return of Spontaneous Circulation in Patients Undergoing Cardiopulmonary Resuscitation. *J Exp Clin Med*. 2022;39:1157–1162.
 35. Matsuyama T, Ohta B, Kiyohara K, Kitamura T. Intra-arrest partial carbon dioxide level and favorable neurological outcome after out-of-hospital cardiac arrest: a nationwide multicenter observational study in Japan (the JAAM-OHCA registry). *Eur Heart J Acute Cardiovasc Care*. 2023;12:14–21. doi: 10.1093/ehjacc/zuac152
 36. Hong SI, Kim JS, Kim YJ, Kim WY. Dynamic changes in arterial blood gas during cardiopulmonary resuscitation in out-of-hospital cardiac arrest. *Sci Rep*. 2021;11:23165. doi: 10.1038/s41598-021-02764-4
 37. Farzi S, H F, Wallner S, Spindelböck W, Prause G, Gemes G. Prehospital measurement of arterial base excess and its role as a possible predictor of outcome after out-of-hospital cardiac arrest. *Emergencias*. 2013;25:47–50.
 38. Lin CH, Yu S, Chen CY, Huang FW, Chen WK, Shih HM. Early blood pH as an independent predictor of neurological outcome in patients with out-of-hospital cardiac arrest: A retrospective observational study. *Medicine (Baltimore)*. 2021;100:25724. doi: 10.1097/MD.00000000000025724
 39. Sutton RM, French B, Meaney PA, Topjian AA, Parshuram CS, Edelson DP, Schexnayder S, Abella BS, Merchant RM, Bembea M, et al; American Heart Association's Get With The Guidelines–Resuscitation Investigators. Physiologic monitoring of CPR quality during adult cardiac arrest: A propensity-matched cohort study. *Resuscitation*. 2016;106:76–82. doi: 10.1016/j.resuscitation.2016.06.018
 40. Murphy RA, Bobrow BJ, Spaite DW, Hu C, McDannold R, Vadeboncoeur TF. Association between Prehospital CPR Quality and End-Tidal Carbon Dioxide Levels in Out-of-Hospital Cardiac Arrest. *Prehosp Emerg Care*. 2016;20:369–377. doi: 10.3109/109031272015.1115929
 41. De Roos E, Vanwulpen M, Hachimi-Idrissi S. Chest compression release velocity: An independent determinant of end-tidal carbon dioxide in out-of-hospital cardiac arrest. *Am J Emerg Med*. 2022;54:71–75. doi: 10.1016/j.ajem.2022.01.053
 42. Sheak KR, Wiebe DJ, Leary M, Babaeizadeh S, Yuen TC, Zive D, Owens PC, Edelson DP, Daya MR, Idris AH, et al. Quantitative relationship between end-tidal carbon dioxide and CPR quality during both in-hospital and out-of-hospital cardiac arrest. *Resuscitation*. 2015;89:149–154. doi: 10.1016/j.resuscitation.2015.01.026
 43. Paiva EF, Paxton JH, O'Neil BJ. The use of end-tidal carbon dioxide (ETCO₂) measurement to guide management of cardiac arrest: A systematic review. *Resuscitation*. 2018;123:1–7. doi: 10.1016/j.resuscitation.2017.12.003
 44. Rivers EP, Lozon J, Enriquez E, Havstad JV, Martin GB, Lewandowski CA, Goetting MG, Rosenberg JA, Paradis NA, Nowak RM. Simultaneous radial, femoral, and aortic arterial pressures during human cardiopulmonary resuscitation. *Crit Care Med*. 1993;21:878–883. doi: 10.1097/00003246-199306000-00016
 45. Paradis NA, Martin GB, Rivers EP, Goetting MG, Appleton TJ, Feingold M, Nowak RM. Coronary perfusion pressure and the return of spontaneous circulation in human cardiopulmonary resuscitation. *JAMA*. 1990;263:1106–1113.
 46. Varney J, Motawea KR, Mostafa MR, AbdelQadir YH, Abouleenein M, Kandil OA, Ibrahim N, Hashim HT, Murry K, Jackson G, et al. Efficacy of head-up CPR compared to supine CPR positions: Systematic review and meta-analysis. *Health Sci Rep*. 2022;5:e644. doi: 10.1002/hsr2.644
 47. Jaeger D, Kosmopoulos M, Voicu S, Kalra R, Gaisendrees C, Schlartenberger G, Bartos JA, Yannopoulos D. Cerebral hemodynamic effects of head-up CPR in a porcine model. *Resuscitation*. 2023;193:110039. doi: 10.1016/j.resuscitation.2023.110039
 48. Pepe PE, Schepke KA, Antevy PM, Crowe RP, Millstone D, Coyle C, Prusansky C, Garay S, Ellis R, Fowler RL, et al. Confirming the Clinical Safety and Feasibility of a Bundled Methodology to Improve Cardiopulmonary Resuscitation Involving a Head-Up/Torso-Up Chest Compression Technique. *Crit Care Med*. 2019;47:449–455. doi: 10.1097/CCM.0000000000003608
 49. Moore JC, Pepe PE, Schepke KA, Lick C, Duval S, Holley J, Salverda B, Jacobs M, Nystrom P, Quinn R, et al. Head and thorax elevation during cardiopulmonary resuscitation using circulatory adjuncts is associated with improved survival. *Resuscitation*. 2022;179:9–17. doi: 10.1016/j.resuscitation.2022.07.039
 50. Bachista KM, Moore JC, Labarere J, Crowe RP, Emanuelson LD, Lick CJ, Debaty GP, Holley JE, Quinn RP, Schepke KA, et al. Survival for Non-shockable Cardiac Arrests Treated With Noninvasive Circulatory Adjuncts and Head/Thorax Elevation. *Crit Care Med*. 2024;52:170–181. doi: 10.1097/CCM.0000000000000605

TERMINATION OF RESUSCITATIVE MEASURES

Recommendations for Termination of Resuscitation		
COR	LOE	Recommendations
1	B-NR	1. When TOR is being considered, BLS prehospital professionals should use the BLS TOR rule for adult patients in OHCA, where ALS is not available or may be significantly delayed.
2a	B-NR	2. It is reasonable for ALS prehospital professionals to use the ALS TOR rule to terminate resuscitation efforts in the field for adult patients with OHCA.
2a	B-NR	3. In a tiered EMS system with both ALS and BLS professionals, it is reasonable to use the Universal TOR rule for adult patients with OHCA.
2b	C-LD	4. In intubated adult patients, failure to achieve an end-tidal CO ₂ of greater than 10 mm Hg by waveform capnography after 20 minutes of ALS resuscitation may be considered as a component of a multimodal approach to decide when to end resuscitative efforts.
3: Harm	C-EO	5. In nonintubated adult patients, a specific end-tidal CO ₂ cutoff value at any time during CPR should not be used as an indication to end resuscitative efforts.

Synopsis

OHCA is a resource-intensive condition associated with low rates of survival. It is important for EMS professionals to differentiate patients in whom continued resuscitation is futile from patients with a chance of survival who should receive continued resuscitation and transportation to hospital. This will aid in both resource use and optimizing a patient's chance for survival. Using a validated TOR rule will help ensure accuracy in determining futility in cardiac arrest patients. Though many TOR rules have been proposed,³ TOR rules that have been highly studied in North America include the BLS, ALS, and UTOR rules (Figures 3 and 4).

TOR rules are applicable for the specific scope of practice of the EMS agency upon which they were validated. For example, an EMS agency with only BLS-trained professionals (first responders or emergency medical technicians [EMTs]) the BLS TOR rule is appropriate;




Figure 3. BLS/Universal Termination of Resuscitation Rules. AED indicates automated external defibrillator; and BLS, basic life support.




Figure 4. ALS Termination of Resuscitation Rule. ALS indicates advanced life support; and CPR, cardiopulmonary resuscitation.

for an EMS agency with only ALS-trained professionals, the ALS TOR rule is appropriate; and for an agency or system with both BLS- and ALS-trained professionals (commonly referred to as a tiered-response system) the UTOR rule is appropriate.

These TOR rules are inappropriate for populations in which they have not been validated, such as for patients with overdose, trauma, or IHCA. Further research is needed to evaluate the appropriate application of existing TOR rules to gain a clearer understanding of the risks and benefits associated with their use.

Recommendation-Specific Supportive Text

1. In a meta-analysis of 7 published BLS TOR studies primarily in North America (33 795 patients pooled from retrospective and prospective observational studies), only 0.13% (95% CI, 0.03%–0.58%) of patients who fulfilled the BLS termination criteria survived to hospital discharge.¹ A 2024 international meta-analysis² of 6 reclassified retrospective validation studies from EMS systems primarily in Asia suggests that the specificity of the BLS rule, when generalized outside of North America may be lower than previously reported. The analysis reclassified EMS agencies, included heterogeneous methodologies and response characteristics, and relied heavily on retrospective data. These limitations highlight the importance of further rigorous, prospective research to validate the specificity and applicability of BLS TOR rules by diverse EMS agencies.
2. In a meta-analysis of 2 published retrospective observational studies (10 178 patients) of the ALS TOR rule, only 0.01% (95% CI, 0.00%–0.07%) of patients who fulfilled the ALS termination criteria survived to hospital discharge.¹ The same 2024 international meta-analysis described above² analyzed 17 published studies on ALS TOR patients, finding a pooled specificity of 96% (95% CI, 0.93–0.99).

3. The UTOR rule, which uses the same criteria as the BLS rule (ie, arrest not witnessed by EMS professionals; no shock delivered; no ROSC), has been prospectively validated in combined BLS/ALS, or tiered response, EMS agencies.³ Although the rule did not have adequate specificity after 6 minutes of resuscitation (false-positive rate: 2.1%) it did achieve better than 99% specificity after approximately 15 minutes of attempted resuscitation while still reducing transportation by half. A retrospective analysis found that the application of the UTOR rule at 20 minutes of resuscitation was able to predict futility, identifying over 99% of survivors and patients with good neurological outcome.⁴ Another meta-analysis² also examined the UTOR rule in 19 studies, many of which had initially been classified as employing the BLS TOR rule in tiered BLS + ALS EMS systems. After reclassification, the pooled specificity of the UTOR rule was reported to be 88% (95% CI, 0.88–0.94). Notably, 10 of the 11 reclassified studies were conducted outside of North America.

4. In intubated patients, an ETCO₂ measurement of less than 10 mm Hg indicates little to no cardiac output. Several small retrospective studies provide evidence showing that an ETCO₂ less than 10 mm Hg after 20 minutes of ALS resuscitation is strongly but not perfectly predictive of futility.^{5–8} These small observational studies suffer from a high risk of bias. Alternative ETCO₂ thresholds, time points, and changes in ETCO₂ over time have been proposed.^{9–12} The use of ETCO₂ alone to predict patient outcome needs to be validated in a large prospective study.² No data is available that addresses ETCO₂ measurement with supraglottic airways and predicts poor outcomes. Additionally, isolated ETCO₂ values greater than 20 mm Hg, at later stages of resuscitation, should not be used in isolation as a potential indicator for continued resuscitation, particularly when other prognostic indicators are not favorable. High-quality studies assessing the role of ETCO₂ for termination of resuscitation in patients on mechanical CPR have not been conducted and are an active area of research.

5. No studies were found that specifically examined the use of ETCO₂ in cardiac arrest patients without an advanced airway. It is not known whether ETCO₂ values during bag-mask ventilation are as reliable as those with an advanced airway in place. There is insufficient evidence to support a universal “cutoff value” of ETCO₂ for decisions about TOR in a non-intubated patient.

REFERENCES

1. Ebell MH, Vellinga A, Masterson S, Yun P. Meta-analysis of the accuracy of termination of resuscitation rules for out-of-hospital cardiac arrest. *Emerg Med J*. 2019;36:479–484. doi: 10.1136/emermed-2018-207833

2. Kwong JL, Drennan IR, Turner L, Cheskes S. Predefibrillation end-tidal CO2 and defibrillation success in out-of-hospital cardiac arrest: an observational cohort study. *Emerg Med J*. 2023;40:48–55. doi: 10.1136/emermed-2021-211951

3. Grunau B, Taylor J, Scheuermeyer FX, Stenstrom R, Dick W, Kawano T, Barbic D, Drennan I, Christenson J. External Validation of the Universal Termination of Resuscitation Rule for Out-of-Hospital Cardiac Arrest in British Columbia. *Ann Emerg Med*. 2017;70:374–381.e1. doi: 10.1016/j.annemergmed.2017.01.030

4. Drennan IR, Case E, Verbeek PR, Reynolds JC, Goldberger ZD, Jasti J, Charleston M, Herren H, Idris AH, Leslie PR, et al; Resuscitation Outcomes Consortium Investigators. A comparison of the universal TOR Guideline to the absence of prehospital ROSC and duration of resuscitation in predicting futility from out-of-hospital cardiac arrest. *Resuscitation*. 2017;111:96–102. doi: 10.1016/j.resuscitation.2016.11.021

5. Wayne MA, Levine RL, Miller CC. Use of end-tidal carbon dioxide to predict outcome in prehospital cardiac arrest. *Ann Emerg Med*. 1995;25:762–767. doi: 10.1016/s0196-0644(95)70204-0

6. Levine RL, Wayne MA, Miller CC. End-tidal carbon dioxide and outcome of out-of-hospital cardiac arrest. *N Engl J Med*. 1997;337:301–306. doi: 10.1056/NEJM199707313370503

7. Akinci E, Ramadan H, Yuzbasioglu Y, Coskun F. Comparison of end-tidal carbon dioxide levels with cardiopulmonary resuscitation success presented to emergency department with cardiopulmonary arrest. *Pak J Med Sci*. 2014;30:16–21. doi: 10.12669/pjms.301.4024

8. Ahrens T, Schallom L, Bettorf K, Ellner S, Hurt G, O'Mara V, Ludwig J, George W, Marino T, Shannon W. End-tidal carbon dioxide measurements as a prognostic indicator of outcome in cardiac arrest. *Am J Crit Care*. 2001;10:391–398.

9. Sun KF, Poon KM, Lui CT, Tsui KL. Clinical prediction rule of termination of resuscitation for out-of-hospital cardiac arrest patient with pre-hospital defibrillation given. *Am J Emerg Med*. 2021;50:733–738. doi: 10.1016/j.ajem.2021.09.058

10. Nassal MMJ, Elola A, Aramendi E, Jaureguibeitia X, Powell JR, Idris A, Raya Krishnamoorthy BP, Daya MR, Aufderheide TP, Carlson JN, et al. Temporal Trends in End-Tidal Capnography and Outcomes in Out-of-Hospital Cardiac Arrest: A Secondary Analysis of a Randomized Clinical Trial. *JAMA Netw Open*. 2024;7:e2419274. doi: 10.1001/jamanetworkopen.2024.19274

11. Crickmer M, Drennan IR, Turner L, Cheskes S. The association between end-tidal CO2 and return of spontaneous circulation after out-of-hospital cardiac arrest with pulseless electrical activity. *Resuscitation*. 2021;167:76–81. doi: 10.1016/j.resuscitation.2021.08.014

12. Cereceda-Sánchez FJ, Ponce-Taylor J, Montero-París P, Unzaga-Ercilla I, Martínez-Cuellar N, Molina-Mula J. Initial capnography values and resuscitation outcomes of patients assisted by basic life support units in first instance; descriptive prospective study. *Signa Vitae*. 2021;17:86–91. doi: 10.22514/sv.2021.099

ADVANCED AIRWAY PLACEMENT DURING RESUSCITATION

Recommendations for Advanced Airway Placement During Resuscitation		
COR	LOE	Recommendations
1	B-NR	1. Frequent experience or frequent retraining is recommended for health care professionals who perform endotracheal intubation in adults with cardiac arrest.
1	C-LD	2. If advanced airway placement will interrupt chest compressions, health care professionals should defer insertion of the airway until the adult in cardiac arrest fails to respond to initial CPR and defibrillation attempts or obtains ROSC.
1	C-LD	3. Continuous waveform capnography is recommended in addition to clinical assessment as the most reliable method of confirming and monitoring correct placement of an endotracheal tube in adults with cardiac arrest.

Downloaded from http://ahajournals.org by on July 20, 2026

Recommendations for Advanced Airway Placement During Resuscitation (Continued)		
COR	LOE	Recommendations
1	C-EO	4. Emergency medical services systems that perform prehospital intubation in adults with cardiac arrest should provide a program of ongoing quality improvement to minimize complications and track overall supraglottic airway and endotracheal tube placement success rates.
2a	B-R	5. If an advanced airway is used for adults with OHCA, a supraglottic airway can be acceptable in settings with low endotracheal intubation success rate or minimal training opportunities.
2a	B-R	6. If an advanced airway is used, either a supra-glottic airway or endotracheal tube can be used for adults with OHCA in settings with high endotracheal intubation success rates or optimal training opportunities for endotracheal tube placement.
2a	B-NR	7. If an advanced airway is used for adults in cardiac arrest in the in-hospital setting by expert health care professionals trained in these procedures, either a supraglottic airway or endotracheal tube can be used.
2b	B-R	8. Either bag-mask ventilation or an advanced airway strategy may be considered during CPR for adult cardiac arrest in any setting depending on the situation and skill set of the professional.
2b	C-LD	9. If an advanced airway is in place, it may be reasonable for the clinician to deliver 1 breath every 6 s (10 breaths/min) while continuous chest compressions are being performed.

Synopsis

Airway management during cardiac arrest usually commences with a basic strategy such as BMV. In addition, it may be helpful for professionals to become proficient in one advanced airway strategy—endotracheal intubation (ETI) or supraglottic airway (SGA) insertion—as well as a second (backup) strategy for use if they are unable to establish the first-choice airway adjunct. Because placement of an advanced airway may result in interruption of chest compressions, a malpositioned device, or hyperventilation, professionals need weigh these risks against potential benefits. The 2019 focused update on use of advanced airways and Part 3: Adult Basic and Advanced Life Support from the 2020 Guidelines addressed the use of advanced airways in cardiac arrest. The 2020 Guidelines state that BMV or an advanced airway strategy may be considered during CPR for adult cardiac arrest in any setting.^{1,2} A key aspect of airway management decision-making is the health care professional's airway management skill and experience, frequent retraining and ongoing quality improvement to minimize airway management complications. Thus, the ultimate decision of the use, type, and timing of an advanced airway will require consideration of a host of patient and professional characteristics that are not easily defined in a global recommendation (Figure 5).

Recommendation-Specific Supportive Text

1. Frequent retraining is important to maintain a health care professional's psychomotor airway management skills.^{3,4} The manner in which this training is best accomplished is unclear. Research is necessary to address the specific type, amount, and frequency of training experiences that optimize airway management performance during cardiac arrest.
2. Although an advanced airway can be placed without interrupting chest compressions,⁵ interruptions frequently occur. Therefore, health care professionals should weigh the potential benefits of an advanced airway with the benefits of maintaining a high chest compression fraction.^{6–8} This is particularly true in cardiac arrest patients with shockable rhythms where CPR and defibrillation must take priority due to the strong association of early CPR and defibrillation with enhanced survival with favorable neurological outcome. In contrast, in arrests with non-shockable rhythms, early airway management is possible in resuscitations where CPR performance and epinephrine administration have already been optimized.
3. In a small clinical trial and several observational studies, waveform capnography was 100% specific for confirming endotracheal tube position during cardiac arrest.^{9–11} The sensitivity of waveform capnography decreases after a prolonged cardiac arrest.^{9–11} The use of waveform capnography to assess the placement of other advanced airways (eg, Combitube, laryngeal mask airway) has not been studied.
4. The writing group acknowledges the critical roles of tracking the overall success rate for systems performing ETI. This data will allow systems to make informed decisions as to whether practice should allow for ETI, move toward SGA, or simply use BMV for patients in cardiac arrest; recommendations will vary depending on the overall success rate in a given system. There is a clear need for further research with both patient- and professional- centered outcomes to assess the impact of prehospital quality improvement on selecting and maintaining the most appropriate airway management approach for EMS professionals within an agency.
- 5 and 6. One RCT of OHCA comparing SGA (iGel) to ETI in a non-physician-based EMS system (ETI success, 69%) found no difference in survival or survival with favorable neurologic outcome at hospital discharge.¹² A second RCT of OHCA comparing SGA (laryngeal tube) to ETI in a non-physician-based EMS system (ETI success,

Downloaded from http://ahajournals.org by on July 20, 2026




Figure 5. Schematic representation of ALS recommendations for use of advanced airways during CPR.

*Frequent experience or frequent retraining is recommended for health care professionals who perform endotracheal intubation. ALS indicates advanced life support; CPR, cardiopulmonary resuscitation; and EMS, emergency medical services.

52%) found both higher survival to hospital discharge and higher survival to hospital discharge with good neurologic outcome in the patients managed with SGA.¹³ A recent systematic review noted that while SGAs may be associated with faster placement and increased rate of ROSC, there is no clear survival benefit to SGA or ETI, and the effect of SGA versus ETI on neurological

survival remains uncertain.¹⁴ Outcomes likely depend on patient and professional characteristics, especially EMS professional exposure to initial and ongoing advanced airway training and experience. Precise thresholds for high or low ETI success rates have not been identified. The decision to place an advanced airway requires an understanding of patient and professional

characteristics that are not easily defined in a global recommendation. There is a need for further research, specifically on the interface between patient factors and the experience, training, tools and skills of the professional.

7. IHCA recommendations on advanced airway management have been extrapolated from OHCA. There is currently no high-certainty evidence that suggests benefit for any specific airway management strategy for IHCA. There is a need for further research to identify the IHCA patient and professional characteristics in airway management that impact clinical outcomes.
8. A 2024 ILCOR CoSTR summary¹⁵ demonstrated no new data to change the recommendation regarding the use of BMV or advanced airway strategies. One large RCT of OHCA comparing BMV to ETI in a physician-based EMS system showed no significant benefit for either technique for 28-day survival or survival with favorable neurological outcome.¹⁶ The success rate of ETI in this study was 98%, suggesting a relatively optimal setting for the potential success of ETI as an intervention. Further research is required to determine equivalence or superiority between the two approaches for acute airway management, as well as any potential benefit of a stepwise approach (eg, BVM followed by advanced airway placement).
9. A 2017 systematic review identified 1 observational human study and 10 animal studies comparing different ventilation rates after advanced airway placement.¹⁷ While no clear benefit from a rate of 10 breaths per minute was identified, no other rate was found to be superior. A 2017 ILCOR systematic review did not identify any new evidence to alter this recommendation, which was reiterated in the “2017 AHA Focused Update on Adult BLS and CPR Quality: An Update to the AHA Guidelines for CPR and Emergency Cardiovascular Care.”^{18,19}

REFERENCES

1. Panchal AR, Berg KM, Hirsch KG, Kudenchuk PJ, Del Rios M, Cabanas JG, Link MS, Kurz MC, Chan PS, Morley PT, et al. 2019 American Heart Association Focused Update on Advanced Cardiovascular Life Support: Use of Advanced Airways, Vasopressors, and Extracorporeal Cardiopulmonary Resuscitation During Cardiac Arrest: An Update to the American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2019;140:e881–e894. doi: 10.1161/CIR.0000000000000732
2. Panchal AR, Bartos JA, Cabanas JG, Donnino MW, Drennan IR, Hirsch KG, Kudenchuk PJ, Kurz MC, Lavonas EJ, Morley PT, et al; Adult Basic and Advanced Life Support Writing Group. Part 3: Adult Basic and Advanced Life Support: 2020 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2020;142:S366–S468. doi: 10.1161/CIR.0000000000000916
3. Wong ML, Carey S, Mader TJ, Wang HE; American Heart Association National Registry of Cardiopulmonary Resuscitation Investigators. Time to invasive airway placement and resuscitation outcomes after in

- hospital cardiopulmonary arrest. *Resuscitation*. 2010;81:182–186. doi: 10.1016/j.resuscitation.2009.10.027
4. Warner KJ, Carlblom D, Cooke CR, Bulger EM, Copass MK, Sharar SR. Paramedic training for proficient prehospital endotracheal intubation. *Prehosp Emerg Care*. 2010;14:103–108. doi: 10.3109/10903120903144858
 5. Gatward JJ, Thomas MJ, Nolan JP, Cook TM. Effect of chest compressions on the time taken to insert airway devices in a manikin. *Br J Anaesth*. 2008;100:351–356. doi: 10.1093/bja/aem364
 6. Vaillancourt C, Everson-Stewart S, Christenson J, Andrusiek D, Powell J, Nichol G, Cheskes S, Aufderheide TP, Berg R, Stiell IG; Resuscitation Outcomes Consortium Investigators. The impact of increased chest compression fraction on return of spontaneous circulation for out-of-hospital cardiac arrest patients not in ventricular fibrillation. *Resuscitation*. 2011;82:1501–1507. doi: 10.1016/j.resuscitation.2011.07.011
 7. Talikowska M, Tohira H, Finn J. Cardiopulmonary resuscitation quality and patient survival outcome in cardiac arrest: A systematic review and meta-analysis. *Resuscitation*. 2015;96:66–77. doi: 10.1016/j.resuscitation.2015.07.036
 8. Christenson J, Andrusiek D, Everson-Stewart S, Kudenchuk P, Hostler D, Powell J, Callaway CW, Bishop D, Vaillancourt C, Davis D, et al; Resuscitation Outcomes Consortium Investigators. Chest compression fraction determines survival in patients with out-of-hospital ventricular fibrillation. *Circulation*. 2009;120:1241–1247. doi: 10.1161/CIRCULATIONAHA.109.852202
 9. Tanigawa K, Takeda T, Goto E, Tanaka K. Accuracy and reliability of the self-inflating bulb to verify tracheal intubation in out-of-hospital cardiac arrest patients. *Anesthesiology*. 2000;93:1432–1436. doi: 10.1097/00000542-200012000-00015
 10. Takeda T, Tanigawa K, Tanaka H, Hayashi Y, Goto E, Tanaka K. The assessment of three methods to verify tracheal tube placement in the emergency setting. *Resuscitation*. 2003;56:153–157. doi: 10.1016/s0300-9572(02)00345-3
 11. Grmec S. Comparison of three different methods to confirm tracheal tube placement in emergency intubation. *Intensive Care Med*. 2002;28:701–704. doi: 10.1007/s00134-002-1290-x
 12. Bengier JR, Kirby K, Black S, Brett SJ, Clout M, Lazaroo MJ, Nolan JP, Reeves BC, Robinson M, Scott LJ, et al. Effect of a Strategy of a Supraglottic Airway Device vs Tracheal Intubation During Out-of-Hospital Cardiac Arrest on Functional Outcome: The AIRWAYS-2 Randomized Clinical Trial. *JAMA*. 2018;320:779–791. doi: 10.1001/jama.2018.11597
 13. Wang HE, Schmicker RH, Daya MR, Stephens SW, Idris AH, Carlson JN, Colella MR, Herren H, Hansen M, Richmond NJ, et al. Effect of a Strategy of Initial Laryngeal Tube Insertion vs Endotracheal Intubation on 72-Hour Survival in Adults With Out-of-Hospital Cardiac Arrest: A Randomized Clinical Trial. *JAMA*. 2018;320:769–778. doi: 10.1001/jama.2018.7044
 14. Forestell B, Ramsden S, Sharif S, Centofanti J, Al Lawati K, Fernando SM, Welsford M, Nichol G, Nolan JP, Rochweg B. Supraglottic Airway Versus Tracheal Intubation for Airway Management in Out-of-Hospital Cardiac Arrest: A Systematic Review, Meta-Analysis, and Trial Sequential Analysis of Randomized Controlled Trials. *Crit Care Med*. 2024;52:e89–e99. doi: 10.1097/CCM.00000000000006112
 15. Greif R, Bray JE, Djarr T, Drennan IR, Liley HG, Ng KC, Cheng A, Douma MJ, Scholefield BR, Smyth M, et al. 2024 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations: Summary From the Basic Life Support; Advanced Life Support; Pediatric Life Support; Neonatal Life Support; Education, Implementation, and Teams; and First Aid Task Forces. *Circulation*. 2024;150:e580–e687. doi: 10.1161/CIR.0000000000001288
 16. Jabre P, Penaloza A, Pinero D, Duchateau FX, Borron SW, Javaudin F, Richard O, de Longueville D, Bouilleau G, Devaud ML, et al. Effect of Bag-Mask Ventilation vs Endotracheal Intubation During Cardiopulmonary Resuscitation on Neurological Outcome After Out-of-Hospital Cardiorespiratory Arrest: A Randomized Clinical Trial. *JAMA*. 2018;319:779–787. doi: 10.1001/jama.2018.0156
 17. Vissers G, Soar J, Monsieus KG. Ventilation rate in adults with a tracheal tube during cardiopulmonary resuscitation: A systematic review. *Resuscitation*. 2017;119:5–12. doi: 10.1016/j.resuscitation.2017.07.018
 18. Ashoor HM, Lillie E, Zarin W, Pham B, Khan PA, Nincic V, Yazdi F, Ghassemi M, Ivory J, Cardoso R, et al; ILCOR Basic Life Support Task Force. Effectiveness of different compression-to-ventilation methods for cardiopulmonary resuscitation: A systematic review. *Resuscitation*. 2017;118:112–125. doi: 10.1016/j.resuscitation.2017.05.032
 19. Kleinman ME, Goldberger ZD, Rea T, Swor RA, Bobrow BJ, Brennan EE, Terry M, Hemphill R, Gazmuri RJ, Hazinski MF, et al. 2017 American Heart Association Focused Update on Adult Basic Life Support and Cardiopulmonary Resuscitation Quality: An Update to the American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2018;137:e7–e13. doi: 10.1161/CIR.0000000000000539

WIDE-COMPLEX TACHYCARDIA

Recommendations for the Management of Adult Patients With Wide-Complex Tachycardia		
COR	LOE	Recommendations
1	B-NR	1. Synchronized cardioversion is recommended for acute treatment of adult patients with hemodynamically unstable wide-complex tachycardia.
1	B-NR	2. Synchronized cardioversion is recommended for acute treatment of adult patients with hemodynamically stable wide-complex tachycardia when vagal maneuvers and pharmacological therapy is ineffective or contraindicated.
2b	B-R	3. Administration of IV amiodarone, procainamide, or sotalol may be considered for the treatment of adults with wide-complex tachycardia.
2b	B-NR	4. In hemodynamically stable adult patients with regular monomorphic wide-complex tachycardia, IV adenosine may be considered for treatment or aiding rhythm diagnosis when the cause of the rhythm cannot be determined.
3: Harm	B-NR	5. Verapamil and diltiazem should not be administered for adult patients with wide-complex tachycardia.
3: Harm	C-LD	6. Adenosine should not be administered for adult patients with hemodynamically unstable, irregularly irregular, or polymorphic wide-complex tachycardia.

Synopsis

Wide-complex tachycardia (WCT) is generally defined as a rhythm greater than 150/min with a QRS duration of 0.12 second or more.^{1,2} There are 4 means of achieving a wide QRS complex: (1) conduction of ventricular origin, (2) left or right bundle branch block, (3) conduction through a bypass tract, and (4) ventricular pacing. Rarely, hyperkalemia or sodium channel blockade can present with a WCT. Thus, WCTs can be ventricular (VT or VF), or supraventricular tachycardias resulting from mechanisms 2, 3, or 4. WCT can have a monomorphic or nonmonomorphic (ie, polymorphic) QRS. Polymorphic QRS complexes are typical of torsades de pointes, VF, or AF with a bypass tract. Most other rhythms are monomorphic. Patients with WCT can be hemodynamically stable or unstable. Hemodynamically stable patients with WCT allow the professional time to obtain a 12-lead electrocardiogram, place an IV, and administer IV drugs, while patients who are unstable require prompt cardioversion (Figure 6).

Recommendation-Specific Supportive Text

1. In hemodynamically unstable patients with WCT it is critical to promptly restore sinus rhythm. If the patient is unstable as a result of the WCT (eg, systolic blood pressure <80 mm Hg or altered mentation) immediate synchronized cardioversion is warranted.³
2. In stable patients with WCT, vagal maneuvers or adenosine may be attempted; however, if the patient remains in WCT following these therapies, synchronized cardioversion is recommended. If

time and clinical stability permits, administration of procedural sedation may be appropriate.³

3. Amiodarone has a relatively slow onset of action, and some solutes may cause hypotension. IV procainamide can also cause hypotension and must be infused no faster than 50 mg/min. IV sotalol is a newer agent without these concerns for hypotension. Direct comparisons of efficacy are generally between lidocaine and these three antiarrhythmic medications. Lidocaine's kinetic properties are less effective for VT at hemodynamically tolerated rates than amiodarone, procainamide, or sotalol,⁴ and it is unclear if lidocaine is more efficacious than placebo. Conclusions based on limited available evidence from small heterogeneous human studies for the treatment of stable, monomorphic VT, procainamide and sotalol each appear to have similar efficacy; amiodarone was not more effective than procainamide.^{5–9} Avoiding co-administration of these drugs is important because their combined effects may be arrhythmogenic. When any of these drugs are administered, the immediate availability of a defibrillator is encouraged because each drug may convert a WCT to a more rapid, malignant (ie, hemodynamically unstable) form, depending upon its underlying cause.
4. Adenosine blocks the atrioventricular (AV) node and decreases automaticity. Thus, WCTs in which the AV node is part of the circuit will be terminated by an effective dose of adenosine as described in therapies for narrow-complex tachycardia (NCT). Adequate adenosine administration may momentarily block the AV node, allowing the underlying rhythm to be correctly assessed. Therefore, adenosine can be diagnostic and therapeutic in a stable, regular, monomorphic WCT.^{10,11}
5. Verapamil is a calcium channel blocker that slows AV node conduction, shortens the refractory period of accessory pathways, and acts both as a negative inotrope and vasodilator. The vasodilatory effects are relatively profound, and thus, hypotension is common with administration. Verapamil will not terminate a WCT of ventricular origin and may cause profound hypotension leading to shock and cardiac arrest.^{12,13}
6. Adenosine will not terminate a WCT of ventricular origin, may cause profound hypotension leading to shock, or precipitate VF and cardiac arrest.^{14,15}

REFERENCES

1. Page RL, Joglar JA, Caldwell MA, Calkins H, Conti JB, Deal BJ, Estes NA 3rd, Field ME, Goldberger ZD, Hammill SC, et al; Evidence Review Committee Chair†. 2015 ACC/AHA/HRS Guideline for the Management of Adult Patients With Supraventricular Tachycardia: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *Circulation*. 2016;133:e506–e574. doi: 10.1161/CIR.0000000000000311
2. Al-Khatib SM, Stevenson WG, Ackerman MJ, Bryant WJ, Callans DJ, Curtis AB, Deal BJ, Dickfeld T, Field ME, Fonarow GC, et al. 2017 AHA/ACC/

Downloaded from <http://ahajournals.org> by on July 20, 2026




Figure 6. Adult Tachyarrhythmia With a Pulse Algorithm.

CHF indicates congestive heart failure; ECG, electrocardiogram; IV, intravenous; NS, normal saline; and VT, ventricular tachycardia.

HRS guideline for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *Heart Rhythm*. 2018;15:e73–e189. doi: 10.1016/j.hrthm.2017.10.036

- Roth A, Elkayam I, Shapira I, Sander J, Malov N, Kehati M, Golovner M. Effectiveness of prehospital synchronous direct-current cardioversion for supraventricular tachyarrhythmias causing unstable hemodynamic states. *Am J Cardiol*. 2003;91:489–491. doi: 10.1016/s0002-9149(02)03257-5
- Josephson ME. Lidocaine and sustained monomorphic ventricular tachycardia: fact or fiction. *Am J Cardiol*. 1996;78:82–83. doi: 10.1016/s0002-9149(96)00271-8
- Somberg JC, Bailin SJ, Haffajee CI, Paladino WP, Kerin NZ, Bridges D, Timar S, Molnar J, Amio-Aqueous I. Intravenous lidocaine versus intravenous amiodarone (in a new aqueous formulation) for incessant ventricular tachycardia. *Am J Cardiol*. 2002;90:853–859. doi: 10.1016/s0002-9149(02)02707-8
- Ortiz M, Martin A, Arribas F, Coll-Vinent B, Del Arco C, Peinado R, Almendral J, Investigators PS. Randomized comparison of intravenous procainamide vs. intravenous amiodarone for the acute treatment of tolerated wide QRS tachycardia: the PROCAMIO study. *Eur Heart J*. 2017;38:1329–1335. doi: 10.1093/eurheartj/ehw230
- Ho DS, Zecchin RP, Richards DA, Uther JB, Ross DL. Double-blind trial of lignocaine versus sotalol for acute termination of spontaneous sustained ventricular tachycardia. *Lancet*. 1994;344:18–23. doi: 10.1016/s0140-6736(94)91048-0
- Gorgels AP, van den Dool A, Hofs A, Mulleneers R, Smeets JL, Vos MA, Wellens HJ. Comparison of procainamide and lidocaine in terminating sustained monomorphic ventricular tachycardia. *Am J Cardiol*. 1996;78:43–46. doi: 10.1016/s0002-9149(96)00224-x
- deSouza IS, Martindale JL, Sinert R. Antidysrhythmic drug therapy for the termination of stable, monomorphic ventricular tachycardia: a systematic review. *Emerg Med J*. 2015;32:161–167. doi: 10.1136/emmermed-2013-202973
- Rankin AC, Oldroyd KG, Chong E, Rae AP, Cobbe SM. Value and limitations of adenosine in the diagnosis and treatment of narrow and broad complex tachycardias. *Br Heart J*. 1989;62:195–203. doi: 10.1136/hrt.62.3.195
- Marill KA, Wolfram S, Desouza IS, Nishijima DK, Kay D, Setnik GS, Stair TO, Ellnor PT. Adenosine for wide-complex tachycardia: efficacy and safety. *Crit Care Med*. 2009;37:2512–2518. doi: 10.1097/CCM.0b013e3181a93661
- Stewart RB, Bardy GH, Greene HL. Wide complex tachycardia: misdiagnosis and outcome after emergent therapy. *Ann Intern Med*. 1986;104:766–771. doi: 10.7326/0003-4819-104-6-766
- Buxton AE, Marchlinski FE, Doherty JU, Flores B, Josephson ME. Hazards of intravenous verapamil for sustained ventricular tachycardia. *Am J Cardiol*. 1987;59:1107–1110. doi: 10.1016/0002-9149(87)90857-5
- Gupta AK, Shah CP, Maheshwari A, Thakur RK, Hayes OW, Lokhandwala YY. Adenosine induced ventricular fibrillation in Wolff-Parkinson-White syndrome. *Pacing Clin Electrophysiol*. 2002;25:477–480. doi: 10.1046/j.1460-9592.2002.00477.x

15. Gulamhusein S, Ko P, Carruthers SG, Klein GJ. Acceleration of the ventricular response during atrial fibrillation in the Wolff-Parkinson-White syndrome after verapamil. *Circulation*. 1982;65:348–354. doi: 10.1161/01.cir.65.2.348

POLYMORPHIC VENTRICULAR
TACHYCARDIA

Recommendations for Treatment of Adults With Polymorphic Ventricular Tachycardia		
COR	LOE	Recommendations
1	B-NR	1. Immediate unsynchronized shock is recommended for adults with sustained polymorphic ventricular tachycardia.
2b	C-LD	2. Magnesium may be considered for treatment of adults with recurrences of polymorphic ventricular tachycardia associated with a long QT interval (torsades de pointes).
2b	C-LD	3. IV lidocaine, amiodarone, and measures to treat myocardial ischemia may be considered to treat recurrences of polymorphic VT in adults in the absence of a prolonged QT interval.
3: No Benefit	C-LD	4. Routine use of magnesium is not recommended for the treatment of polymorphic VT in adults with a normal QT interval.

Synopsis

Polymorphic VT refers to a wide-complex tachycardia of ventricular origin with differing configurations of the QRS complex from beat to beat. However, the most critical feature in the diagnosis and treatment of polymorphic VT is not the morphology of the rhythm but rather what is known (or suspected) about the patient's underlying QT interval. It is often difficult to know whether the tachycardia is pVT or ventricular fibrillation (VF). Most VF begins with pVT and if pVT is prolonged it will usually degenerate into VF (Figure 6).

Torsades de pointes is a form of polymorphic VT that is associated with a prolonged heart rate–corrected QT interval when the rhythm is normal and VT is not present. The risk for developing torsades increases when the corrected QT interval is greater than 500 milliseconds and accompanied by bradycardia.¹ Torsades can be due to an inherited genetic abnormality² or can be caused by drugs and electrolyte imbalances that lengthen the QT interval.³

Conversely, polymorphic VT not associated with a long QT is most often due to acute myocardial ischemia.^{4,5} Other potential causes include catecholaminergic polymorphic VT, a genetic abnormality in which polymorphic VT is provoked by exercise or emotion⁶; “short QT” syndrome, a form of polymorphic VT associated with an unusually short QT interval (corrected QT interval less than 330–370 milliseconds)^{7,8}; idiopathic right ventricular outflow tract polymorphic VT; and bidirectional VT seen in digitalis toxicity in which the axis of alternate QRS complexes shifts by 180°.⁹ There are no RCTs examining the acute pharmacological treatment of polymorphic VT. Supportive data, with and without long corrected QT interval, are largely based on case reports and case series.

Recommendation-Specific Supportive Text

1. Regardless of the underlying QT interval, all forms of polymorphic VT are considered hemodynamically and electrically unstable. Episodes of polymorphic VT may repeatedly recur and remit spontaneously, become sustained, or degenerate to VF. When the QRS complex of a VT is of uniform morphology, electric cardioversion synchronized to the QRS complex minimizes the risk of provoking VF during the vulnerable period of the cardiac cycle (T wave). In contrast, polymorphic VT cannot be synchronized reliably because of the differing characteristics of each QRS complex and requires high-energy (maximum manufacturer's setting) unsynchronized shock. Assessment of a patient's mental status is important when the appropriateness of sedation is considered before defibrillation. While effective in terminating polymorphic VT, defibrillation may not prevent its recurrence, for which pharmacological therapies are often required and the primary focus of subsequent recommendations.
2. Torsades de pointes typically presents in a recurring pattern of self-terminating, hemodynamically unstable polymorphic VT in context of a known or suspected long QT abnormality, often with an associated bradycardia. Termination of torsades by defibrillation may not prevent its recurrence, which requires additional pharmacological interventions. In small case series, IV magnesium has been effective in suppressing and preventing recurrences.^{10–13} Magnesium is believed to suppress fluctuations in the myocardial action potential that can trigger torsades.¹⁴ Correcting electrolyte abnormalities, particularly hypokalemia, is also advisable.
3. Treatment of torsades with antiarrhythmic medications may prolong the QT interval and promote the arrhythmia. When given acutely, β -adrenergic blockers can also precipitate torsades by causing or worsening bradycardia. In patients with bradycardia or pause-precipitated torsades, expert consultation is best sought for additional measures such as overdrive pacing or isoproterenol.¹⁵ There is literature to support instances of medication induced torsades, so prevention of drug induced prolonged QT is an important priority for professionals.
3. Polymorphic VT that is clearly not associated with QT prolongation, is most often triggered by acute myocardial ischemia or infarction and often rapidly degenerates into VF.^{3,5} Polymorphic VT may be difficult to differentiate from VF initially, but, as with other ventricular arrhythmias (VT and VF), is terminated with immediate defibrillation. However, termination of polymorphic VT with defibrillation may not prevent its recurrence, requiring additional pharmacological measures. No RCTs have been performed to determine the optimal pharmacological management of

Downloaded from <http://ahajournals.org> by on July 20, 2026

polymorphic VT. Treatment of causative myocardial ischemia (eg, β -adrenergic blockers or emergent coronary intervention) as well as lidocaine and amiodarone,^{16–22} in concert with defibrillation, may be synergetic when the arrhythmia is sustained. β -Adrenergic blockers have also been shown to reduce the incidence of ventricular arrhythmias in acute coronary syndromes.^{23,24}

Expert consultation is advisable when alternative causes of polymorphic VT are suspected, for which β -adrenergic blockers and antiarrhythmics may also have efficacy.^{6,25} Since the 2018 AHA Focused Update on Advanced Cardiovascular Life Support Use of Antiarrhythmic Drugs During and Immediately After Cardiac Arrest,²⁶ no new evidence has been identified.

4. In the absence of long QT, magnesium has not been shown to be effective in the treatment of polymorphic VT¹⁰ or other ventricular tachyarrhythmias.¹² A single case series with 5 patients showed no benefit of magnesium administration for polymorphic VT with normal QT. Since the 2018 AHA focused update on antiarrhythmic drugs during and immediately after cardiac arrest no new evidence has been identified.

REFERENCES

1. Chan A, Isbister GK, Kirkpatrick CM, Dufful SB. Drug-induced QT prolongation and torsades de pointes: evaluation of a QT nomogram. *QJM*. 2007;100:609–615. doi: 10.1093/qjmed/hcm072

2. Saprungruang A, Khongphatthanayothin A, Mauleekoonphairoj J, Wandee P, Kanjanathai S, Bhuiyan ZA, Wilde AAM, Poovorawan Y. Genotype and clinical characteristics of congenital long QT syndrome in Thailand. *Indian Pacing Electrophysiol J*. 2018;18:165–171. doi: 10.1016/j.ipej.2018.07.007

3. Drew BJ, Ackerman MJ, Funk M, Gibler WB, Kligfield P, Menon V, Philippides GJ, Roden DM, Zareba W; American Heart Association Acute Cardiac Care Committee of the Council on Clinical Cardiology. Prevention of torsade de pointes in hospital settings: a scientific statement from the American Heart Association and the American College of Cardiology Foundation. *J Am Coll Cardiol*. 2010;55:934–947. doi: 10.1016/j.jacc.2010.01.001

4. Wolfe CL, Nibley C, Bhandari A, Chatterjee K, Scheinman M. Polymorphous ventricular tachycardia associated with acute myocardial infarction. *Circulation*. 1991;84:1543–1551. doi: 10.1161/01.cir.84.4.1543

5. Pogwizd SM, Corr PB. Electrophysiologic mechanisms underlying arrhythmias due to reperfusion of ischemic myocardium. *Circulation*. 1987;76:404–426. doi: 10.1161/01.cir.76.2.404

6. Liu N, Ruan Y, Priori SG. Catecholaminergic polymorphic ventricular tachycardia. *Prog Cardiovasc Dis*. 2008;51:23–30. doi: 10.1016/j.pcad.2007.10.005

7. Gollob MH, Redpath CJ, Roberts JD. The short QT syndrome: proposed diagnostic criteria. *J Am Coll Cardiol*. 2011;57:802–812. doi: 10.1016/j.jacc.2010.09.048

8. Cross B, Homoud M, Link M, Foote C, Garlitski AC, Weinstock J, Estes NA 3rd. The short QT syndrome. *J Interv Card Electrophysiol*. 2011;31:25–31. doi: 10.1007/s10840-011-9566-0

9. Chapman M, Hargreaves M, Schneider H, Royle M. Bidirectional ventricular tachycardia associated with digoxin toxicity and with normal digoxin levels. *Heart Rhythm*. 2014;11:1222–1225. doi: 10.1016/j.hrthm.2014.03.050

10. Tzivoni D, Keren A, Cohen AM, Loebel H, Zahavi I, Chenzbraun A, Stern S. Magnesium therapy for torsades de pointes. *Am J Cardiol*. 1984;53:528–530. doi: 10.1016/0002-9149(84)90025-0

11. Tzivoni D, Banai S, Schuger C, Benhorin J, Keren A, Gottlieb S, Stern S. Treatment of torsade de pointes with magnesium sulfate. *Circulation*. 1988;77:392–397. doi: 10.1161/01.cir.77.2.392

12. Manz M, Jung W, Luderitz B. [Effect of magnesium on sustained ventricular tachycardia]. *Herz*. 1997;22:51–55. doi: 10.1007/BF03042655

13. Hoshino K, Ogawa K, Hishitani T, Isobe T, Etoh Y. Successful uses of magnesium sulfate for torsades de pointes in children with long QT syndrome. *Pediatr Int*. 2006;48:112–117. doi: 10.1111/j.1442-200X.2006.02177.x

14. Baker WL. Treating arrhythmias with adjunctive magnesium: identifying future research directions. *Eur Heart J Cardiovasc Pharmacother*. 2017;3:108–117. doi: 10.1093/ehjcvp/pvw028

15. DiSegni E, Klein HO, David D, Libhaber C, Kaplinsky E. Overdrive pacing in quinidine syncope and other long QT-interval syndromes. *Arch Intern Med*. 1980;140:1036–1040.

16. Vrana M, Pokorny J, Marcian P, Fejfar Z. Class I and III antiarrhythmic drugs for prevention of sudden cardiac death and management of postmyocardial infarction arrhythmias. A review. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub*. 2013;157:114–124. doi: 10.5507/bp.2013.030

17. Steg PG, James SK, Atar D, Badano LP, Blomstrom-Lundqvist C, Borger MA, Di Mario C, Dickstein K, Ducrocq G, et al.; Task Force on the management of ST-segment elevation ESC. ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J*. 2012;33:2569–2619. doi: 10.1093/eurheartj/ehs215

18. Nalliah CJ, Zaman S, Narayan A, Sullivan J, Kovoor P. Coronary artery reperfusion for ST elevation myocardial infarction is associated with shorter cycle length ventricular tachycardia and fewer spontaneous arrhythmias. *Europace*. 2014;16:1053–1060. doi: 10.1093/europace/eut307

19. Brady W, Meldon S, DeBehnke D. Comparison of prehospital monomorphic and polymorphic ventricular tachycardia: prevalence, response to therapy, and outcome. *Ann Emerg Med*. 1995;25:64–70. doi: 10.1016/s0196-0644(95)70357-8

20. Brady WJ, DeBehnke DJ, Laundrie D. Prevalence, therapeutic response, and outcome of ventricular tachycardia in the out-of-hospital setting: a comparison of monomorphic ventricular tachycardia, polymorphic ventricular tachycardia, and torsades de pointes. *Acad Emerg Med*. 1999;6:609–617. doi: 10.1111/j.1553-2712.1999.tb00414.x

21. Gorenek B, Lundqvist CB, Terradellas JB, Camm AJ, Hindricks G, Huber K, Kirchhof P, Kuck KH, Kudaiberdieva G, Lin T, et al. Cardiac arrhythmias in acute coronary syndromes: position paper from the joint EHRA, ACCA, and EAPCI task force. *Eur Heart J Acute Cardiovasc Care*. 2015;4:386. doi: 10.1177/2048872614550583

22. Luqman N, Sung RJ, Wang CL, Kuo CT. Myocardial ischemia and ventricular fibrillation: pathophysiology and clinical implications. *Int J Cardiol*. 2007;119:283–290. doi: 10.1016/j.ijcard.2006.09.016

23. Al-Khatib SM, Stevenson WG, Ackerman MJ, Bryant WJ, Callans DJ, Curtis AB, Deal BJ, Dickfeld T, Field ME, Fonarow GC, et al. 2017 AHA/ACC/HRS guideline for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *Heart Rhythm*. 2018;15:e73–e189. doi: 10.1016/j.hrthm.2017.10.036

24. Chatterjee S, Chaudhuri D, Vedanthan R, Fuster V, Ibanez B, Bangalore S, Mukherjee D. Early intravenous beta-blockers in patients with acute coronary syndrome—a meta-analysis of randomized trials. *Int J Cardiol*. 2013;168:915–921. doi: 10.1016/j.ijcard.2012.10.050

25. Van Houzen NE, Alsheikh-Ali AA, Garlitski AC, Homoud MK, Weinstock J, Link MS, Estes NA 3rd. Short QT syndrome review. *J Interv Card Electrophysiol*. 2008;23:1–5. doi: 10.1007/s10840-008-9201-x

26. Panchal AR, Berg KM, Kudenchuk PJ, Del Rios M, Hirsch KG, Link MS, Kurz MC, Chan PS, Cabañas JG, Morley PT, et al. 2018 American Heart Association Focused Update on Advanced Cardiovascular Life Support Use of Antiarrhythmic Drugs During and Immediately After Cardiac Arrest: An Update to the American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2018;138:e740–e749. doi: 10.1161/CIR.0000000000000613

REGULAR NARROW-COMPLEX TACHYCARDIA

Recommendations for the Treatment of Regular Narrow-Complex Tachycardia in Adults		
COR	LOE	Recommendations
1	B-R	1. Vagal maneuvers are recommended for acute treatment of adult patients with regular narrow-complex tachycardias.

Recommendations for the Treatment of Regular Narrow-Complex Tachycardia in Adults (Continued)		
COR	LOE	Recommendations
1	B-R	2. Adenosine is recommended for acute treatment of adult patients with regular narrow-complex tachycardias.
1	B-NR	3. Synchronized cardioversion is recommended for acute treatment of adult patients with hemodynamically unstable narrow-complex tachycardias.
1	B-NR	4. Synchronized cardioversion is recommended for acute treatment of adult patients with hemodynamically stable narrow-complex tachycardias when vagal maneuvers and pharmacological therapy is ineffective or contraindicated.
2a	B-R	5. Intravenous diltiazem or verapamil can be effective for acute treatment of adult patients with hemodynamically stable narrow-complex tachycardias that are regular.
2a	C-LD	6. Intravenous β -adrenergic blockers are reasonable for acute treatment of adult patients with hemodynamically stable narrow-complex tachycardias that are regular.

Synopsis

In most patients narrow-complex tachycardias (NCTs), which include sinus tachycardia, atrial flutter, AV nodal reentrant tachycardia, AV reentrant tachycardia, and other atrial tachycardia, and previously referred to imprecisely as supraventricular tachycardia, generally are not hemodynamically unstable, and health care professionals have an opportunity to attempt vagal maneuvers. Adenosine may also be effective in terminating arrhythmias in these patients.

For hemodynamically unstable patients with NCTs, electrical therapies are warranted (Figure 6). However, there must be a reasonable expectation that the NCT is causing instability rather than being a manifestation of instability caused by another clinical diagnosis (eg, sepsis). Hemodynamically unstable patients with NCT may have hypotension, acutely altered mental status, signs of shock, chest pain, or acute heart failure. Most patients with NCTs have conversion success rates of up to 98% with vagal maneuvers, pharmacological management, or both, described below (eg, adenosine, diltiazem, verapamil).

If nasally administered calcium channel blockers become available in the future, they will be evaluated for their efficacy and safety and considered for inclusion in these guidelines and the corresponding treatment algorithm.^{1–3}

These recommendations are supported by the 2015 American College of Cardiology, AHA, and Heart Rhythm Society Guidelines for the Management of Adult Patients With Supraventricular Tachycardia and a 2025 evidence update.

Recommendation-Specific Supportive Text

1. The Valsalva maneuver is effective in terminating 20 to 30% of NCTs.^{4,5} With modified techniques, termination improves to up to 50%.^{6–9} Vagal maneuvers

- include having the patient blow into a 10 cc syringe, bearing down, and carotid massage. However, more effective approaches have been described, such as passive leg raise, supine repositioning, and passive leg elevation immediately after Valsalva, blowing into a syringe attached to a sphygmomanometer, and pressing the epigastric region 10 seconds after Valsalva. When performing carotid massage, caution is advised because the technique can precipitate a stroke in patients with carotid atherosclerosis.
2. Adenosine has demonstrated very high rates of conversion of AV nodal reentrant tachycardia, AV reentrant tachycardia, and many atrial tachycardias.¹⁰ It also blocks the AV node, confirming a diagnosis of atrial flutter or fibrillation. Adenosine has a very short half-life and may be less effective if infused in a peripheral vein. Adequate dosing is assured by alteration of the NCT, including termination or the patient's symptoms of flushing and a feeling of dread. Adenosine may have profound effects in patients with central lines or post-heart transplant patients; 1 mg IV may be adequate to terminate the NCT. Adenosine can cause severe bronchospasm in patients with asthma, and thus, is contraindicated.
3. In hemodynamically unstable patients with NCT, it is critical to promptly restore sinus rhythm. If unstable, immediate cardioversion is warranted.¹¹
4. Cardioversion can be used to treat stable patients with NCT not responsive to vagal maneuvers and pharmacological therapy.¹¹ Assessment of a patient's mental status and clinical stability is important when the appropriateness of sedation is considered before cardioversion.
5. IV diltiazem and verapamil convert NCT to normal sinus rhythm in 64% to 98% of patients. The efficacy of verapamil is similar to adenosine but with significant concerns for inducing hypotension.^{12–18} Diltiazem and verapamil are particularly useful in patients with recurrent supraventricular tachycardia after treatment with adenosine because they are longer acting. Hypotension is a common side effect, and these medications must be infused slowly.¹⁹ For patients with NCT and suspicion of systolic heart failure, diltiazem and verapamil are best avoided.
6. Intravenous β -adrenergic blockers are generally safe and have been shown to terminate NCT in adult patients with hemodynamically stable narrow-complex tachycardias that are regular.

REFERENCES

1. Ip JE, Coutu B, Bennett MT, Pandey AS, Stambler BS, Sager P, Chen M, Shardonofsky S, Plat F, Camm AJ. Etipamil Nasal Spray for Conversion of Repeated Spontaneous Episodes of Paroxysmal Supraventricular Tachycardia During Long-Term Follow-Up: Results From the NODE-302 Study. *J Am Heart Assoc*. 2023;12:e028227. doi: 10.1161/JAHA.122.028227

2. Stambler BS, Camm AJ, Alings M, Dorian P, Heidbuchel H, Houtgraaf J, Kowey PR, Merino JL, Mondesert B, Piccini JP, et al; RAPID Investigators. Self-

Downloaded from http://ahajournals.org by on July 20, 2026

administered intranasal etipamil using a symptom-prompted, repeat-dose regimen for atrioventricular-nodal-dependent supraventricular tachycardia (RAPID): a multicentre, randomised trial. *Lancet*. 2023;402:118–128. doi: 10.1016/S0140-6736(23)00776-6

3. Stambler BS, Plat F, Sager PT, Shardonofsky S, Wight D, Potvin D, Pandey AS, Ip JE, Coutu B, Mondesert B, et al. First Randomized, Multicenter, Placebo-Controlled Study of Self-Administered Intranasal Etipamil for Acute Conversion of Spontaneous Paroxysmal Supraventricular Tachycardia (NODE-301). *Circ Arrhythm Electrophysiol*. 2022;15:e010915. doi: 10.1161/CIRCEP.122.010915

4. Lim SH, Anantharaman V, Teo WS, Goh PP, Tan AT. Comparison of treatment of supraventricular tachycardia by Valsalva maneuver and carotid sinus massage. *Ann Emerg Med*. 1998;31:30–35.

5. Smith GD, Fry MM, Taylor D, Morgans A, Cantwell K. Effectiveness of the Valsalva Manoeuvre for reversion of supraventricular tachycardia. *Cochrane Database Syst Rev*. 2015;2015:CD009502. doi: 10.1002/14651858.CD009502.pub3

6. Appelboom A, Reuben A, Mann C, Gagg J, Ewings P, Barton A, Lobban T, Dayer M, Vickery J, Bengier J; REVERT trial collaborators. Postural modification to the standard Valsalva manoeuvre for emergency treatment of supraventricular tachycardias (REVERT): a randomised controlled trial. *Lancet*. 2015;386:1747–1753. doi: 10.1016/S0140-6736(15)61485-4

7. Chen C, Tam TK, Sun S, Guo Y, Teng P, Jin D, Xu L, Liu X. A multicenter randomized controlled trial of a modified Valsalva maneuver for cardioversion of supraventricular tachycardias. *Am J Emerg Med*. 2020;38:1077–1081. doi: 10.1016/j.ajem.2019.158371

8. Corbacioglu SK, Akinci E, Cevik Y, Aytar H, Oncul MV, Akkan S, Uzunosmanoglu H. Comparing the success rates of standard and modified Valsalva maneuvers to terminate PSVT: A randomized controlled trial. *Am J Emerg Med*. 2017;35:1662–1665. doi: 10.1016/j.ajem.2017.05.034

9. Lan Q, Han B, Wu F, Peng Y, Zhang Z. Modified Valsalva maneuver for treatment of supraventricular tachycardias: A Meta-analysis. *Am J Emerg Med*. 2021;50:507–512. doi: 10.1016/j.ajem.2021.08.067

10. DiMarco JP, Sellers TD, Berne RM, West GA, Belardinelli L. Adenosine: electrophysiologic effects and therapeutic use for terminating paroxysmal supraventricular tachycardia. *Circulation*. 1983;68:1254–1263. doi: 10.1161/01.cir.68.6.1254

11. Roth A, Elkayam I, Shapira I, Sander J, Malov N, Kehati M, Golovner M. Effectiveness of prehospital synchronous direct-current cardioversion for supraventricular tachyarrhythmias causing unstable hemodynamic states. *Am J Cardiol*. 2003;91:489–491. doi: 10.1016/s0002-9149(02)03257-5

12. Ahmad F, Abu Sneh M, Patel RS, Rohit Reddy S, Llukmani A, Hashim A, Haddad DR, Gordon DK. In The Line of Treatment: A Systematic Review of Paroxysmal Supraventricular Tachycardia. *Cureus*. 2021;13:e15502. doi: 10.7759/cureus.15502

13. Alabed S, Sabouni A, Providencia R, Atallah E, Qintar M, Chico TJ. Adenosine versus intravenous calcium channel antagonists for supraventricular tachycardia. *Cochrane Database Syst Rev*. 2017;10:CD005154. doi: 10.1002/14651858.CD005154.pub4

14. Delaney B, Loy J, Kelly AM. The relative efficacy of adenosine versus verapamil for the treatment of stable paroxysmal supraventricular tachycardia in adults: a meta-analysis. *Eur J Emerg Med*. 2011;18:148–152. doi: 10.1097/MEJ.0b013e3283400ba2

15. DiMarco JP, Miles W, Akhtar M, Milstein S, Sharma AD, Platia E, McGovern B, Scheinman MM, Govier WC. Adenosine for paroxysmal supraventricular tachycardia: dose ranging and comparison with verapamil. Assessment in placebo-controlled, multicenter trials. The Adenosine for PSVT Study Group. *Ann Intern Med*. 1990;113:104–110. doi: 10.7326/0003-4819-113-2-104

16. Lim SH, Anantharaman V, Teo WS. Slow-infusion of calcium channel blockers in the emergency management of supraventricular tachycardia. *Resuscitation*. 2002;52:167–174. doi: 10.1016/s0300-9572(01)00459-2

17. Madsen CD, Pointer JE, Lynch TG. A comparison of adenosine and verapamil for the treatment of supraventricular tachycardia in the prehospital setting. *Ann Emerg Med*. 1995;25:649–655. doi: 10.1016/s0196-0644(95)70179-6

18. Pachchipulusu VK, Paudel Y, Patel A, Najam B, Desai HN, Illango J, Seffah KD, Kumar M, Naveen N, Khan S. Efficacy of Adenosine Versus Verapamil in Terminating Acute Attacks of Paroxysmal Supraventricular Tachycardia: A Systematic Review. *Cureus*. 2023;15:e45946. doi: 10.7759/cureus.45946

19. Lim SH, Anantharaman V, Teo WS, Chan YH. Slow infusion of calcium channel blockers compared with intravenous adenosine in the emergency treatment of supraventricular tachycardia. *Resuscitation*. 2009;80:523–528. doi: 10.1016/j.resuscitation.2009.01.017

ATRIAL FIBRILLATION OR FLUTTER WITH RAPID VENTRICULAR RESPONSE: PART 1 ELECTRICAL THERAPIES

Recommendations for Electrical Therapies for Atrial Fibrillation/Flutter		
COR	LOE	Recommendations
1	C-LD	1. In adult patients with hemodynamic instability attributable to atrial fibrillation or atrial flutter with rapid ventricular rates, immediate electrical cardioversion should be performed to restore sinus rhythm.
1	C-LD	2. Urgent direct-current cardioversion of new onset atrial fibrillation or atrial flutter with rapid ventricular rates in the setting of acute coronary syndrome is recommended for adult patients with hemodynamic compromise, ongoing ischemia, or inadequate rate control.
2a	B-R	3. For synchronized cardioversion of atrial fibrillation in adults using any currently US-approved biphasic waveform defibrillator, an initial energy setting of at least 200 J is reasonable and incremented in the event of shock failure, depending on the biphasic defibrillator used.
2b	B-R	4. The usefulness of double synchronized cardioversion of atrial fibrillation in adults as an initial treatment strategy is uncertain.
2b	C-LD	5. For synchronized cardioversion of atrial flutter in adults, an initial energy setting of 200 J may be reasonable and incremented in the event of shock failure, depending on the biphasic defibrillator used.
2b	C-LD	6. The usefulness of double synchronized cardioversion as a rescue treatment for shock-refractory atrial fibrillation in adults is uncertain.

Synopsis

Atrial fibrillation (AF) is a disorganized atrial rhythm due to multiple reentry (self-circulating) circuits throughout atrial myocardium. Atrial flutter is a more organized tachyarrhythmia caused by a single macroreentrant atrial circuit. The two arrhythmias are common, often co-exist, and are treated similarly. Treating AF/flutter with rapid ventricular rates depends on hemodynamic stability. Unstable patients and those with rate-related ischemia often need urgent electrical cardioversion (Figure 7). Stable patients can be treated pharmacologically to achieve rate or rhythm control. Conversion of AF/flutter greater than 48 hours in nonanticoagulated patients risks thromboembolic events. For acute AF/flutter assessed as less than 48 hours, decision-making is more nuanced, requiring expert consultation unless emergent cardioversion is needed. Pre-excited AF/flutter (Wolff-Parkinson-White) is covered under Wide-Complex Tachycardias.¹

Electrical cardioversion is influenced by many factors. Technical features (beyond energy setting) affecting shock success include pad dimensions, pad location, minimizing impedance barriers, shock waveform, and timing (proper electrocardiogram




Figure 7. Electrical Cardioversion Algorithm.

IV indicates intravenous; and VT, ventricular tachycardia.

synchronization).^{2,3} Clinical factors influencing cardioversion outcome include body habitus, cardiac status (ventricular function, atrial size), rhythm duration, and concomitant medication use. Any of these, alone or together, can affect outcome as much or more than the energy setting itself.^{4,5} Apart from energy, electrical cardioversion requires regard for the details of how the procedure itself is conducted and the characteristics of the patient in whom it is performed.⁶

These recommendations are supported by the “2023 ACC/AHA/ACCP/HRS Guideline for the Management of Atrial Fibrillation” and a 2025 evidence update.

Recommendation-Specific Supportive Text

1 and 2. Uncontrolled tachycardia may impair ventricular filling, cardiac output, and coronary perfusion while increasing myocardial oxygen demand.

While an expeditious trial of medications, intravenous fluids, or both may be appropriate in some cases, unstable patients or patients with ongoing cardiac ischemia caused or exacerbated by atrial fibrillation/flutter require prompt cardioversion.^{7–9} One should also consider whether the arrhythmia is causative or secondary to a rapid ventricular response driven by underlying factors (eg, sepsis), as this may guide initial pharmacologic strategies for hemodynamic stabilization. Limited data exist on these strategies in hemodynamically unstable patients; however, the hemodynamic benefits of successful cardioversion have been documented.^{10,11} In addition, risk of hypotension and hypoperfusion with negative inotropes has been demonstrated even in normotensive patients.^{12–14}

Hemodynamically unstable patients and those with ongoing cardiac ischemia are likely to benefit from improved hemodynamic status from immediate restoration of sinus rhythm and avoidance of hypotension caused by pharmacological therapies. Depending on the clinical scenario, patients cardioverted from atrial fibrillation/flutter of greater than 48 hours' duration are candidates for anticoagulation. Appropriate selection of patients and medications for anticoagulation are addressed elsewhere and beyond the scope of these guidelines.¹

3. Guidance for monophasic waveform cardioversion of AF/flutter is found in previous guidelines,¹⁵ displaced by biphasic waveforms due to their greater efficacy. Given proprietary differences in electrical characteristics, guidelines previously deferred biphasic cardioversion energy settings to manufacturer specifications.¹⁶ However, interim randomized head-to-head trials together with a network meta-analysis involving >3000 AF patients found that 200-J shocks achieved greater than 90% cumulative cardioversion success across all 3 biphasic platforms currently available in the United States.^{6,17–21}

Previous guidelines advocated lower energy settings over concerns that higher energies posed greater risk for cardiac injury and complications.^{18,22–25} Subsequent clinical studies observed no association between starting biphasic shock energies of up to 360 J with energy-associated elevations in cardiac-specific enzymes, postcardioversion arrhythmias, cardiac dysfunction, or other sequelae.^{26–28} Notably, low-energy monophasic shocks were significantly more likely to provoke VF when cardioverting AF, and AF when cardioverting atrial flutter than a 200-J or greater setting.²⁹

This observation is consistent with the upper limit of vulnerability principle: that an energy threshold exists above which a shock will not induce fibrillation.^{30,31} A higher starting energy was also found to be more efficient than “starting low, titrating higher” approaches, achieved greater first-shock cardioversion success, required fewer overall shocks, and could shorten the period of anesthesia needed for serial shock attempts.^{18,22}

Considering efficacy, safety, efficiency, simplicity, and consistency with other professional guidelines, evidence supports a starting energy of 200 J.^{16–21} Regardless of energy setting, appropriate sedation for cardioversion is advisable in awake patients, given shock is a painful procedure.

4. One approach to synchronized cardioversion is double synchronized cardioversion. Administering 2 simultaneously-synchronized high-energy shocks from separate defibrillators and pads has been described anecdotally, in small case series, and a single randomized trial. As an “initial” cardioversion strategy, a modestly-sized randomized trial comprised of class II obesity patients (39% had prior cardioversion attempts) found first shock efficacy improved with 400 J double synchronized cardioversion versus 200 J synchronized cardioversion (98% versus 87%, $p=0.002$).³² Although not consistently reported, double synchronized cardioversion has not been associated with proarrhythmia, hemodynamic complications, cardiac-specific enzyme elevation,²⁸ or postprocedure discomfort but has caused defibrillator damage.³³

A concern in all studies has been the vagueness of procedural specifications, particularly how 2 independently-initiated shocks from separate defibrillators being reliably simultaneously-synchronized was assured.³⁴ Given the high success of optimal synchronized cardioversion using biphasic waveforms, the incremental benefit from double synchronized cardioversion is modest.

5. Organized rhythms like atrial flutter, caused by a discrete reentry circuit, require less energy for termination than AF, whose multiple reentry circuits broadly engulf the atrial myocardium.³⁵ However, periodic transitions between the two arrhythmias are common. Distinguishing AF from flutter in an emergent setting with only single-lead monitoring can be challenging. Studies evaluating rectilinear and truncated exponential biphasic waveforms available in the United States, while indirect and fewer in number than for AF, have shown comparable cumulative

cardioversion success for atrial flutter between these waveforms at the same energy settings, ranging from 50% to 75% at 50 J, improving to 85% to 90% at 100 J, and approaching 100% at 200 J.^{23,28,36–38} Using a higher (200 J) initial energy setting for flutter cardioversion also affords shorter procedural times without causing cardiac injury or compromising safety,^{23,26,36–38} accompanied by a lower risk of provoking AF or undertreating AF if mistaken as flutter.^{27,36} The likelihood of greater efficacy, efficiency, and simplicity, without safety concerns, supports a starting energy of 200 J for flutter cardioversion using any currently US-marketed biphasic defibrillator; incremented in the event of shock failure, depending on the defibrillator's features.^{23,28,36,37}

6. **Shock-refractory AF**, defined as AF that remains incessant after maximum-energy shocks, is relatively uncommon with biphasic shock waveforms and distinguished from recurrent AF, which returns after an initially successful shock. As in many situations, shock success can be optimized by minimizing transthoracic impedance³ by improving pad-skin contact and applying pad pressure; relocating pads to redirect the shock vector; or adding antiarrhythmic drugs to alter the defibrillation threshold.^{2,5,6,39,40}

Another possibility is the use of double-synchronized cardioversion.⁴¹ Data on this approach are limited, including procedural uncertainties described above (item 4). Observational studies found double-synchronized cardioversion to be successful in 75% to 100% of patients who had previously failed standard cardioversion.^{41–43} Alternatively, a large prospective observational study evaluated a systematic approach to shock-refractory AF. It found applied pad pressure improved synchronized cardioversion success from 87% to 93% and with pad repositioning to 96%, after which the added increment from double-synchronized cardioversion only modestly improved to 98%.²

REFERENCES

1. Joglar JA, Chung MK, Armbruster AL, Benjamin EJ, Chyou JY, Cronin EM, Deswal A, Eckhardt LL, Goldberger ZD, Gopinathannair R, et al; Peer Review Committee Members. 2023 ACC/AHA/ACCP/HRS Guideline for the Diagnosis and Management of Atrial Fibrillation: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2024;149:e1–e156. doi: 10.1161/CIR.0000000000001193
2. Darrat Y, Leung S, Elayi L, Parrott K, Ogunbayo G, Kotter J, Sorrell V, Gupta V, Anaya P, Morales G, et al. A stepwise external cardioversion protocol for atrial fibrillation to maximize acute success rate. *Europace*. 2023;25:828–834. doi: 10.1093/europace/ead009
3. Sadek MM, Chaugai V, Cleland MJ, Zakutney TJ, Birnie DH, Ramirez FD. Association between transthoracic impedance and electrical cardioversion success with biphasic defibrillators: An analysis of 1055 shocks for atrial fibrillation and flutter. *Clin Cardiol*. 2018;41:666–670. doi: 10.1002/clc.22947
4. Brandes A, Crijns H, Rienstra M, Kirchhof P, Grove EL, Pedersen KB, Van Gelder IC. Cardioversion of atrial fibrillation and atrial flutter revisited: current evidence and practical guidance for a common procedure. *Europace*. 2020;22:1149–1161. doi: 10.1093/europace/euaa057
5. Kukendrarajah K, Ahmad M, Carrington M, Ioannou A, Taylor J, Razvi Y, Papageorgiou N, Mead GE, Nevis IF, D'Ascenzo F, et al. External electrical and pharmacological cardioversion for atrial fibrillation, atrial flutter or atrial tachycardias: a network meta-analysis. *Cochrane Database Syst Rev*. 2024;6:CD013255. doi: 10.1002/14651858.CD013255.pub2
6. Nguyen ST, Belley-Cote EP, Ibrahim O, Um KJ, Lengyel A, Adli T, Qiu Y, Wong M, Sibilio S, Benz AP, et al. Techniques improving electrical cardioversion success for patients with atrial fibrillation: a systematic review and meta-analysis. *Europace*. 2023;25:318–330. doi: 10.1093/europace/ead019
7. January CT, Wann LS, Alpert JS, Calkins H, Cigarroa JE, Cleveland JC Jr, Conti JB, Ellinor PT, Ezekowitz MD, Field ME, et al; ACC/AHA Task Force Members. 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines and the Heart Rhythm Society. *Circulation*. 2014;130:e199–e267. doi: 10.1161/CIR.0000000000000041
8. January CT, Wann LS, Calkins H, Chen LY, Cigarroa JE, Cleveland JC Jr, Ellinor PT, Ezekowitz MD, Field ME, Furie KL, et al. 2019 AHA/ACC/HRS Focused Update of the 2014 AHA/ACC/HRS Guideline for the Management of Patients With Atrial Fibrillation: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society in Collaboration With the Society of Thoracic Surgeons. *Circulation*. 2019;140:e125–e151. doi: 10.1161/CIR.0000000000000665
9. McMurray J, Kober L, Robertson M, Dargie H, Colucci W, Lopez-Sendon J, Remme W, Sharpe DN, Ford I. Antiarrhythmic effect of carvedilol after acute myocardial infarction: results of the Carvedilol Post-Infarct Survival Control in Left Ventricular Dysfunction (CAPRICORN) trial. *J Am Coll Cardiol*. 2005;45:525–530. doi: 10.1016/j.jacc.2004.09.076
10. DeMaria AN, Lies JE, King JF, Miller RR, Amsterdam EA, Mason DT. Echographic assessment of atrial transport, mitral movement, and ventricular performance following electroversion of supraventricular arrhythmias. *Circulation*. 1975;51:273–282. doi: 10.1161/01.cir.51.2.273
11. Raymond RJ, Lee AJ, Messineo FC, Manning WJ, Silverman DI. Cardiac performance early after cardioversion from atrial fibrillation. *Am Heart J*. 1998;136:435–442. doi: 10.1016/s0002-8703(98)70217-0
12. Delle Karth G, Geppert A, Neunteufl T, Priglinger U, Haumer M, Gschwandtner M, Siostrzonek P, Heinz G. Amiodarone versus diltiazem for rate control in critically ill patients with atrial tachyarrhythmias. *Crit Care Med*. 2001;29:1149–1153. doi: 10.1097/00003246-200106000-00011
13. Ellenbogen KA, Dias VC, Heywood JT, Mirvis DM. A placebo-controlled trial of continuous intravenous diltiazem infusion for 24-hour heart rate control during atrial fibrillation and atrial flutter: a multicenter study. *J Am Coll Cardiol*. 1991;18:891–897. doi: 10.1016/0735-1097(91)90743-s
14. Platia EV, Michelson EL, Porterfield JK, Das G. Esmolol versus verapamil in the acute treatment of atrial fibrillation or atrial flutter. *Am J Cardiol*. 1989;63:925–929. doi: 10.1016/0002-9149(89)90141-0
15. Guidelines 2000 for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. Part 6: advanced cardiovascular life support: section 2: defibrillation. The American Heart Association in collaboration with the International Liaison Committee on Resuscitation. *Circulation*. 2000;102:I90–I94.
16. Panchal AR, Bartos JA, Cabanas JG, Donnino MW, Drennan IR, Hirsch KG, Kudenchuk PJ, Kurz MC, Lavonas EJ, Morley PT, et al; Adult Basic and Advanced Life Support Writing Group. Part 3: Adult Basic and Advanced Life Support: 2020 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2020;142:S366–S468. doi: 10.1161/CIR.0000000000000916
17. Neal S, Ngarmukos T, Lessard D, Rosenthal L. Comparison of the efficacy and safety of two biphasic defibrillator waveforms for the conversion of atrial fibrillation to sinus rhythm. *Am J Cardiol*. 2003;92:810–814. doi: 10.1016/s0002-9149(03)00888-9
18. Kim ML, Kim SG, Park DS, Gross JN, Ferrick KJ, Palma EC, Fisher JD. Comparison of rectilinear biphasic waveform energy versus truncated exponential biphasic waveform energy for transthoracic cardioversion of atrial fibrillation. *Am J Cardiol*. 2004;94:1438–1440. doi: 10.1016/j.amjcard.2004.07.149

19. Inacio JF, da Rosa Mdos S, Shah J, Rosario J, Vissoci JR, Manica AL, Rodrigues CG. Monophasic and biphasic shock for transthoracic conversion of atrial fibrillation: Systematic review and network meta-analysis. *Resuscitation*. 2016;100:66–75. doi: 10.1016/j.resuscitation.2015.12.009

20. Deakin CD, Connelly S, Wharton R, Yuen HM. A comparison of rectilinear and truncated exponential biphasic waveforms in elective cardioversion of atrial fibrillation: a prospective randomized controlled trial. *Resuscitation*. 2013;84:286–291. doi: 10.1016/j.resuscitation.2012.07.010

21. Alatawi F, Gurevitz O, White RD, Ammash NM, Malouf JF, Bruce CJ, Moon BS, Rosales AG, Hodge D, Hammill SC, et al. Prospective, randomized comparison of two biphasic waveforms for the efficacy and safety of transthoracic biphasic cardioversion of atrial fibrillation. *Heart Rhythm*. 2005;2:382–387. doi: 10.1016/j.hrthm.2004.12.024

22. Resnekov L, McDonald L. Complications in 220 patients with cardiac dysrhythmias treated by phased direct current shock, and indications for electroconversion. *Br Heart J*. 1967;29:926–936. doi: 10.1136/hrt.29.6.926

23. Mortensen K, Risius T, Schwemer TF, Aydin MA, Koster R, Klemm HU, Lutomsky B, Meinertz T, Ventura R, Willems S. Biphasic versus monophasic shock for external cardioversion of atrial flutter: a prospective, randomized trial. *Cardiology*. 2008;111:57–62. doi: 10.1159/000113429

24. DiCola VC, Freedman GS, Downing SE, Zaret BL. Myocardial uptake of technetium-99m stannous pyrophosphate following direct current transthoracic countershock. *Circulation*. 1976;54:980–986. doi: 10.1161/01.cir.54.6.980

25. Dahl CF, Ewy GA, Warner ED, Thomas ED. Myocardial necrosis from direct current countershock. Effect of paddle electrode size and time interval between discharges. *Circulation*. 1974;50:956–961. doi: 10.1161/01.cir.50.5.956

26. Schmidt AS, Lauridsen KG, Torp P, Bach LF, Rickers H, Lofgren B. Maximum-fixed energy shocks for cardioverting atrial fibrillation. *Eur Heart J*. 2020;41:626–631. doi: 10.1093/eurheartj/ehz585

27. Lobo R, White RD, Donato LJ, Wockenfus AM, Kelley BR, Melduni RM, Jaffe AS. Absence of significant myocardial injury following elective direct current cardioversion for atrial fibrillation. *Heart Rhythm O2*. 2023;4:180–186. doi: 10.1016/j.hroo.2022.12.004

28. Lobo R, Jaffe AS, Cahill C, Blake O, Abbas S, Meany TB, Hennessy T, Kiernan TJ. Significance of High-Sensitivity Troponin T After Elective External Direct Current Cardioversion for Atrial Fibrillation or Atrial Flutter. *Am J Cardiol*. 2018;121:188–192. doi: 10.1016/j.amjcard.2017.10.009

29. Gallagher MM, Yap YG, Padula M, Ward DE, Rowland E, Camm AJ. Arrhythmic complications of electrical cardioversion: relationship to shock energy. *Int J Cardiol*. 2008;123:307–312. doi: 10.1016/j.ijcard.2006.12.014

30. Swerdlow CD, Shehata M, Chen PS. Using the upper limit of vulnerability to assess defibrillation efficacy at implantation of ICDs. *Pacing Clin Electrophysiol*. 2007;30:258–270. doi: 10.1111/j.1540-8159.2007.00659.x

31. Swerdlow CD, Martin DJ, Kass RM, Davie S, Mandel WJ, Gang ES, Chen PS. The zone of vulnerability to T wave shocks in humans. *J Cardiovasc Electrophysiol*. 1997;8:145–154. doi: 10.1111/j.1540-8167.1997.tb00776.x

32. Aymond JD, Sanchez AM, Castine MR, Bernard ML, Khatib S, Hiltbold AE, Polin GM, Rogers PA, Dominic PS, Velasco-Gonzalez C, et al. Dual vs Single Cardioversion of Atrial Fibrillation in Patients With Obesity: A Randomized Clinical Trial. *JAMA Cardiol*. 2024;9:641–648. doi: 10.1001/jamacardio.2024.1091

33. Gerstein NS, McLean AR, Stecker EC, Schulman PM. External Defibrillator Damage Associated With Attempted Synchronized Dual-Dose Cardioversion. *Ann Emerg Med*. 2018;71:109–112. doi: 10.1016/j.annemergmed.2017.04.005

34. Bjerregaard P, El-Shafei A, Janosik DL, Schiller L, Quattromani A. Double external direct-current shocks for refractory atrial fibrillation. *Am J Cardiol*. 1999;83:972–4, A10. doi: 10.1016/s0002-9149(98)01048-0

35. Gallagher MM, Guo XH, Poloniecki JD, Guan Yap Y, Ward D, Camm AJ. Initial energy setting, outcome and efficiency in direct current cardioversion of atrial fibrillation and flutter. *J Am Coll Cardiol*. 2001;38:1498–1504. doi: 10.1016/s0735-1097(01)01540-6

36. Stec S, Krynski T, Kulakowski P. Efficacy of low energy rectilinear biphasic cardioversion for regular atrial tachyarrhythmias. *Cardiol J*. 2011;18:33–38.

37. Reisinger J, Gstrein C, Winter T, Zeindhofer E, Hollinger K, Mori M, Schiller A, Winter A, Geiger H, Siostrzonek P. Optimization of initial energy for cardioversion of atrial tachyarrhythmias with biphasic shocks. *Am J Emerg Med*. 2010;28:159–165. doi: 10.1016/j.ajem.2008.10.028

38. Mortensen K, Aydin MA, Schwemer TF, Ventura R, Reppel M, Bode F, Mletzko R, Schunkert H, Risius T. Low energy biphasic cardioversion of atrial flutter: results from a pilot trial. *Int J Cardiol*. 2010;145:368–370. doi: 10.1016/j.ijcard.2010.02.042

39. Ferreira D, Mikhail P, Lim J, Ray M, Dwivedi J, Briennes S, Butel-Simoes L, Meere W, Bland A, Howden N, et al. Manual Chest PRESSURE During Direct Current Cardioversion for Atrial Fibrillation: A Randomized Control Trial (PRESSURE-AF). *JACC Clin Electrophysiol*. 2024;10:2207–2213. doi: 10.1016/j.jacep.2024.05.037

40. Velazquez-Rodriguez E, Perez-Sandoval HA, Rangel-Rojo FJ. Orthogonal electrical cardioversion in atrial fibrillation refractory to biphasic shocks: a case series. *Eur Heart J Case Rep*. 2020;4:1–5. doi: 10.1093/ehjcr/ytaa343

41. Gardner MW, Yadava M, Raitt MH, Elman MR, Zarraga IG, MacMurdy KS, Dalouk KA, Jessel PM. Effectiveness of dual external direct current cardioversion for initial cardioversion in atrial fibrillation. *J Cardiovasc Electrophysiol*. 2019;30:1636–1643. doi: 10.1111/jce.13994

42. Velázquez-Rodríguez E, Pérez-Sandoval HA, Rangel-Rojo FJ. Orthogonal electrical cardioversion in atrial fibrillation refractory to biphasic shocks: a case series. *Eur Heart J Case Rep*. 2020;4:1–5. doi: 10.1093/ehjcr/ytaa343

43. Kabukcu M, Demircioglu F, Yanik E, Minareci K, Ersel-Tuzuner F. Simultaneous double external DC shock technique for refractory atrial fibrillation in concomitant heart disease. *Jpn Heart J*. 2004;45:929–936. doi: 10.1536/jhj.45.929

44. Marrouche NF, Bardy GH, Frielitz HJ, Gunther J, Brachmann J. Quadraple pads approach for external cardioversion of atrial fibrillation. *Pacing Clin Electrophysiol*. 2001;24(Part 1):1321–1324. doi: 10.1046/j.1460-9592.2001.01321.x

ATRIAL FIBRILLATION AND FLUTTER WITH RAPID VENTRICULAR RESPONSE: PART 2 MEDICAL THERAPIES

Recommendations for Medical Therapies for Atrial Fibrillation/Flutter		
COR	LOE	Recommendations
1	B-NR	1. IV administration of a β -adrenergic blocker or nondihydropyridine calcium channel antagonist is recommended to slow the ventricular heart rate in the acute setting in adult patients with atrial fibrillation or atrial flutter with rapid ventricular response without preexcitation.
2a	B-NR	2. IV amiodarone can be useful for rate control in critically ill adult patients with atrial fibrillation with rapid ventricular response without preexcitation.
3: Harm	C-LD	3. In adult patients with preexcitation atrial fibrillation or atrial flutter, digoxin, nondihydropyridine calcium channel antagonists, β -adrenergic blockers, and IV amiodarone should not be administered because they may increase the ventricular response and result in VF.
3: Harm	C-EO	4. Nondihydropyridine calcium channel antagonists and IV β -adrenergic blockers should not be used in adult patients with left ventricular systolic dysfunction and decompensated heart failure because these may lead to further hemodynamic compromise.

Synopsis

Control of rapid heart rates in the emergency setting are achievable using an intravenous nondihydropyridine calcium channel antagonist (eg, diltiazem, verapamil) or β -adrenergic blocker (eg, metoprolol, esmolol). Pharmacologic rhythm control (chemical cardioversion) employs intravenous drugs such as amiodarone, procainamide, or sotalol to convert AF/flutter to sinus rhythm and prevent its recurrence. Amiodarone is unique in that it can also be used for rate control in patients with heart failure in

whom calcium channel or β -adrenergic blockers may not be tolerated.

These recommendations are supported by the “2023 ACC/AHA/ACCP/HRS Guideline for the Diagnosis and Management of Atrial Fibrillation” and a 2025 evidence update.¹

Recommendation-Specific Supportive Text

1. Clinical trial evidence shows that nondihydropyridine calcium channel antagonists (eg, diltiazem, verapamil) and β -adrenergic blockers (eg, metoprolol, esmolol, propranolol), are effective for rate control in patients with atrial fibrillation/flutter.^{2–9} Systematic reviews show that IV diltiazem improves heart rate control in atrial fibrillation/flutter without preexcitation with rapid ventricular response when compared to IV metoprolol.^{10–18} However, most of the included studies were limited by small sample size, retrospective data, and many did not report relevant clinical outcomes and comorbidities. Randomized clinical trials examining verapamil are lacking, thus, diltiazem is generally preferred.
2. Amiodarone is effective for rate control in patients with atrial fibrillation/flutter without preexcitation in patients with contraindications for nondihydropyridine calcium channel antagonists or β -adrenergic blockers.^{2–9} Digoxin is less commonly used in the acute setting because of the slow onset of effect.^{19–21} In an observational study, the optimal amiodarone dosing for heart rate control (<115–80/min) was examined in 80 critically ill patients, most with cardiogenic shock or sepsis.²² The dose needed to achieve heart rate control was nearly twice as high in patients with persistent versus paroxysmal atrial arrhythmias, but patients who

were taking amiodarone prior to intensive care unit admission required lower doses.

3. Based on limited case reports and small case series, there is concern that patients with concomitant preexcitation and atrial fibrillation or atrial flutter may develop VF in response to accelerated ventricular response after the administration of AV nodal blocking agents such as digoxin, nondihydropyridine calcium channel antagonists, β -adrenergic blockers, or IV amiodarone.^{23–26} In this setting, cardioversion is recommended as the most appropriate management.
4. Because of their negative inotropic effect, nondihydropyridine calcium channel antagonists (eg, diltiazem, verapamil) may further decompensate patients with left ventricular systolic dysfunction and symptomatic heart failure.⁷ Patients with heart failure with preserved ejection fraction may tolerate better nondihydropyridine calcium channel antagonists. While β -blockers can be safe in patients with cardiomyopathy, they can be harmful in patients who are decompensated.^{20,21,27} Limited evidence suggests that intravenous diltiazem may cause increased risk of decompensation compared to intravenous β -adrenergic blockers, but more data are needed to support preferential use.²⁸ Multiple studies have demonstrated β -adrenergic blockers are safe for use in patients with chronic obstructive pulmonary disease.²⁹ Refer to Table 2.

REFERENCES

1. Joglar JA, Chung MK, Armbruster AL, Benjamin EJ, Chyou JY, Cronin EM, Deswal A, Eckhardt LL, Goldberger ZD, Gopinathannair R, et al; Peer Review Committee Members. 2023 ACC/AHA/ACCP/HRS Guideline for the Diagnosis and Management of Atrial Fibrillation: A Report of the American College of Cardiology/American Heart Association Joint Committee

Table 2. IV Medications Commonly Used for Acute Rate Control in Atrial Fibrillation and Atrial Flutter

Medication	Bolus dose	Infusion rate	Notes
Nondihydropyridine calcium channel blockers			
Diltiazem	0.25 mg/kg IV bolus over 2 min	5–10 mg/h	Avoid in hypotension, heart failure, cardiomyopathy, and acute coronary syndromes
Verapamil	0.075–0.15 mg/kg IV bolus over 2 min; may give an additional dose after 30 min if no response	0.005 mg/kg per min	Avoid in hypotension, heart failure, cardiomyopathy, and acute coronary syndromes
β -Adrenergic blockers			
Metoprolol	2.5–5 mg over 2 min, up to 3 doses		Avoid in decompensated heart failure
Esmolol	500 mcg/kg IV over 1 min	50–300 mcg/kg per min	Short duration of action; avoid in decompensated heart failure
Propranolol	1 mg IV over 1 min, up to 3 doses		Avoid in decompensated heart failure
Other medications			
Amiodarone	300 mg IV over 1 h	10–50 mg/h over 24 h	Multiple dosing schemes exist for amiodarone
Digoxin	0.25 mg IV, repeated to maximum dose 1.5 mg over 24 h		Typically used as adjunctive therapy with another option from above; caution in patients with renal impairment

Downloaded from http://ahajournals.org by on July 20, 2026

on Clinical Practice Guidelines. *Circulation*. 2024;149:e1–e156. doi: 10.1161/CIR.0000000000001193

2. Siu CW, Lau CP, Lee WL, Lam KF, Tse HF. Intravenous diltiazem is superior to intravenous amiodarone or digoxin for achieving ventricular rate control in patients with acute uncomplicated atrial fibrillation. *Crit Care Med*. 2009;37:2174–9; quiz 2180. doi: 10.1097/CCM.0b013e3181a02f56

3. Salerno DM, Dias VC, Kleiger RE, Tschida VH, Sung RJ, Sami M, Giorgi LV. Efficacy and safety of intravenous diltiazem for treatment of atrial fibrillation and atrial flutter. The Diltiazem-Atrial Fibrillation/Flutter Study Group. *Am J Cardiol*. 1989;63:1046–1051. doi: 10.1016/0002-9149(89)90076-3

4. Platia EV, Michelson EL, Porterfield JK, Das G. Esmolol versus verapamil in the acute treatment of atrial fibrillation or atrial flutter. *Am J Cardiol*. 1989;63:925–929. doi: 10.1016/0002-9149(89)90141-0

5. Hou ZY, Chang MS, Chen CY, Tu MS, Lin SL, Chiang HT, Woosley RL. Acute treatment of recent-onset atrial fibrillation and flutter with a tailored dosing regimen of intravenous amiodarone. A randomized, digoxin-controlled study. *Eur Heart J*. 1995;16:521–528. doi: 10.1093/oxfordjournals.eurheartj.a060945

6. Ellenbogen KA, Dias VC, Plumb VJ, Heywood JT, Mirvis DM. A placebo-controlled trial of continuous intravenous diltiazem infusion for 24-hour heart rate control during atrial fibrillation and atrial flutter: a multicenter study. *J Am Coll Cardiol*. 1991;18:891–897. doi: 10.1016/0735-1097(91)90743-s

7. Delle Karth G, Geppert A, Neunteufl T, Priglinger U, Haumer M, Gschwandtner M, Siostrzonek P, Heinz G. Amiodarone versus diltiazem for rate control in critically ill patients with atrial tachyarrhythmias. *Crit Care Med*. 2001;29:1149–1153. doi: 10.1097/00003246-200106000-00011

8. Abrams J, Allen J, Allin D, Anderson J, Anderson S, Blanski L, Chadda K, DiBianco R, Favrot L, Gonzalez J. Efficacy and safety of esmolol vs propranolol in the treatment of supraventricular tachyarrhythmias: a multicenter double-blind clinical trial. *Am Heart J*. 1985;110:913–922. doi: 10.1016/0002-8703(85)90185-1

9. Clemp H, Wood MA, Gilligan DM, Ellenbogen KA. Intravenous amiodarone for acute heart rate control in the critically ill patient with atrial tachyarrhythmias. *Am J Cardiol*. 1998;81:594–598. doi: 10.1016/s0002-9149(97)00962-4

10. Ward SM, Radke J, Calhoun C, Caporossi J, Hall GA, Matuskowitz AJ, Weeda ER, Weant KA. Weight-based versus non-weight-based diltiazem dosing in the setting of atrial fibrillation with rapid ventricular response. *Am J Emerg Med*. 2020;38:2271–2276. doi: 10.1016/j.ajem.2019.09.020

11. Sharda SC, Bhatia MS. Comparison of diltiazem and metoprolol for atrial fibrillation with rapid ventricular rate: Systematic review and meta-analysis. *Indian Heart J*. 2022;74:494–499. doi: 10.1016/j.ihj.2022.10.195

12. Martindale JL, deSouza IS, Silverberg M, Freedman J, Sinert R. beta-Blockers versus calcium channel blockers for acute rate control of atrial fibrillation with rapid ventricular response: a systematic review. *Eur J Emerg Med*. 2015;22:150–154. doi: 10.1097/MEJ.0000000000000227

13. Lan Q, Wu F, Han B, Ma L, Han J, Yao Y. Intravenous diltiazem versus metoprolol for atrial fibrillation with rapid ventricular rate: A meta-analysis. *Am J Emerg Med*. 2022;51:248–256. doi: 10.1016/j.ajem.2021.08.082

14. Kuang P, Mah ND, Barton CA, Miura AJ, Tanas LR, Ran R. Achieving ventricular rate control using metoprolol in beta-blocker-naïve patients vs patients on chronic beta-blocker therapy. *Am J Emerg Med*. 2016;34:606–608. doi: 10.1016/j.ajem.2015.12.084

15. Hirschy R, Ackerbauer KA, Peksa GD, O'Donnell EP, DeMott JM. Metoprolol vs. diltiazem in the acute management of atrial fibrillation in patients with heart failure with reduced ejection fraction. *Am J Emerg Med*. 2019;37:80–84. doi: 10.1016/j.ajem.2018.04.062

16. Feeney ME, Rowe SLB, Mah ND, Barton CA, Ran R. Achieving ventricular rate control in patients taking chronic beta-blocker therapy. *Am J Emerg Med*. 2018;36:110–113. doi: 10.1016/j.ajem.2017.09.013

17. Burum A, Carino J, McBeth M, Samuel N, Hintze TD. A Systematic Review of Weight-Based Metoprolol for Acute Atrial Fibrillation with Rapid Ventricular Rate. *Emerg Med Int*. 2023;2023:3138064. doi: 10.1155/2023/3138064

18. Bolton A, Paudel B, Adhaduk M, Alsuhailani M, Samuelson R, Schweizer ML, Hodgson-Zingman D. Intravenous Diltiazem Versus Metoprolol in Acute Rate Control of Atrial Fibrillation/Flutter and Rapid Ventricular Response: A Meta-Analysis of Randomized and Observational Studies. *Am J Cardiovasc Drugs*. 2024;24:103–115. doi: 10.1007/s40256-023-00615-3

19. Shojaaee M, Feizi B, Miri R, Etemadi J, Feizi AH. Intravenous Amiodarone versus Digoxin in Atrial Fibrillation Rate Control; a Clinical Trial. *Emerg (Tehran)*. 2017;5:e29.

20. January CT, Wann LS, Calkins H, Chen LY, Cigarroa JE, Cleveland JC Jr, Ellinor PT, Ezekowitz MD, Field ME, Furie KL, et al. 2019 AHA/ACC/

HRS Focused Update of the 2014 AHA/ACC/HRS Guideline for the Management of Patients With Atrial Fibrillation: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society in Collaboration With the Society of Thoracic Surgeons. *Circulation*. 2019;140:e125–e151. doi: 10.1161/CIR.0000000000000665

21. January CT, Wann LS, Alpert JS, Calkins H, Cigarroa JE, Cleveland JC Jr, Conti JB, Ellinor PT, Ezekowitz MD, Field ME, et al; ACC/AHA Task Force Members. 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines and the Heart Rhythm Society. *Circulation*. 2014;130:e199–e267. doi: 10.1161/CIR.0000000000000041

22. Salem JE, El-Aissaoui M, Alazard M, Hulot JS, Aissaoui N, Le-Heuzey JY, Funck-Brentano C, Faisy C, Urien S. Modeling of Amiodarone Effect on Heart Rate Control in Critically Ill Patients with Atrial Tachyarrhythmias. *Clin Pharmacokinet*. 2016;55:991–1002. doi: 10.1007/s40262-016-0372-9

23. Kim RJ, Gerling BR, Kono AT, Greenberg ML. Precipitation of ventricular fibrillation by intravenous diltiazem and metoprolol in a young patient with occult Wolff-Parkinson-White syndrome. *Pacing Clin Electrophysiol*. 2008;31:776–779. doi: 10.1111/j.1540-8159.2008.01086.x

24. Jacob AS, Nielsen DH, Gianelly RE. Fatal ventricular fibrillation following verapamil in Wolff-Parkinson-White syndrome with atrial fibrillation. *Ann Emerg Med*. 1985;14:159–160. doi: 10.1016/s0196-0644(85)81080-5

25. Gulamhusein S, Ko P, Carruthers SG, Klein GJ. Acceleration of the ventricular response during atrial fibrillation in the Wolff-Parkinson-White syndrome after verapamil. *Circulation*. 1982;65:348–354. doi: 10.1161/01.cir.65.2.348

26. Boriani G, Biffi M, Frabetti L, Azzolini U, Sabbatani P, Bronzetti G, Capucci A, Magnani B. Ventricular fibrillation after intravenous amiodarone in Wolff-Parkinson-White syndrome with atrial fibrillation. *Am Heart J*. 1996;131:1214–1216. doi: 10.1016/s0002-8703(96)90098-8

27. Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr, Drazner MH, Fonarow GC, Geraci SA, Horwich T, Januzzi JL, et al. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association task force on practice guidelines. *Circulation*. 2013;128:e240–e327. doi: 10.1161/CIR.0b013e31829e8776

28. Hasbrouck M, Nguyen TT. Acute management of atrial fibrillation in congestive heart failure with reduced ejection fraction in the emergency department. *Am J Emerg Med*. 2022;58:39–42. doi: 10.1016/j.ajem.2022.03.058

29. Salpeter S, Ormiston T, Salpeter E. Cardioselective beta-blockers for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*. 2005;2005:CD003566. doi: 10.1002/14651858.CD003566.pub2

INITIAL MANAGEMENT OF BRADYCARDIA

Recommendations for Initial Management of Bradycardia		
COR	LOE	Recommendations
1	C-EO	1. In adult patients presenting with acute symptomatic bradycardia, evaluation and treatment of reversible causes are recommended.
2a	B-NR	2. In adult patients with acute bradycardia associated with hemodynamic compromise, administration of atropine is reasonable to increase heart rate.
2a	C-LD	3. In adult patients with persistent hemodynamically unstable bradycardia refractory to medical therapy, temporary transvenous pacing is reasonable to increase heart rate and improve symptoms.
2b	C-LD	4. If bradycardia is unresponsive to atropine, IV adrenergic agonists with rate-accelerating effects (eg, epinephrine, dopamine) or transcutaneous pacing may be effective while the adult patient is prepared for emergent transvenous temporary pacing if required.
2b	C-EO	5. Immediate pacing might be considered in unstable adult patients with high-degree AV block when IV/IO access is not available.

Synopsis

Bradycardia has historically been defined as a ventricular heart rate of less than 60/min. However, the “2018 ACC/AHA/HRS Guideline on the Evaluation and Management of Patients With Bradycardia and Cardiac Conduction Delay” adopted a lower threshold of less than 50/min, reflecting population studies that commonly use this lower cutoff.¹ Bradycardia can be a normal finding in healthy adults, athletes, or during sleep.¹ The clinical manifestation of bradycardia ranges from stable (asymptomatic) through symptomatic to unstable (symptomatic and associated with signs of poor perfusion). Patients with symptomatic bradycardia can present with dizziness or light-headedness, acute altered mental status, ischemic chest discomfort, acute heart failure, hypotension, or syncope due to decreased cardiac output and tissue hypoperfusion. Clinicians must evaluate bradycardia within the clinical context to determine whether it is a manifestation of another underlying disease process (eg, myocardial ischemia, hypoxia, electrolyte disturbance, acidosis, hypothermia) or the primary driver of the clinical presentation. Immediate treatment of symptomatic bradycardia as described in Figure 8 is focused on both supporting perfusion and identifying and reversing the underlying cause(s). The 2018 ACC/AHA/HRS Guideline provides comprehensive recommendations on the evaluation and management of chronic stable and unstable bradycardia.

These recommendations remain supported by the 2018 ACC/AHA/HRS Guideline and a 2025 evidence update.

Recommendation-Specific Supportive Text

1. Symptomatic bradycardia may be caused by a number of potentially reversible or treatable causes, including myocardial ischemia, hypoxemia, hypothyroidism, infections, structural heart disease, increased vagal tone, metabolic derangement, electrolyte abnormalities, or medications/toxins.¹ Bradycardia may be difficult to resolve without addressing the underlying cause(s), underscoring the importance of evaluating potential etiologies while simultaneously initiating emergent stabilization measures. A large retrospective review of prehospital patients undergoing transcutaneous pacing identified hypoxia and initial nonbradycardic rhythms as factors associated with pacing failure and increased mortality, emphasizing the critical need to address underlying causes alongside stabilization efforts.²
2. Atropine has been shown to be effective for the treatment of symptomatic bradycardia in observational studies.^{3–7} In a study in patients with OHCA and a nonshockable heart rhythm (asystole or pulseless electrical activity), the addition of atropine following epinephrine was a predictor of survival to

hospital admission.⁸ In a study among patients with IHCA, the administration of atropine for nonshockable heart rhythms did not improve survival.⁹

3. When symptomatic bradycardia is refractory to medical management, cardiac pacing is a reasonable alternative. Transcutaneous pacing is noninvasive, can be performed quickly at bedside, and is typically used as a bridge to placement of a temporary transvenous pacing catheter. Transcutaneous pacing is often painful in the conscious patient and procedural tolerance may necessitate sedation. Evidence for transvenous pacing consists of limited observational studies, many of which have focused on indications and the relatively high complication rate (including bloodstream infections and pneumothorax, among others).^{10–13} When the heart rate does not improve with medications and shock persists, transvenous pacing can improve the heart rate and symptoms until more definitive treatment (correction of underlying cause or permanent pacemaker placement) can be implemented.
4. If atropine is ineffective, alternative agents to increase heart rate and blood pressure or transcutaneous pacing are reasonable alternatives. For medical management of a periarrest patient, epinephrine has gained popularity, including IV infusion and use of “push-dose” administration for acute bradycardia and hypotension. Studies on push-dose epinephrine for bradycardia specifically are lacking, although limited data support its use for hypotension.^{14,15} Use of push-dose vasopressors require careful attention to correct dosing. Medication errors and adverse hemodynamic effects have been reported.¹⁶ Dopamine infusion can also increase heart rate.¹⁷ There are limited studies comparing medications to transcutaneous pacing for the treatment of bradycardia. A randomized feasibility study in patients failing atropine compared dopamine to transcutaneous pacing and found no difference in survival to discharge.¹⁷ Whether to trial transcutaneous pacing, epinephrine, dopamine, or other vasoactive agent will likely, therefore, depend on clinician experience and resources available.
5. For severe symptomatic bradycardia causing shock, if no IV or IO access is available, immediate transcutaneous pacing while access is being pursued may be undertaken. A 2006 systematic review involving seven studies of transcutaneous pacing for symptomatic bradycardia and bradysystolic cardiac arrest in the prehospital setting did not find a benefit from pacing compared with standard ALS, although a subgroup analysis from a single trial suggested a possible benefit in patients with symptomatic bradycardia.¹⁸ More recent studies have underscored persistent challenges for




Figure 8. Adult Bradycardia With a Pulse Algorithm.

clinicians in achieving and maintaining both electrical and mechanical capture during pacing.^{19,20} These difficulties may limit patient benefit in certain scenarios, emphasizing the need for improved methods to distinguish true electrical capture from false capture. Future research is needed to determine optimal pad placement and develop reliable methods for confirming initial and sustained capture, such as the practicality and preferred location of pulse palpation, hemodynamic findings, pulse

oximetry characteristics, continuous ETCO₂ monitoring, and ultrasound.

REFERENCES

1. Kusumoto FM, Schoenfeld MH, Barrett C, Edgerton JR, Ellenbogen KA, Gold MR, Goldschlager NF, Hamilton RM, Joglar JA, Kim RJ, et al. 2018 ACC/AHA/HRS Guideline on the Evaluation and Management of Patients With Bradycardia and Cardiac Conduction Delay: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *Circulation*. 2019;140:e382–e482. doi: 10.1161/CIR.0000000000000628

2. Smida T, Voges L, Crowe R, Scheidler J, Bardes J. Prehospital Transcutaneous Cardiac Pacing in the United States: Treatment Epidemiology, Predictors of Treatment Failure, and Associated Outcomes. *Prehosp Emerg Care*. 2024;29:578–585. doi: 10.1080/10903127.2024.2393768

3. Swart G, Brady WJ Jr, DeBehnke DJ, Ma OJ, Aufderheide TP. Acute myocardial infarction complicated by hemodynamically unstable bradyarrhythmia: prehospital and ED treatment with atropine. *Am J Emerg Med*. 1999;17:647–652. doi: 10.1016/s0735-6757(99)90151-1

4. Smith I, Monk TG, White PF. Comparison of transesophageal atrial pacing with anticholinergic drugs for the treatment of intraoperative bradycardia. *Anesth Analg*. 1994;78:245–252. doi: 10.1213/00000539-199402000-00009

5. Chadda KD, Lichstein E, Gupta PK, Kourtesis P. Effects of atropine in patients with bradyarrhythmia complicating myocardial infarction. Usefulness of an optimum dose for overdrive. *Am J Med*. 1977;63:503–510. doi: 10.1016/0002-9343(77)90194-2

6. Chadda KD, Lichstein E, Gupta PK, Choy R. Bradycardia-hypotension syndrome in acute myocardial infarction. Reappraisal of the overdrive effects of atropine. *Am J Med*. 1975;59:158–164. doi: 10.1016/0002-9343(75)90349-6

7. Brady WJ, Swart G, DeBehnke DJ, Ma OJ, Aufderheide TP. The efficacy of atropine in the treatment of hemodynamically unstable bradycardia and atrioventricular block: prehospital and emergency department considerations. *Resuscitation*. 1999;41:47–55. doi: 10.1016/s0300-9572(99)00032-5

8. Yano T, Kawana R, Yamauchi K, Endo G, Nagamine Y. The Additive Effect of Atropine Sulfate during Cardiopulmonary Resuscitation in Out-of-hospital Non-traumatic Cardiac Arrest Patients with Non-shockable Rhythm. *Intern Med*. 2019;58:1713–1721. doi: 10.2169/internalmedicine.1932-18

9. Holmberg MJ, Moskowitz A, Wiberg S, Grossestreuer AV, Yankama T, Witten L, Perman SM, Donnino MW, Andersen LW; American Heart Association's Get With The Guidelines®-Resuscitation Investigators. Guideline removal of atropine and survival after adult in-hospital cardiac arrest with a non-shockable rhythm. *Resuscitation*. 2019;137:69–77. doi: 10.1016/j.resuscitation.2019.02.002

10. McCann P. A review of temporary cardiac pacing wires. *Indian Pacing Electrophysiol J*. 2007;7:40–49.

11. Jou YL, Hsu HP, Tuan TC, Wang KL, Lin YJ, Lo LW, Hu YF, Kong CW, Chang SL, Chen SA. Trends of temporary pacemaker implant and underlying disease substrate. *Pacing Clin Electrophysiol*. 2010;33:1475–1484. doi: 10.1111/j.1540-8159.2010.02893.x

12. Ferguson JD, Banning AP, Bashir Y. Randomised trial of temporary cardiac pacing with semirigid and balloon-flotation electrode catheters. *Lancet*. 1997;349:1883. doi: 10.1016/S0140-6736(97)24026-2

13. Sofi NU, Sinha SK, Ali A, Samrat S, Razi MM, Sharma AK, Sachan M, Pandey U, Thakur R. Unguided temporary pacing via jugular/subclavian

vein in an emergency department of a high-volume tertiary care hospital of India: its safety, efficacy, and practicability. *Egypt Heart J*. 2022;74:33. doi: 10.1186/s43044-022-00271-z

14. Singer S, Pope H, Fuller BM, Gibson G. The safety and efficacy of push dose vasopressors in critically ill adults. *Am J Emerg Med*. 2022;61:137–142. doi: 10.1016/j.ajem.2022.08.055

15. Nawrocki PS, Poremba M, Lawner BJ. Push Dose Epinephrine Use in the Management of Hypotension During Critical Care Transport. *Prehosp Emerg Care*. 2020;24:188–195. doi: 10.1080/10903127.2019.1588443

16. Cole JB, Knack SK, Karl ER, Horton GB, Satpathy R, Driver BE. Human Errors and Adverse Hemodynamic Events Related to "Push Dose Pressors" in the Emergency Department. *J Med Toxicol*. 2019;15:276–286. doi: 10.1007/s13181-019-00716-z

17. Morrison LJ, Long J, Vermeulen M, Schwartz B, Sawadsky B, Frank J, Cameron B, Burgess R, Shield J, Bagley P, et al. A randomized controlled feasibility trial comparing safety and effectiveness of prehospital pacing versus conventional treatment: 'PrePACE'. *Resuscitation*. 2008;76:341–349. doi: 10.1016/j.resuscitation.2007.08.008

18. Sherbino J, Verbeek PR, MacDonald RD, Sawadsky BV, McDonald AC, Morrison LJ. Prehospital transcutaneous cardiac pacing for symptomatic bradycardia or bradyasystolic cardiac arrest: a systematic review. *Resuscitation*. 2006;70:193–200. doi: 10.1016/j.resuscitation.2005.11.019

19. Moayed S, Patel P, Brady N, Witting M, Dickfeld TL. Anteroposterior pacer pad position is better than anterolateral for transcutaneous cardiac pacing. *Resuscitation*. 2022;181:140–146. doi: 10.1016/j.resuscitation.2022.11.009

20. Kimbrell J, Kreinbrook J, Poke D, Kalosza B, Geldner J, Shekhar AC, Miele A, Bouthillet T, Vega J. False Electrical Capture in Prehospital Transcutaneous Pacing by Paramedics: A Case Series. *Prehosp Emerg Care*. 2024;28:928–936. doi: 10.1080/10903127.2024.2321287

ARTICLE INFORMATION

The American Heart Association requests that this document be cited as follows: Wigginton JG, Agarwal S, Bartos JA, Coute RA, Drennan IR, Haamid A, Kudenchuk RJ, Link MS, Panchal AR, Pelter MM, Del Rios M, Rodriguez AJ, Perman SM, Sanko S, Kotini-Shah P, Kurz MC. Part 9: adult advanced life support: 2025 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*. 2025;152(suppl 2):S538–S577. doi: 10.1161/CIR.0000000000001376

Acknowledgments

The ALS Writing Group would like to acknowledge Drs Matthew Neth and Sean Morgan of the BLS Writing Group for their contributions to recommendations regarding Adjuncts for Performance of CPR.

Disclosures

Appendix 1. Writing Group Disclosures

Writing Group Member	Employment	Research Grant	Other Research Support	Speakers' Bureau/ Honoraria	Expert Witness	Ownership Interest	Consultant/ Advisory Board	Other
Michael C. Kurz	University of Chicago	None	None	None	None	None	None	None
Jane G. Wigginton	UT Dallas	None	None	None	None	None	None	None
Sachin Agarwal	Columbia University Irving Medical Center	NIH†	None	None	None	None	None	None
Jason A. Bartos	University of Minnesota	None	None	None	None	None	None	None
Ryan A. Coute	University of Alabama at Birmingham	NIH†; AHA*; NIH*; Zoll Foundation*	None	None	None	None	None	None
Ian R. Drennan	University of Toronto; Ornge Critical Care and Transport Medicine; Sunnybrook Health Sciences (Canada)	None	None	ZOLL Medical†	None	None	None	None
Ameera Haamid	University of Chicago	None	None	None	None	None	None	None
Peter J. Kudenchuk	University of Washington Medical Center	NIH/NINDS†	None	None	None	None	None	None

(Continued)

Appendix 1. Continued

Writing Group Member	Employment	Research Grant	Other Research Support	Speakers' Bureau/Honoraria	Expert Witness	Ownership Interest	Consultant/Advisory Board	Other
Mark S. Link	UT Southwestern Medical Center	None	None	None	None	None	None	None
Ashish R. Panchal	The Ohio State University Wexner Medical Center	None	None	None	None	None	None	None
Michele M. Pelter	UCSF Physiological Nursing	NIH†; UCSF*	None	None	None	None	Cardinal Health*	None
Marina Del Rios	The University of Chicago Medicine	NIH*	None	None	None	None	None	None
Amber J. Rodriguez	American Heart Association National Center	None	None	None	None	None	None	None
Sarah M. Perman	Yale School of Medicine	NIH/NHLBI†	None	None	None	None	None	None
Stephen Sanko	Keck School of Medicine of USC	None	None	None	None	None	None	None
Pavitra Kotini-Shah	UIC Emergency Medicine	NIH K award†; Zoll Foundation*; WorldPoint*	None	None	None	None	Worldpoint*	Illinois Heart Rescue*

This table represents the relationships of writing group members that may be perceived as actual or reasonably perceived conflicts of interest as reported on the Disclosure Questionnaire, which all members of the writing group are required to complete and submit. A relationship is considered to be "significant" if (a) the person receives \$5000 or more during any 12-month period, or 5% or more of the person's gross income; or (b) the person owns 5% or more of the voting stock or share of the entity, or owns \$5000 or more of the fair market value of the entity. A relationship is considered to be "modest" if it is less than "significant" under the preceding definition.

*Modest.

†Significant.

Appendix 2. Reviewer Disclosures

Reviewer	Employment	Research Grant	Other Research Support	Speakers' Bureau/Honoraria	Expert Witness	Ownership Interest	Consultant/Advisory Board	Other
William Brady	University of Virginia	None	None	None	None	None	None	None
Daniel Davis	Logan Health	None	None	None	None	None	Philips Healthcare*	None
Michael W. Donnino	Beth Israel Deaconess Med Center	NIH (Consultant on NIH funded study on supraglottic airway versus intubation)†	None	None	None	None	None	None
Bryan McNally	Emory University School of Medicine	CDC CARES Expansion and Modernization Grant (Funding for the Cardiac Arrest Registry to Enhance Survival (CARES))†	None	None	None	None	Qilu Hospital†; St Lukes Hospital*; Duke University*; Cedars Sinai Hospital*	None
Robert W. Neumar	University of Michigan	NIH (Research Grants)†; AHA (Research Grant)†; Laerdal Foundation (Research Grant)*	None	None	None	None	None	None
Robert A. Swor	William Beaumont Hospital	None	None	None	None	None	None	None

This table represents the relationships of reviewers that may be perceived as actual or reasonably perceived conflicts of interest as reported on the Disclosure Questionnaire, which all reviewers are required to complete and submit. A relationship is considered to be "significant" if (a) the person receives \$5000 or more during any 12-month period, or 5% or more of the person's gross income; or (b) the person owns 5% or more of the voting stock or share of the entity, or owns \$5000 or more of the fair market value of the entity. A relationship is considered to be "modest" if it is less than "significant" under the preceding definition.

*Modest.

†Significant.