


SE5 (Os06g0603000) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 10 SNPs, MAF>=0.05)




Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

Hd1 (Os06g0275000) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 6 SNPs, MAF>=0.05)





Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

Hd6 (Os03g0762000) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 18 SNPs, MAF>=0.05)



Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).




















OsFKF1 (Os11g0547000) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 52 SNPs, MAF>=0.05)




Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n<=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity by genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (aw p).



Ghd7 (Os07g0261200) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 13 SNPs, MAF>=0.05)




osFTL8 (Os01g0202700) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 22 SNPs, MAF>=0.05)




Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

osFTL1 (Os01g0218500) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 17 SNPs, MAF>=0.05)




Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

osFTL9 (Os01g0748800) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 8 SNPs, MAF>=0.05)



Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).




osFTL6 (Os04g0488400) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 12 SNPs, MAF>=0.05)




Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).


osFTL10 (Os05g0518000) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 7 SNPs, MAF>=0.05)




Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).




osFTL12 (Os06g0552900) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 4 SNPs, MAF>=0.05)



Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).



osFTL11 (Os11g0293800) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 5 SNPs, MAF>=0.05)



Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

rsFTL7 chr12:7,235,920 T>A

