

SE5 (Os06g0603000) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 10 SNPs, MAF>=0.05)


● Green point = photosensitive accession ● Orange point = non-photosensitive accession ■ Lower bar: green = % photosensitive ■ orange = % non-photosensitive



Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).


Hd1 (Os06g0275000) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 6 SNPs, MAF>=0.05)

● Green point = photosensitive accession ● Orange point = non-photosensitive accession ■ Lower bar: green = % photosensitive ■ orange = % non-photosensitive




Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

Hd6 (Os03g0762000) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 18 SNPs, MAF>=0.05)



Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

Hd3a (Os06g0157700) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 18 SNPs, MAF>=0.05)



Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

Ehd1 (Os10g0463400) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 18 SNPs, MAF>=0.05)




Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).






Green point = photosensitive accession Orange point = non-photosensitive accession Lower bar: green = % photosensitive orange = % non-photosensitive






OsGI (Os01g0182600) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 22 SNPs, MAF>=0.05)




Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).




Hd2/OsPRR37 (Os07g0695100) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 31 SNPs, MAF>=0.05)

● Green point = photosensitive accession ● Orange point = non-photosensitive accession ■ Lower bar: green = % photosensitive ■ orange = % non-photosensitive



Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

Ghd7 (Os07g0261200) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 13 SNPs, MAF>=0.05)




Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

osFTL8 (Os01g0202700) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 22 SNPs, MAF>=0.05)




Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

osFTL1 (Os01g0218500) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 17 SNPs, MAF>=0.05)



Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

osFTL9 (Os01g0748800) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 8 SNPs, MAF>=0.05)



Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

osFTL13 (Os02g0232300) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 15 SNPs, MAF>=0.05)




Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

osFTL6 (Os04g0488400) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 12 SNPs, MAF>=0.05)




Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

osFTL10 (Os05g0518000) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 7 SNPs, MAF>=0.05)



Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).


osFTL3(RFT1) (Os06g0157500) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 17 SNPs, MAF>=0.05)



Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).


osFTL12 (Os06g0552900) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 4 SNPs, MAF>=0.05)

● Green point = photosensitive accession ● Orange point = non-photosensitive accession ■ Lower bar: green = % photosensitive ■ orange = % non-photosensitive




Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

osFTL4 (Os09g0513500) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 20 SNPs, MAF>=0.05)




Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

osFTL11 (Os11g0293800) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 5 SNPs, MAF>=0.05)



Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).

osFTL7 (Os12g0232300) — flowering date and photosensitivity by genotype of SNPs in the gene (n=278 accessions; 24 SNPs, MAF>=0.05)



Each point = one accession (green = photosensitive, orange = non-photosensitive); box = quartiles, thick line = median; violin if n>=10, box if n>=3; ALT = allele differing from Nipponbare reference. ANOVA/KW on flowering date over genotype groups with n>=3; lower bar = fraction photosensitive (green); chi-square/Fisher/permutation test of photosensitivity vs genotype. Unadjusted for population structure (exploratory). Stars on ANOVA: * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns = not significant (raw p).