


The result of each trial (0 : failure ;1: success):

$$X_i \in \{0, 1\} \sim \text{Bernoulli}(p), p \in [0, 1]$$

Bernoulli Process ($t; p$)

Parameters are discrete, states are discrete



The total number of successes in n trials

$$Y_i \in \{0, 1, 2, \dots, n\} \sim \text{Binomial}(n, p), p \in [0, 1], n \in \mathbb{Z}_{>0}$$

The number of trials between $(i-1)^{th}$ and i^{th} success:

$$N_i \in \{1, 2, \dots\} \sim \text{Geometric}(p), p \in [0, 1]$$

The number of trials to get the k^{th} success:

$$W_k \in \{1, 2, \dots\} \sim \text{NegativeBinomial}(p, k), p \in [0, 1], k \in \mathbb{Z}_{>0}$$

$$\text{PMF } p_{X_i}(x) = p^x(1-p)^{(1-x)} \text{ or } px + (1-p)(1-x), x \in \{0, 1\}$$

$$F_{X_i}(x) = \begin{cases} 0, & x < 0 \\ 1-p, & x \in [0, 1] \\ 1, & x \geq 1 \end{cases}$$

$$\mu_{X_i} = p, \sigma_{X_i} = \sqrt{np(1-p)}$$

$$\text{PMF } p_Y(y) = C_n^y p^y (1-p)^{n-y}$$

$$C_n^y = \frac{n!}{y!(n-y)!}$$

$$\mu_Y = p, \sigma_Y = \sqrt{np(1-p)}$$

$$\text{PMF } p_{N_i}(n) = p(1-p)^{n-1}$$

$$\text{CDF } F_{N_i}(n) = 1 - p(1-p)^n$$

$$\mu_{N_i} = \frac{1}{p}, \sigma_{N_i} = \frac{\sqrt{1-p}}{p}$$

$$\text{PMF } p_{W_k}(w) = C_{w-1}^{k-1} p^k (1-p)^{w-k}$$

$$\text{CDF } F_{W_k}(w) = \sum_{i=1}^w p_{W_k}(i)$$

$$\mu_{W_k} = \frac{k}{p}, \sigma_{W_k} = \frac{\sqrt{k(1-p)}}{p}$$