Minimum Widths for Unit Squares in Thin Rectangles

Hiroaki Hosono (working draft)

25 October 2026

Abstract

We present a computer-assisted determination of the minimum width needed to pack N unit squares, with arbitrary rotations and pairwise disjoint interiors, in a rectangle of height $1851/1000 \le a < 2$. For odd $N \ge 11$, the width is $4+2(N-2)\sqrt{2}-(N-1)a$. For even $N \ge 12$, it is the minimum of an explicit rationally parametrised function, attained by a staggered configuration with two upright end caps. The representative minimising angle is unique. The proof separates a finite certificate restricting every candidate packing to a specified configuration class from analytical minimisation and attainment. We describe the exact acceptance obligations and the retained two-system evidence.

1. Introduction and notation

Let $\nu(x,a)$ be the maximum number of closed unit squares contained in $[0,x]\times[0,a]$ with pairwise disjoint interiors, allowing arbitrary orientations. Contacts are permitted. Write

$$B_N(a) = \inf\{x > 0 : \nu(x,a) \ge N\}.$$

Friedman's rectangle problem [1] records constructions and boundary pieces for small numbers of squares. Related counting bounds for unit squares in rectangles are studied in [2,3]. The lower bound in the present paper uses the finite verification and analytical argument described below.

Our question fixes a height strictly below two and determines the width uniformly in the number of squares.

No priority claim is made; a broader literature search remains future work.

2. Main theorem

Measure θ counterclockwise from the positive horizontal axis to a side direction of a unit square. Orientations differing by $\pi/2$ describe the same square. Represent an orientation by $t = \tan(\theta/2)$ and let

$$\mathcal{O} = \left[2\arctan\frac{4339}{12800},\ 2\arctan\frac{4553}{8800}\right].$$

For $\theta \in \mathcal{O}$, put $c = \cos\theta$, $s = \sin\theta$, $h = c + s$, $b = 3a/2 - 1$, and

$$P = \frac{2h-a}{h^2-1}, \qquad Q = 1 + \frac{2h-b}{h^2-1}, \qquad A = \frac{2-a}{4}(\tan\theta - \cot\theta).$$

Define

$$G_N(\theta; a) = (N-4)P + 2Q - 2|A|.$$

All denominators are positive on this angle interval. The symbol h denotes a square's bounding-box side length, whereas a is the container height.

Theorem 2.1 (exact thin-strip width; computer-assisted). For every $1851/1000 \le a < 2$:

1. For odd $N \ge 11$,
$$B_N(a) = 4 + 2(N-2)\sqrt{2} - (N-1)a.$$
2. For even $N \ge 12$,
$$B_N(a) = \min_{\theta \in \mathcal{O}} G_N(\theta; a).$$
The representative minimiser $\theta_N(a) > \pi/4$ is unique and lies below $2\arctan(43/100)$. Its reflection $\pi/2 - \theta_N(a)$ gives the other minimising orientation. The value is attained by a staggered packing.

Both formulas are attained by two upright end caps and $N-2$ interior squares at one common angle, placed on alternating walls. For odd N, the two caps touch the same wall and the optimal common angle is $\pi/4$. For even N, the caps touch opposite walls and the optimal angle is selected by the one-variable minimum in the theorem. Figure 1 shows examples of this general construction; the proof of attainment for every size is given in Section 5.

An interactive illustration in the online version allows the number of squares and the height to vary. Decimal widths and angles in that illustration are floating-point approximations; the exact claims are the formulas above. The finite range of the controls does not limit the theorem. The illustration displays the attaining family and is not a proof computation.

The theorem combines the retained two-system results on $[1851/1000, 19/10]$ and $[19/10, 2)$, including the separate coverage for $N = 11$. The mathematical argument is given below, with its finite proof obligations exposed in Section 6.

3. Certified restriction to a configuration class

First order the squares by nondecreasing horizontal centre coordinate; ties may be ordered arbitrarily. Use square symmetry to represent each angle by $0 \le \theta \le \pi/2$, with $t = \tan(\theta/2)$. In this representation, **near-upright** means

$$t \in \widehat{\mathcal{U}} = [0, 7/256] \cup [125/132, 1].$$

For a square with centre height y and bounding-box side $h_\theta = \cos\theta + \sin\theta$, define its normalised height

$$\rho = \frac{y - h_\theta/2}{a - h_\theta} \in [0, 1].$$

The denominator is positive in the theorem range. The closed lower and upper height labels are $\rho \in [0, 1/2]$ and $\rho \in [1/2, 1]$. A midpoint may receive either label in the closed cover. A **sequence in the configuration class** (a *type sequence* in the source records) has the half-angle parameters of its two end squares in $\widehat{\mathcal{U}}$, every interior angle in $\mathcal{O}$, and alternating height labels along all consecutive pairs, including the pairs at the caps. The class restriction is the assertion that every packing at or below the threshold stated next has such a type sequence.

Put $g = 2\sqrt{2} - a$ and $F_0 = 4 - 2\sqrt{2}$. The retained certificate establishes the following assertion for every $N \geq 11$: a packing of width at most

$$(N-1)g + F_0$$

has two near-upright end caps, its $N-2$ interior squares have angles in $\mathcal{O}$, and the centres of the interior squares alternate between the closed lower and upper halves of the container (the midpoint belongs to both). Every sequence outside the class has strictly greater width.

This is an all-sequence assertion. The verification covers angle/height cells and all transitions, not just configurations found in a search. The separating-axis theorem (SAT), which tests disjointness by projections onto edge normals, gives exact bounds for transitions outside the class. Rational potentials, found by linear programming (LP) and checked in exact arithmetic, redistribute these bounds along a sequence. Calibrated transitions within the class use angular variation. A saturated dynamic-programming (DP) layer and its checked fixed point cover arbitrarily large N.

The extension from finitely many sizes to every larger size is an induction, not an inference from sampled values. After the chosen edge layer, consider the finite augmented state graph, including the flags and previous-edge information needed by the transition rules. Let b_z be the bound at entry to the saturated layer, $w_{zz'}$ a verified transition cost, and V_z the final saturated bound. The exact fixed-point check establishes

$$V_z \leq b_z, \qquad V_{z'} \leq V_z + w_{zz'} \quad \text{for every legal transition } z \to z'.$$

At entry the accumulated path cost is at least V_z. If that holds at one state, adding the next verified transition cost proves it at the next state. Thus it holds after any number of further transitions. Adding each legal terminal bound gives the required residual-width bound for arbitrarily long sequences outside the class. The start and end conventions convert edge counts to square counts; their two end rules are why a saturated edge layer must not be labelled as a finite count of squares. The finite sizes before this tail are checked separately, as listed in Section 6. Every legal transition, including entry into a run of compatible interior states, must be present for this induction to apply. This argument uses the checked finite vector inequalities, and does not assume that all chord-transition costs are nonnegative.

The threshold is sufficient for the equality theorem as follows. The 45° construction has width $(N-1)g + F_0$. For even N, $G_N(\pi/4; a)$ equals that threshold, so $\min G_N$ is no larger. An arbitrary packing wider than the threshold already satisfies the claimed lower bound. Any packing at or below the threshold is restricted to this class, where Section 4 applies. Section 5 constructs a packing attaining the resulting value. Thus the class restriction alone is not being substituted for minimisation or attainment.

The finite verification includes transitions into runs of compatible interior states as well as transitions within those runs. Near height two it uses a corrected absolute-value inequality for nearly upright squares. Earlier computations with the superseded signed inequality are excluded from the evidence.

4. The lower bound within the configuration class

4.1. Pair and cap inequalities

For interior angles $\theta_1, \ldots, \theta_{N-2}$ define

$$\mathrm{TV} = \sum_{i=1}^{N-3} |\theta_{i+1} - \theta_i|.$$

The retained pair calibration gives a lower bound per radian of θ. Its constant can be chosen as

$$k = \frac{1}{5} \quad \left(\frac{1851}{1000} \le a \le \frac{19}{10}\right), \qquad k = \frac{227}{2000} \quad \left(\frac{19}{10} \le a < 2\right).$$

Here is the summed inequality explicitly. Put $r = N - 2$ and label the interiors from left to right. The cap centres have horizontal coordinates x_0 and x_{r+1}. Let $\sigma_0 = +1$ when the first cap-to-interior step goes from a lower-labelled square to an upper-labelled one, and $\sigma_0 = -1$ for the reverse step. Set

$$\sigma_i = (-1)^i \sigma_0 \quad (0 \le i \le r), \qquad d_i = x_{i+1} - x_i.$$

The gap functions $f_\pm$ and $c_\pm$ are defined explicitly in Section 5. The calibrated interior-pair inequality is

$$d_i \ge \frac{f_{\sigma_i}(\theta_i) + f_{\sigma_i}(\theta_{i+1})}{2} + k|\theta_{i+1} - \theta_i| \quad (1 \le i < r).$$

Thus calibration means a uniform strengthening of the common-angle gap bound by a term proportional to angle difference. If h_L, h_R are the bounding-box side lengths of the original two caps, the cap comparison gives

$$\frac{h_L}{2} + d_0 \ge \frac{1}{2} + c_{\sigma_0}(\theta_1), \qquad d_r + \frac{h_R}{2} \ge c_{\sigma_r}(\theta_r) + \frac{1}{2}.$$

The identities

$$P = \frac{f_+ + f_-}{2}, \qquad T_\sigma = \frac{1}{2} + c_\sigma + \frac{1}{2} f_{-\sigma} = Q + \sigma A$$

then give the occupied-width bound

$$W \ge \sum_{j=2}^{r-1} P(\theta_j) + Q(\theta_1) + Q(\theta_r) + \sigma_0 A(\theta_1) + \sigma_r A(\theta_r) + k\,\mathrm{TV}.$$

Because $\sigma_r = (-1)^r \sigma_0$, the endpoint signs agree for even N and are opposite for odd N. This is the sign rule used in the next subsection. The same certified bounds give $|\partial_\theta A| \le k$.

The cap comparison is applied to pairs at their original positions. It does not assume that simultaneously moving arbitrary caps preserves a packing. The lower-height cap comparison is a separate certified dependency; the upper-height argument must not be extrapolated below 1.9 without it.

4.2. Reduction to a common angle

Choose an interior angle $\bar{\theta}$ closest to $\pi/4$. Both P and Q decrease as h increases. Indeed,

$$\frac{d}{dh}\frac{2h-z}{h^2-1} = -\frac{2(h^2-zh+1)}{(h^2-1)^2} < 0$$

for $z = a$ or $z = b$, since $z < 2$ in the claimed range. Consequently each P- and Q-term is at least its value at $\bar{\theta}$.

For even N, the two endpoint A-terms have the same sign pattern. Replacing them by $-2|A(\bar{\theta})|$ loses at most

$$k(|\theta_1 - \bar{\theta}| + |\theta_{N-2} - \bar{\theta}|) \le k\,\mathrm{TV}.$$

The variation contribution absorbs this loss. The occupied width is therefore at least $G_N(\bar{\theta}; a)$, and hence at least its minimum.

For odd N, the endpoint A-terms have opposite signs. Their difference is absorbed by $k\,\mathrm{TV}$. The remainder is minimised when $h = \sqrt{2}$, giving

$$(N-4)P(\pi/4) + 2Q(\pi/4) = 4 + 2(N-2)\sqrt{2} - (N-1)a.$$

4.3. Uniqueness and the critical equation

In this paragraph differentiation is with respect to t, not per radian of θ. On

$$I = [\sqrt{2} - 1, 4553/8800],$$

the retained exact polynomial checks establish

$$P_{tt} > 0, \qquad P_{tt} + 2Q_{tt} - 2A_{tt} > 0.$$

On the open upper-angle side $t > \sqrt{2} - 1$, $A \ge 0$, and

$$G_{N,tt} = (N-5)P_{tt} + (P_{tt} + 2Q_{tt} - 2A_{tt}) > 0.$$

The full function has an absolute-value cusp at $t_0 = \sqrt{2} - 1$. The derivative here is the right derivative of the upper-angle branch:

$$G_{N,t}(t_0+; a) = -(2-a)(2+\sqrt{2}) < 0,$$

while the checked derivative at $t = 43/100$ is positive for $N \ge 12$. This yields a unique representative critical point below that endpoint. The reflection of the lower-angle part of $\mathcal{O}$ lies in the checked upper interval, so it adds no smaller value. Reflection gives the other minimising orientation.

The rational parametrisation of c, s turns the critical-point equation into a polynomial equation in t, with rational polynomial coefficients in a, after multiplication by positive denominators. The minimising parameter is therefore an algebraic function of a. When a is algebraic, its parameter and minimum width are algebraic numbers. For an arbitrary real a, no algebraicity

over the rationals is asserted, and the angle $2\arctan t$ is not being called an algebraic number. Convexity and endpoint signs here are certificate dependencies, not deductions from a plot.

5. Attaining configurations

Use two upright caps and $N-2$ interiors at one angle, on alternating walls. Let

$$f_+ = \frac{1-s(a-h)}{c}, \qquad f_- = \frac{1-c(a-h)}{s},$$

$$c_+ = \frac{(1+h)/2 - s(a-(1+h)/2)}{c}, \qquad c_- = \frac{(1+h)/2 - c(a-(1+h)/2)}{s}.$$

Interior centres have height $h/2$ or $a-h/2$. Each cap has centre height $1/2$ or $a-1/2$, on the wall opposite its neighbour. Starting with cap centre $x = 1/2$, take consecutive horizontal increments $c_\pm$ at the caps and $f_\pm$ between interiors, according to the alternating wall pattern.

Adjacent pairs touch on an appropriate face normal. On the angle interval containing every minimiser, the retained attainment checks establish

$$f_\pm > h/2, \qquad c_\sigma + f_{\sigma'} > (1+h)/2, \qquad c_\sigma > (h-1)/2$$

for the required sign choices. The first two inequalities separate nonadjacent pairs horizontally; longer chains add positive increments. The last makes the caps outermost. The cap–cap pair is also separated, since a cap increment and two interior increments already exceed one.

Containment follows from $h \le \sqrt{2} < a$. Thus the width is exactly one plus the sum of the horizontal increments. For even N, choose the starting wall to obtain

$$1 + 2c_{\sigma_0} + f_{-\sigma_0} + (N-4)(f_+ + f_-)/2 = (N-4)P + 2Q - 2|A|.$$

For odd N, use $\theta = \pi/4$; the same sum is the formula in Theorem 2.1. Choosing the critical orientation in the even case gives exact attainment. The alternating sequence puts the two upright caps on the same wall for odd numbers of squares and on opposite walls for even numbers (Figure 1).

For fixed t, the individual expressions in the analysis and attainment checks are affine in a. The accepted endpoint certificates therefore extend pointwise through the two height intervals. This is applied to individual inequalities, not to a presumed endpoint extremum of an arbitrary minimum of affine branches.

A nearby rational-half-angle placement certificate may have a small positive width slack. Such a certificate is evidence of a packing near the optimum, not equality of its width with the critical value.




Figure 1. The construction of Section 5 at height 1.9: eleven squares at 45 degrees (the odd optimum), and twelve squares at the illustrative rational half-angle parameter 427/1000. Orange squares are the two upright end caps; blue squares alternate between the walls. The even panel illustrates a feasible construction at the displayed angle, not the exact critical-angle optimum. Coordinates are rounded only for drawing; the figure is not a packing certificate.




Figure 2. Height and width on the theorem interval. The odd curve is the exact formula for eleven squares. The even curve is the width of the twelve-square construction at the fixed half-angle parameter 427/1000; it is an upper bound, while the theorem takes the minimum over all admissible angles. The right endpoint at height two is excluded. Lines are drawn from the stated formulas and do not establish the finite lower-bound computation.

6. Exact computations and reproducibility

Appendix A lists the requirements for independent reproduction. The mathematical coverage and the obligations for a complete verification are set out here.

6.1. Verification coverage

The lower-height verification records document checks by a first implementation and a specification-derived second implementation, followed by mathematical review. It reports complete interval and Sturm checks separately from the 36 finite placement tests. Those placements are not a proof of all heights or all N.

For $a \in [1851/1000, 1852/1000]$, the combined finite-layer and tail verification establishes $N \geq 24$ using its individual finite layers together with its saturated tail; the first-system output labels that tail $N \geq 26$, rather than $N \geq 24$. Separate fixed-N checks cover $11 \leq N \leq 26$. Their union covers every $N \geq 11$. On $[1852/1000, 1854/1000]$, the chord runs cover $N \geq 12$ and separate runs cover $N = 11$. Later-height runs cover the remainder. The corrected absolute-form final interval and the separate $N = 11$ subdivision at height 1.9 must be included.

The verification records document that the second implementation was written from the specifications without using the first implementation's code.

6.2. Acceptance obligations

A complete replay must check the closed state cover, all SAT branches, potential coverage, every transition (including entry to and exit from runs of compatible interior states), the integer rounding model, and the saturated-layer fixed point. It must also check the full-domain calibration and cap inequalities, the polynomial signs and root counts, the affinity argument and the attainment inequalities. Positive output on selected intervals, floating LP feasibility, placement samples or a root count alone are insufficient.

6.3. Availability of the computational supplement

The implementations use Python 3, SymPy and NumPy. A self-contained computational supplement must specify their versions, the complete source code, and the identities of every accepted input and output. That version-pinned supplement is not yet available with this paper. The public paper and its TeX bundle provide the exposition and figures. The complete verification-code and input-data archive remains to be released. Appendix A states what a complete archive must contain and what each verification must establish.

7. Scope and computational supplement

The theorem covers every fixed height $1851/1000 \leq a < 2$ and every integer $N \geq 11$, with the odd and even formulas stated in Theorem 2.1. The left height endpoint is included and height two is excluded. Other heights and smaller values of N are outside the statement.

This is a computer-assisted working paper and has not been externally peer reviewed. The assembled text has received a limited independent consistency review against the verification records. A self-contained, version-pinned computational supplement and a complete replay inventory remain to be released. Preparing the text and illustrations did not include a new execution of the proof computations.

8. References

[1] Erich Friedman, *Problem of the Month (June 2010)*, Math Magic. https://erich-friedman.github.io/mathmagic/0610.html.

[2] Hiroshi Nagamochi, *Packing Unit Squares in a Rectangle*, The Electronic Journal of Combinatorics **12** (2005), Research Paper R37. https://doi.org/10.37236/1934.

[3] Hakan Karakuş, *A counterexample to Nagamochi's scoring lemma and a new rectangle packing bound*, arXiv:2609.37410v1 (2026). https://arxiv.org/abs/2609.37410v1.

Appendix A. Requirements for independent reproduction

A.1. Verification obligations

A complete computational supplement must connect each proof ingredient to a precisely specified verification and an unambiguous set of accepted input and output data. The following obligations describe the mathematical content; the complete archive has not yet been released.

Ingredient	Required verification
state cover, exact bounds, DP and saturation rules	prove closed coverage, sound rounding and the fixed-vector inequalities
upper-height analysis and attainment	verify polynomial signs and nonintersection throughout the stated intervals
upper-height coverage, including the final interval and $N = 11$	record both the exceptional finite cases and the arbitrarily large tail
lower-height configuration-class restriction	check every legal transition, including entries into interior runs
lower-height calibration and cap comparison	verify the quantitative pair inequalities and cap bounds at original positions
lower-height analysis and attainment	verify the angle interval, affine-height extension and all required separation bounds
upper-height calibration	verify the common-angle and angle-variation bounds uniformly

A.2. Archive and replay requirements

The released archive must specify the full source code and dependency versions, the exact state covers and potential data, all accepted inputs and outputs, the height intervals and finite-count ranges, and the checked saturation conditions. It must document how the two implementations were developed independently and identify which output establishes each proof obligation.

Reproduction must cover all separating-axis branches, all legal transitions and terminal rules, exact integer rounding, every fixed-point inequality, the analytical polynomial checks, and the attaining configurations. A selection of examples or floating-point optimization results does not replace these checks.

In particular, fixed-N checks do not replace the separately required saturation argument. The final height interval $[1.999, 2)$ requires the absolute-value form of the near-upright inequality. These requirements must be explicit in the released verification data.