

Why $K = 0.95$: the physics behind the sub- $\pi/4$ area trick

A first-principles derivation of the finite- V_{ct} correction to the OR-protocol pulse area.

TriQG examples/Smaller_VDD_energy · 2026-05-15

Punchline. The protocol’s “area = $\pi/4$ ” was derived in the $V_{ct} \rightarrow \infty$ limit, where the EIT shield Ω_R couples $|P\rangle$ to a fully-decoupled $|R\rangle$. At finite V_{ct} , $|P\rangle$ and $|R\rangle$ **hybridize**. The probe still drives only $|P\rangle$, but now sees $|P\rangle$ -amplitude distributed across two dressed states at different energies. The effective two-photon Raman rate on $|+\rangle = (|A\rangle + |B\rangle)/\sqrt{2}$ is **enhanced** by a few percent. To hit the design flip phase $\varphi_+ = \pi$ exactly, the pulse area must be under-rotated by the same few percent. At rev. 4 parameters that comes out to $K_{\text{opt}} = 0.93$; the empirical 0.95 is 2% off due to higher-order corrections and the decoherence/gate-time trade-off.

1 The clean limit and what changes

The OR protocol acts on a target $|A\rangle, |B\rangle$ via an off-resonant Raman through $|P\rangle$ (detuning Δ) with EIT shielding Ω_R on $|P\rangle \leftrightarrow |R\rangle$. Because the probe drives both $|A\rangle \leftrightarrow |P\rangle$ and $|B\rangle \leftrightarrow |P\rangle$ with the same Ω_p , the target lives in a $|+\rangle / |-\rangle$ basis:

$$|+\rangle = (|A\rangle + |B\rangle)/\sqrt{2} \quad (\text{bright, coupling } \sqrt{2}\Omega_p \text{ to } |P\rangle), \quad (1)$$

$$|-\rangle = (|A\rangle - |B\rangle)/\sqrt{2} \quad (\text{dark, decoupled}). \quad (2)$$

A π rotation of $|+\rangle$ relative to $|-\rangle$ swaps $|A\rangle$ and $|B\rangle$ (the flip). In **blockaded** cells (≥ 1 control in $|r\rangle$), $|R\rangle$ is shifted up by V_{ct} , so the target effectively becomes a 3-level system:

$$\underbrace{|+\rangle}_{E_+=0} \xleftrightarrow{\sqrt{2}\Omega_p} \underbrace{|P\rangle}_{E_P=\Delta} \xleftrightarrow{\Omega_R} \underbrace{|R\rangle}_{E_R=V_{ct}}. \quad (3)$$

In the **clean limit** $V_{ct} \rightarrow \infty$, $|R\rangle$ is fully decoupled, the $|+\rangle \leftrightarrow |P\rangle$ Raman runs with rate $\Omega_p^2/(2\Delta)$, and integrating to area $\pi/4$ (in the code’s convention) gives $\varphi_+ = \pi$ exactly. That is the **design**.

2 Dressing of $|P\rangle$ at finite V_{ct}

At finite V_{ct} , the $(|P\rangle, |R\rangle)$ sub-block of the Hamiltonian

$$H_{PR} = \begin{pmatrix} \Delta & \Omega_R/2 \\ \Omega_R/2 & V_{ct} \end{pmatrix} \quad (4)$$

has **exact** eigenvalues and eigenvectors

$$\varepsilon_{\pm} = \frac{\Delta + V_{ct}}{2} \pm \sqrt{\frac{(V_{ct} - \Delta)^2}{4} + \frac{\Omega_R^2}{4}}, \quad \tan 2\theta = \frac{\Omega_R}{V_{ct} - \Delta}. \quad (5)$$

The two dressed states $|\tilde{P}_{\pm}\rangle$ at energies $\varepsilon_{\pm}$ each carry a fraction of $|P\rangle$ character, $\cos^2 \theta$ and $\sin^2 \theta$ respectively. The probe still only couples to bare $|P\rangle$, so its matrix element to each dressed state is reduced by that overlap:

$$\langle + | H | \tilde{P}_{\pm} \rangle = \frac{\Omega_p}{\sqrt{2}} \langle P | \tilde{P}_{\pm} \rangle. \quad (6)$$

3 Effective Raman rate on $|+\rangle$

The AC Stark shift on $|+\rangle$ – which **is** the phase-accumulation rate that flips $|A\rangle \leftrightarrow |B\rangle$ – is the sum of the two-level contributions from both dressed states:

$$\Omega_{\text{eff}} = \frac{\Omega_p^2}{2} \left[\frac{|\langle P | \tilde{P}_-\rangle|^2}{\varepsilon_-} + \frac{|\langle P | \tilde{P}_+\rangle|^2}{\varepsilon_+} \right]. \quad (7)$$

In the $V_{ct} \rightarrow \infty$ limit, $\varepsilon_- \rightarrow \Delta$, $|\langle P | \tilde{P}_-\rangle|^2 \rightarrow 1$, the upper term dies, and Equation 7 collapses to $\Omega_p^2/(2\Delta)$. That is the $\pi/4$ design baseline.

At finite V_{ct} , Equation 7 differs from the baseline by a **parameter-dependent** factor. The pulse area is calibrated assuming the baseline; the actual phase rotation is then larger (or smaller) by that factor. To hit $\varphi_+ = \pi$ exactly, the area must be scaled by the **inverse** of that factor:

$$K_{\text{opt}} = \frac{\Omega_p^2/(2\Delta)}{\Omega_{\text{eff}}} = \frac{1}{1 + (V_{ct}\text{-correction})}. \quad (8)$$

4 Numerical check at rev. 4

Plug in rev. 4 FINAL parameters $\Omega_p = 2\pi \times 50$ MHz, $\Omega_R = 2\pi \times 175$ MHz, $\Delta = 2\pi \times 500$ MHz, $V_{ct} = 2\pi \times 226$ MHz:

Lower eigenvalue ε_-	200.4
Upper eigenvalue ε_+	525.6
$ P\rangle$ -fraction of $ \tilde{P}_-\rangle$	0.079
$ P\rangle$ -fraction of $ \tilde{P}_+\rangle$	0.922
$\Omega_{\text{eff}}/(\Omega_p^2/2)$	$0.079/200.4 + 0.922/525.6 = 2.149 \times 10^{-3}$
Ideal $1/\Delta$	2.000×10^{-3}
Ratio (enhancement)	1.074 (+7.4%)
Predicted $K_{\text{opt}} = 1/1.074$	0.931

Table 1: Exact diagonalization of H_{PR} at rev. 4 parameters and the corresponding Ω_{eff} via Equation 7. Numbers are in MHz $/(2\pi)$.

The full brute-force / parameter-scan optimum is $K = 0.95$ (rev. 4 report, Fig. 1). **Theory and experiment agree to 2%.**

5 Where does the enhancement come from physically?

The non-obvious feature: at rev. 4, $V_{ct} = 226 < \Delta = 500$. The $|R\rangle$ state sits **below** $|P\rangle$ in energy. Dressing by Ω_R pushes the bare $|P\rangle$ **up** (to $\varepsilon_+ = 525.6$, above Δ) and creates a **low-energy** dressed state at $\varepsilon_- = 200.4$ that carries 7.9% of $|P\rangle$ character.

The probe sees this distribution and gets a Stark shift from **both** dressed states. The high-lying $|\tilde{P}_+\rangle$ contributes slightly **less** than the ideal $|P\rangle$ would ($0.922/525.6 < 1/500$), but the low-lying $|\tilde{P}_-\rangle$ contributes a substantial $0.079/200.4 \approx 1/2540$ on top. The low-energy contribution **wins**: total Raman rate +7%.

In one sentence. When $V_{ct} < \Delta$, the EIT-dressed $|R\rangle$ effectively donates a fraction of $|P\rangle$ character to a low-energy state, and the probe gets a cheaper Raman process through it.

6 The 2% gap between $K = 0.93$ and $K = 0.95$

The leading-order prediction is $K_{\text{opt}} = 0.93$ at rev. 4, but the empirical optimum is $K = 0.95$. Three corrections explain the gap:

1. **Higher-order in Ω_p .** The Schrieffer-Wolff treatment is second order in Ω_p ; the next correction is $\sim \Omega_p^4/\Delta^3$, which at rev. 4 is order $(50)^4/(500)^3 \cdot (2\pi) \approx 0.05$ MHz – non-negligible at the 10^{-3} infidelity level.
2. **Pulse shaping.** The super-Gaussian envelope makes $\Omega_p^2(t)/\Delta_{\text{eff}}(t)$ vary through the pulse. The correction factor is **instantaneous**, so the area-weighted average differs slightly from the peak value used above.
3. **Decoherence trade-off.** Shorter pulses (smaller K , since $T_f \propto K$ at fixed α) suffer less Cs $|r\rangle$ decay. The fidelity optimum has to balance phase-mismatch infidelity $\sim \delta^2/4$ against the linear decay loss, which shifts K_{opt} slightly **upward** from the pure phase-matching value.

7 Sign flip at $\Delta \approx V_{ct}$: why $K = 1$ wins at the small- Δ champion

The brute-force $K = 1$ champion sits at $\Delta = 400$ MHz, $V_{ct} = 442$ MHz – **strong-mixing** regime where $|\Delta - V_{ct}| = 42$ MHz is smaller than $\Omega_R/2 = 90$ MHz. Perturbation theory in Ω_R breaks down; exact diagonalization gives:

ε_- ($ P\rangle$ -like, since $\Delta < V_{ct}$)	328.6
ε_+ ($ R\rangle$ -like)	513.4
$ P\rangle$ -fraction of $ \tilde{P}_-\rangle$	0.615
$ P\rangle$ -fraction of $ \tilde{P}_+\rangle$	0.385
$\Omega_{\text{eff}}/\Omega_{\text{ideal}}$	1.049 (+4.9%)
Predicted K_{opt}	0.954

Table 2: Exact diagonalization of H_{PR} at the $K = 1$ champion cell ($\Delta = 400$, $V_{ct} = 442$, $\Omega_R = 180$ MHz $/(2\pi)$). The dressed states are **symmetrically split** by ~ 92 MHz around the centroid because $|\Delta - V_{ct}|$ is small.

The two dressed states are roughly **symmetric** around the bare-energy centroid: one pushed down to 328.6 (P-like), one pushed up to 513.4 (R-like, but heavily mixed). Their contributions to Equation 7 partially cancel – the net enhancement is only 4.9%, much smaller than the 7.4% at rev. 4.

A 5% phase mismatch at $K = 1$ costs $\sin^2(0.05\pi/2) \approx 6 \times 10^{-3}$ in the $|1, 1, * \rangle$ branch. A $K = 0.95$ pulse has 5% longer T_f , costing roughly $2 \cdot 0.05 \cdot 220$ ns $/77 \mu\text{s} \approx 3 \times 10^{-4}$ additional Cs decay. Within 10^{-3} fidelity differences, **the two effects nearly cancel**, and the small residual favors $K = 1$ in this cell ($\Delta < V_{ct}$, weaker enhancement).

Bottom line. The “optimal K ” is not a protocol-level constant; it is a function $K_{\text{opt}}(\Delta, \Omega_R, V_{ct})$ that the brute-force sweep measures empirically. The closed-form prediction is

$$K_{\text{opt}} = \left[(\Omega_p^2/2) \left[\frac{|\langle P | \tilde{P}_-\rangle|^2}{\varepsilon_-} + \frac{|\langle P | \tilde{P}_+\rangle|^2}{\varepsilon_+} \right] / (\Omega_p^2/(2\Delta)) \right]^{-1} \quad (9)$$

with the exact-eigenvalue $\varepsilon_{\pm}$ and overlaps from $H_{PR} = \begin{pmatrix} \Delta & \Omega_R/2 \\ \Omega_R/2 & V_{ct} \end{pmatrix}$.

At rev. 4: $K_{\text{opt}} = 0.93$ (theory) vs 0.95 (empirical). At the $K = 1$ champion cell: $K_{\text{opt}} = 0.95$ (theory) vs numerically a near-tie between $K = 0.95$ and $K = 1$ where decoherence trade-offs tip the balance to $K = 1$.

The “0.95” is not a fundamental constant – it is a **good compromise value** near the rev. 4 cell that survives the empirical optimization, and the brute force shows it generalizes to a wide plateau of cells where $\Delta > V_{ct}$ holds.