1. OR Gate Probe Pulse Shape Comparison

Cubic Super-Gaussian vs Hanning Window

1.1. Pulse Shape Definitions

The two-photon effective pulse area, which governs the target qubit's Rabi cycle, is defined as

$$\text{area} = \int_{T_c}^{T_c+2T_f} \frac{\left(\Omega_p(t)\right)^2}{2\delta}\,\mathrm{d}t \tag{1}$$

where δ is the single-photon detuning from $|P\rangle$. Both pulse shapes below are tuned so that area $= \pi$.

1.1.1. Cubic Super-Gaussian (Current)

$$\Omega_p^{(\mathrm{CSG})}(t) = \Omega_{p,0} \exp\left(-\left(\frac{(t-t_c)^3}{\sigma}\right)^2\right), \quad t \in \left[T_c, T_c + 2T_f\right] \tag{2}$$

where $t_c = T_c + T_f$ is the pulse center. The cubic exponent $(t-t_c)^3$ produces an asymmetric envelope that rises more steeply on the leading edge.

Coefficients for area $= \pi$ (with T_f enlarged so edges $< 0.05\%$ of peak):

Parameter	**Value**
$\Omega_{p,0}/2\pi$	50 MHz
$\delta/2\pi$	500 MHz
σ	$1.771 \times 10^{-3}\ \mu s^3$
T_f	0.17 μs
Probe duration $2T_f$	340 ns
Total gate time	360 ns
Edge amplitude	$< 0.05\%$ of peak

1.1.2. Hanning (sin²) Window

$$\Omega_p^{(\mathrm{Han})}(t) = \Omega_{p,0} \sin^2\left(\pi \frac{t-T_c}{2T_f}\right), \quad t \in \left[T_c, T_c + 2T_f\right] \tag{3}$$

The Hanning window is symmetric, smoothly starts from exactly zero and returns to exactly zero at both edges. Its area efficiency (fraction of the rectangular-pulse area) is $3/8 = 37.5\%$.

Coefficients for area $= \pi$:

Parameter	**Value**
$\Omega_{p,0}/2\pi$	50 MHz
$\delta/2\pi$	500 MHz
T_f	0.2667 μs
Probe duration $2T_f$	533 ns
Total gate time	553 ns

1.2. Pulse Envelopes

Figure 1 overlays the two pulse envelopes, both tuned to area $= \pi$. Both pulses start and end at zero (or negligibly close). The cubic super-Gaussian reaches peak amplitude faster but has an asymmetric shape due to the cubic exponent. The Hanning pulse is wider (533 vs 340 ns) but is perfectly symmetric with smooth transitions at both boundaries.



Figure 1: Probe pulse envelopes for the cubic super-Gaussian (blue, 340 ns) and Hanning $\sin^2$ window (orange, 533 ns). Both are tuned to area $= \pi$ with $\Omega_{p,0}/2\pi = 50$ MHz and $\delta/2\pi = 500$ MHz. Both pulses start and end at zero.

1.3. Average Gate Fidelity

Following Yu *et al.* (arXiv:2203.14302, Eq. 7), the average gate fidelity is computed over all $2^{n+1} = 8$ computational basis states:

$$\bar{\mathcal{F}}_n = \frac{1}{2^{n+1}} \sum_{k=1}^{2^{n+1}} F\left(\rho_{\text{out}}^{(k)}, \rho_{\text{et}}^{(k)}\right) \tag{4}$$

For the OR gate, the ideal output equals the input for all basis states (the target's computational population is preserved). The simulation uses `sesolve` (coherent dynamics, no decoherence) with the full 36-dimensional Hilbert space ($3 \times 3 \times 4$, two Cs controls and one Rb target).

1.3.1. Per-State Results

Input	Cubic SG	Hanning	ΔF
$\|0,0,A\rangle$	0.8715	**0.9961**	+0.125
$\|0,0,B\rangle$	0.8715	**0.9961**	+0.125
$\|0,1,A\rangle$	0.9998	**1.0000**	+0.000
$\|0,1,B\rangle$	0.9998	**1.0000**	+0.000
$\|1,0,A\rangle$	0.9998	**1.0000**	+0.000
$\|1,0,B\rangle$	0.9998	**1.0000**	+0.000
$\|1,1,A\rangle$	0.9962	**0.9983**	+0.002
$\|1,1,B\rangle$	0.9962	**0.9983**	+0.002
$\bar{\mathcal{F}}$	**0.9668**	**0.9986**	**+0.032**



Figure 2: Per-state fidelity for the two pulse shapes. The Hanning pulse (orange) achieves uniformly higher fidelity across all 8 basis inputs. The dominant improvement is in the unblocked $|0,0\rangle$ inputs.

1.3.2. Summary

Metric	**Cubic SG**	**Hanning**
Average gate fidelity $\bar{\mathcal{F}}$	0.9668	**0.9986**
Gate infidelity $1-\bar{\mathcal{F}}$	3.32×10^{-2}	$\mathbf{1.42\times10^{-3}}$
Probe duration	340 ns	533 ns
Total gate time	360 ns	553 ns
Worst-case state fidelity	0.871	**0.996**
$P(\|R\rangle)$ leakage ($\|0,0\rangle$ inputs)	6.9%	**0.0%**

The Hanning pulse reduces the gate infidelity by **23×** (from 3.3% to 0.14%) at the cost of a 1.6 × longer gate time (553 vs 360 ns). Both pulses now start and end at zero, ensuring a fair comparison. The improvement is almost entirely due to the unblocked $|0,0\rangle$ inputs, where the cubic super-Gaussian leaves 6.9% of the population stranded in the Rydberg state $|R\rangle$, while the Hanning pulse achieves complete Rabi return ($P(|R\rangle)\approx 0$).

The smooth, symmetric rise and fall of the Hanning window ensures that the effective two-photon Rabi cycle on the bright state $(|A\rangle+|B\rangle)/\sqrt{2}$ completes a full rotation and returns all population to the computational subspace. The cubic super-Gaussian, despite starting and ending at zero, has an asymmetric envelope due to the cubic exponent $(t-t_c)^3$. This asymmetry causes the accumulated bright-state Rabi angle to miss an exact multiple of 2π, leaving population in $|R\rangle$ at the end of the pulse.