



Platelets Are Not Just for Clots

James D. McFadyen, Zane S. Kaplan *

Australian Centre for Blood Diseases, Monash University, Melbourne, Australia



ARTICLE INFO

Available online 12 January 2015

Keywords:

Platelets
Inflammation
Thrombosis
Leukocytes
Cancer

ABSTRACT

Although the role of platelets as central mediators of hemostasis and thrombosis has been the primary focus of research into platelet biology for more than a century, over the last decade, nonhemostatic functions of platelets have been increasingly defined. As such, a large body of experimental evidence now exists, which places the platelet as a key player in mediating a diverse range of immune, inflammatory, and malignant disease processes. This review outlines the central mechanisms that underpin the nonhemostatic role of platelets and provides a summary of evidence demonstrating a role for platelets in mediating selected inflammatory, immune, and malignant disease processes.

Crown Copyright © 2015 Published by Elsevier Inc. All rights reserved.

Contents

The Classical Role of Platelets	111
Evolving Concepts in Platelet Biology—Linking Hemostasis, Inflammation, And Immunity	111
Platelets as Sentinels in Antimicrobial Surveillance and Response to Invading Pathogens	111
Platelet Immunoreceptors and Toll-Like Receptors.	111
Platelet CD40 Ligand, a Central Modulator of the Immune Response.	112
Platelet-Endothelial Adhesive Interactions in Inflammation.	113
Platelet-Derived Chemical Signals in Endothelial Activation	113
Platelet-Leukocyte Crosstalk in Inflammation	114
Platelets as Mediators of Inflammatory Disease	115
Atherosclerosis	115
Transfusion-Related Acute Lung Injury	115
Immune Thrombocytopenic Purpura (ITP)	115
Transfusion-Related Immunomodulation	116
Sepsis	116
Cancer	116
Rheumatoid Arthritis	117
Summary and Perspectives	117
Conflict of Interest Statement	117
Acknowledgments	117
References	117

The fundamental role of platelets in hemostasis was first identified by the Italian pathologist and pioneering medical researcher Bizzozzero in 1882 [144]. Subsequently, for more than 120 years, science has focused on the central role of platelets as the key cellular components mediating hemostasis and, latterly, as pivotal instigators of pathological thrombosis.

* Corresponding author at: Zane S. Kaplan, MBBS FRACP FRCPA, Clinical Haematologist, Australian Centre for Blood Diseases, Alfred Medical Research and Education Precinct (AMREP), 6th Level Burnet Tower, 89 Commercial Rd, Melbourne, Victoria 3004, Australia.
E-mail address: zane.kaplan@monash.edu (Z.S. Kaplan).

These classical roles of platelets in mediating thrombosis and hemostasis are now well accepted. However, over the past 2 decades, an ever-increasing literature has evolved linking platelets with biological responses beyond hemostasis and thrombosis. Emerging evidence in the field of platelet biology supports a major proinflammatory function for platelets, linked to host defence and a variety of autoimmune, inflammatory, and malignant diseases processes. The weight of this evidence suggests that platelets may be as important for host defenses and pathological inflammatory and malignant diseases as they are for hemostasis and thrombosis. The purpose of this review is to therefore

provide a brief overview of the classical role of platelets followed by a discussion of the interactions between platelets and the immune system, and the emerging role for platelets in disorders beyond hemostasis.

The Classical Role of Platelets

The hemostatic function of platelets is of fundamental importance in order to maintain the integrity of a closed, high-pressure circulatory system [1]. Ordinarily platelets circulate in a quiescent state, near the endothelial cells lining the blood vessels without forming stable adhesions. However, after breaches of the vasculature, a number of highly reactive subendothelial matrix proteins become exposed including von Willebrand factor (VWF), collagen, fibrinogen, and fibronectin. Initial platelet adhesion is mediated principally by platelet surface receptors interacting with their cognate ligands. Initial platelet adhesive interactions are mediated by platelet surface glycoproteins (GPs) including GPVI and GPIIb α interacting with endothelial bound collagen and VWF, respectively [2,3]. These receptor-ligand interactions initiate a cascade of intracellular responses resulting in amplification of platelet activation responses through the secretion of soluble agonists including thromboxane A₂ (TXA₂) and ADP, which act in concert with the engaged platelet receptors to mobilize intracellular Ca²⁺, which instigates platelet shape change, degranulation, and up-regulation of adhesive function of integrin $\alpha_{IIb}\beta_3$. Integrin $\alpha_{IIb}\beta_3$, in its active conformation, can then bind fibrinogen with high affinity allowing a hemostatic platelet aggregate or plug to form [4]. The fundamental role of VWF, GPIIb α , and $\alpha_{IIb}\beta_3$ in normal hemostatic responses is evidenced by the bleeding diathesis observed in patients with von Willebrand disease, Bernard-Soulier syndrome [5], and Glanzmann thrombasthenia [6], who have quantitative or qualitative abnormalities of VWF, GPIIb α , and $\alpha_{IIb}\beta_3$ respectively. The more detailed aspects of the hemostatic role of platelets have been extensively reviewed elsewhere [3,7,8].

Evolving Concepts in Platelet Biology—Linking Hemostasis, Inflammation, And Immunity

Platelets are the second most abundant cell type in the circulation after red blood cells, with an average of 1 trillion platelets present at any one time [3,9]. In addition to their vast number, the small size of platelets aids in their role as guardians of the vasculature and sentinels for infection. The average discoid platelet is approximately 2 μ m in diameter. Owing to this small size, under laminar flow conditions, platelets are “pushed” to the periphery by larger erythrocytes and leukocytes, forcing them into close contact with the vessel wall [9]. This allows for the rapid detection of any breaches in the endothelium in addition to sensing any invading pathogens.

Platelets contain 3 principal types of cytoplasmic granules, including α -granules, dense granules, and lysosomes. These granules possess a vast array of proteins that have autocrine and paracrine functions and include adhesive proteins, coagulation factors, chemokines, cytokines, growth factors, proteoglycans, immunoglobulins, and proteases and protease inhibitors. In addition to prestored mediators, recent evidence from proteomic studies suggests that platelets have an extensive transcriptome and proteome [10,11]. Although platelets lack a nucleus and hence genomic DNA, they do contain messenger RNA (mRNA) and all the required translational machinery required for protein synthesis. Recent evidence suggests that after platelet activation by potent agonists such as thrombin, the platelet releasate contains in excess of 300 proteins [11,12]. Consequently, when platelets become activated, they degranulate releasing a vast array of cytokines and chemokines that play a key role in modulating immune responses. Thus, conceptually, the immune response to tissue injury is guided, in part, by the thrombotic response by way of the numerous proteins stored within platelet dense and α -granules (see Table). Hence, it is becoming increasingly recognized that platelets may be as important for host immunity and inflammation as they are for hemostasis. Below we will detail the mechanisms by

which platelets may act as inflammatory and immune mediators followed by a summary of the current evidence demonstrating a central role for platelets in mediating the pathogenesis of a diverse range of clinical conditions.

Platelets as Sentinels in Antimicrobial Surveillance and Response to Invading Pathogens

Platelets play an essential role in pathogen surveillance, containment, and wound healing [13]. Platelets are the first and most abundant cell type to accumulate at sites of intravascular infection [14], where they likely play a central role in surveillance and recognition of microbial invasion [15]. In addition, platelets are able to orchestrate complex immune and inflammatory events through a combination of direct cell-cell contacts as well as through the release of a vast arsenal of bioactive molecules (see above and below) that facilitate “information transfer” with immune effector cells [15]. Platelets have been shown to have a role in aiding the immune defense against infections by bacteria, parasites, viruses, fungi, protozoa, and spirochetes [15]. Furthermore, platelet granules contain platelet microbicidal proteins and peptides including the classical chemokines (kinocidins and thrombo-cidins), platelet factor 4 (PF4), CTAP-III, regulated on activation normal T expressed and secreted (RANTES), fibrinopeptide B, and thymosin β -4, which act on a broad range of pathogens, including some strains of *Staphylococcus aureus* (*S. aureus*), viridans streptococci, and *Candida albicans* [16]. In addition, platelet microbicidal proteins have also been shown to potentiate the effect of antibiotic therapies [15].

Activated platelets have been shown by Youssefian et al [17] to engulf both HIV particles and *S. aureus* in vacuoles that became a prominent site of α -granule release. These studies suggest that activated platelets expressing P-selectin may assist the phagocytosis and limit the spread of microorganisms by guiding leukocytes to areas of tissue injury where invading microorganisms are trapped by platelets. However, it is conceivable that the HIV virus and the bacteria use platelet engulfment to evade the immune system, as neither killing nor digestion of microorganisms has been demonstrated within platelets. The interaction between platelets and *S. aureus* has been implicated in the pathogenesis of infective endocarditis [16]. In protozoal infection by *Toxoplasma gondii*, platelets have been observed to kill the parasites via a mechanism involving TXA₂ synthesis [18]. In malarial infections, platelets are able to kill *Plasmodium falciparum*-infected human red cells in vitro, and in a murine malaria model, platelet depletion or aspirin treatment reduced survival of *P. falciparum* parasites [19]. This, however, presents a biological paradox as platelets are implicated in the pathogenesis of cerebral malaria, a condition associated with a high mortality. Platelets may orchestrate this deleterious effect either via binding to and activating the endothelium, or through the formation of aggregates between parasitized red cells and platelets, which can obstruct blood vessels [20–23].

Platelet Immunoreceptors and Toll-Like Receptors

Many of the immunologic functions of platelets involve interactions with platelet immunoreceptors including members of the Fc receptor (FcR) and toll-like receptor (TLR) families.

Platelets express FcRs, which are able to recognize IgG, IgE, and IgA immunoglobulin subclasses on their surface. The FcRs expressed by human platelets include Fc ϵ RII, Fc γ RIIA, and Fc α RI [24]. Fc γ RIIA has been implicated in the development of heparin-induced thrombocytopenia [25] as well as the pathogenesis of bacterial endocarditis where platelet activation occurs via parallel binding between Fc γ RIIA and IgG antibodies directed against the bacteria and fibrinogen binding to $\alpha_{IIb}\beta_3$ [16]. The platelet collagen receptor GPVI is also a member of the Ig superfamily. GPVI forms a complex with the immunoreceptor tyrosine activation motif-containing Fc γ R, which is critical for GPVI signaling [24,26]. An additional immunoreceptor tyrosine activation motif

Table

Key thromboinflammatory modulators released by platelets [15,64,69,85,139–143]

Localization within the platelet	Released molecule	Target cell	Function
Dense granule	• Histamine	Endothelial cells, monocytes, neutrophils, NK cells, T and B cells and eosinophils	Increased endothelial permeability, expression of adhesive ligands on the endothelial surface and endothelial cytokine and chemokine release
	• 5HT	Monocytes and T cells	Activation of leukocytes—proinflammatory
α -Granules			Monocyte and T-cell activation—proinflammatory
• Adhesion receptors	• $\alpha_{IIb}\beta_{III}$	Platelet	Platelet aggregation, platelet-endothelial, and platelet-leukocyte adhesion
	• GPIb α	Platelet	Platelet adhesion under shear. Platelet-leukocyte interaction via GPIb α -Mac-1 binding
• Adhesive ligands	• P-selectin	Platelets, endothelial cell, and leukocytes—PSGL-1	Adhesion of leukocytes to platelets and endothelial cells
• Chemokines	• CXCL1–GRO- α	Neutrophils and monocytes	Induces neutrophil chemotaxis and monocyte adhesion under flow
	• CXCL4-PF4	Neutrophils and monocytes	Synergizes with other cytokines, leukocyte adhesion, and monocyte differentiation
	• CXCL5–ENA-78	Neutrophils	Induces neutrophil chemotaxis
	• CXCL7–NAP-2	Neutrophils	Enhances neutrophil and monocyte adhesion and neutrophil transendothelial migration
	• CXCL8–IL-8	Neutrophils, T cells, and basophils	Induces chemotaxis and neutrophil activation
	• CCL2–MCP-1	Monocytes and T cells	Monocyte chemotaxis and infiltration and T-cell production of IL-4
	• CCL3–MIP-1 α	Monocytes, NK cells, T and B cells, eosinophils, and basophils	Induces chemotaxis and activation
	• CCL5–RANTES	Endothelium/monocytes	Promotes monocytes adhesion to the endothelium and chemokine synthesis
• Growth factors	• PDGF	Vascular smooth muscle cells, monocytes, macrophages, T cells, and B cells	Promotes wound. Stimulates vascular smooth muscle cell migration and proliferation associated with intimal hyperplasia
	• TGF- β	Endothelial cells, monocytes, lymphocytes, NK cells, and neutrophils	Affects multiple regulatory pathways to influence chemotaxis, activation, chemokine release, and cell survival
and• Synthesized cytokines	• IL-1 β	Endothelial cells	Up-regulation of leukocyte adhesion molecules and stimulation of endothelial release of proinflammatory cytokines
α -Granule and platelet membrane			
	• CD40 ligand	Endothelial cell CD40	Up-regulation of leukocyte adhesion molecules and stimulation of endothelial release of proinflammatory cytokines. B cell-class switching
• Proinflammatory lipids	• TXA ₂	Platelets, macrophages, T cells	Platelet activation, promotes inflammation
	• Leukotriene B ₄	Leukocytes—BLT-1 and BLT-2	Chemoattractant and promotion of monocyte differentiation into foam cells and neutrophil activation
	• PAF	Platelet/Leukocytes	Platelet agonist and promotes neutrophil adhesion

5HT, 5 hydroxy-tyramine; TGF- β , transforming growth factor β .

expressed on the surface of platelets is C-type lectin-like receptor 2, which induces potent platelet activation signals via Syk kinase-dependent signaling pathways [27,28]. Activation of platelets through C-type lectin-like receptor 2 binding to its cognate ligand podoplanin facilitates blood/lymphatic vessel separation [29]. This interaction has also been implicated in tumor metastasis [27].

Toll-like receptors are a family of evolutionarily conserved germ-line encoded pattern recognition receptors that recognize conserved molecular motifs expressed by pathogens referred to as pathogen-associated molecular patterns (PAMPs) [30]. Toll-like receptors are divided into subfamilies according to the type of PAMPs they recognize, which include lipoproteins, lipids, proteins, and nucleic acids derived from a wide range of microbes such as bacteria, viruses, parasites, and fungi [31]. Toll-like receptors are not only expressed by professional phagocytic cells (neutrophils, dendritic cells, and macrophages) but are also expressed by platelets [9]. Studies have demonstrated that both murine and human platelets express TLRs1–9, many of which are functional [9] and have been implicated in protective responses to invading pathogens as well as in pathologic immune responses [32–34]. A well-characterized and studied TLR family member is TLR4, which recognizes bacterial lipopolysaccharide (LPS), a component of the outer membrane of gram-negative bacteria, which is a potent immunostimulant that can cause septic shock [31]. Lipopolysaccharide-induced TLR4 signaling in murine models induces platelet adhesion to fibrinogen and platelet sequestration in the microvasculature of the lung, liver, and spleen [35] as well as thrombocytopenia and release of the potent proinflammatory cytokine tissue necrosis factor α (TNF- α) [33]. In addition, recognition of LPS by platelet TLR4 has been shown to induce platelet-neutrophil binding and subsequent robust neutrophil activation resulting in the

formation of neutrophil extracellular traps (NETs) [36]. As discussed below, although NETs may represent a potential bacterial trapping mechanism in severe sepsis, exaggerated NETs formation is deleterious to the host. Expression of functional TLR2 has also been demonstrated in platelets [37]. Platelet TLR2 stimulation by bacteria or viral pathogens via the phosphoinositide 3-kinase signaling pathway is able to directly activate both the prothrombotic and proinflammatory platelet responses [37–39] and thus may exacerbate the thrombotic complications of sepsis. Recently, Podrez and colleagues [40] demonstrated that platelet TLR9 activation is an important factor linking platelet activation with thrombosis, oxidative stress, and innate immunity. Interestingly, platelet TLR9 is uniquely distributed to the electron-dense tubular system-related compartment of the platelet, which has been labeled the “T granule” [41]. In T granules, TLR9 is associated with either VAMP 7 or VAMP 8, its association with the latter is proposed to increase TLR9 surface expression via VAMP 8-mediated T-granule fusion with the open canalicular system [41]. In the model proposed by Thon and colleagues [41], platelets are primed to express TLR9 on their surface during activation, which subsequently facilitates secondary activation through PAMPs expressed on invading pathogens. Therefore, platelet-derived TLRs may be as important in orchestrating platelet thrombotic and inflammatory as they are in alerting the innate immune system to pathogen invasion.

Platelet CD40 Ligand, a Central Modulator of the Immune Response

CD40 ligand (CD40L) is a transmembrane protein with structural homology to TNF- α [42] and is an important bioactive modulator of adaptive immunity. Although CD40L was first identified on the surface

of activated CD4⁺ T cells, it has subsequently been demonstrated that platelets are a rich source of CD40L [43], with greater than 95% of circulating CD40L being of platelet origin [44]. CD40 ligand interacts with its cognate receptor CD40 expressed on numerous cell types including smooth muscle cells, endothelial cells, fibroblasts, dendritic cells, leukocytes, and platelets [42,45]. Through CD40L-dependent mechanisms, platelets are able to activate dendritic cells, resulting in release of proinflammatory cytokines and chemokines as well as activation of naïve T cells, which are fundamental to the establishment of cellular immune responses [9,46]. Within the humoral immune system, CD40L plays a critical role by supporting T-cell-dependent B-cell differentiation and immunoglobulin isotype switching [47]. The importance of this role of CD40L in humoral immunity is evidenced by the severe immunodeficiency state characterized by recurrent bacterial infections observed in patients with X-linked hyper-IgM syndrome who possess a defect in the gene encoding CD40L [48]. Platelet CD40L can also interact with neutrophils, and this interaction may be particularly relevant in the genesis of transfusion-related acute lung injury (TRALI). In this context, platelet-derived soluble CD40L may accumulate during platelet storage, which has the capacity to activate primed neutrophils in the lungs of predisposed patients, resulting in endothelial damage and possibly TRALI, as discussed below [49]. In addition to the aforementioned interactions with hematopoietic cells, CD40L can also modulate inflammatory responses through its interaction with endothelial cells, whereby CD40L binding to its cognate receptor CD40 on endothelial cells results in up-regulation of intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), E-selectin, and P-selectin as well as the release of interleukin (IL)-6 and tissue factor [9]. In addition,

CD40L inhibits endothelial nitric oxide (NO) production by destabilizing NO synthase mRNA [42,50]; thus, CD40L binding to endothelial cells results in a proinflammatory phenotype [42]. Therefore, through its interactions with multiple cell types, CD40L is at the forefront of modulating immune and inflammatory responses.

Platelet-Endothelial Adhesive Interactions in Inflammation

It is increasingly recognized that inflammatory stimuli facilitate sustained endothelial-platelet adhesion by perturbing the antiadhesive properties of quiescent endothelium and inducing increased surface expression of endothelial adhesion molecules [51]. The adhesion of platelets to “inflamed endothelium” involves a similar coordinated multistep process as occurs in hemostasis and thrombosis, including platelet tethering, surface translocation, and firm adhesion (Fig 1) [52]. The expression of the endothelial selectins, P- and E-selectin, the membrane expression of VWF and ICAM-1 bound fibrinogen are all important in mediating platelet adhesion to the inflamed endothelium. The counter-receptors for P-selectin include GPIb α on platelets and P-selectin glycoprotein-1 (PSGL-1) on leukocytes [53,54], which support tethering and rolling, whereas it is thought that stable platelet adhesion is contingent upon platelet $\alpha_{IIb}\beta_3$ binding to fibrinogen.

Platelet-Derived Chemical Signals in Endothelial Activation

Platelets activated through their interactions with inflamed endothelium release a vast array of inflammatory and mitogenic mediators (see Table) into the local microenvironment. These platelet-derived




Fig 1. Platelet endothelial interactions in health and in inflammation. A, The healthy endothelium: the antiadhesive phenotype of endothelial cells is maintained through 3 intrinsic pathways; ecto-ADPase/CD39/NTDPase, which metabolizes ADP, and the PGI₂ and NO pathways, which inhibit platelet activation through stimulation of cAMP and cGMP production, respectively. B, Inflamed endothelium–endothelial activation: in infective or highly oxidative states as well as in the presence of a hyperlipidaemic milieu the endothelium becomes inflamed as a result of modified lipoprotein particles, cytokines, and reactive oxygen species accumulating in the intima. This leads to the expression of adhesive ligands (VWF and P- and E-selectins) on the endothelium. C, Platelet adhesion to the inflamed endothelium and platelet endothelial crosstalk: expression of VWF and P-selectin on the endothelial surface supports platelet tethering and rolling. Subsequently, stable platelet adhesion occurs through fibrinogen bridging of $\alpha_v\beta_3$ or ICAM-1 on endothelial cells with $\alpha_{IIb}\beta_3$ on the platelet surface. Adherent platelets secrete numerous bioactive substances which alter the chemotactic and adhesive properties of the endothelial cells. Platelet-derived IL-1 β induces endothelial secretion of IL-6 and IL-8 as well as surface expression of ICAM-1, $\alpha_v\beta_3$, and MCP-1. Platelet CD40L binds CD40 on endothelial cells resulting in up-regulation of adhesive molecules (ICAM-1, VCAM-1, E-selectin, and P-selectin), cytokine, and tissue factor release and reduction in NO synthesis. Figure adapted from Kaplan and Jackson [85] with permission.

bioactive substances alter the chemotactic and adhesive properties of the endothelial cells, enabling monocytes and other leukocytes to adhere and transmigrate through the endothelium to sites of inflammation [55,56]. In vivo models of atherogenesis (see below), a prototypic disease of thromboinflammatory dysregulation, suggest that these interactions between endothelial cells and activated platelets are likely to be critical in the initiation of atherosclerosis [57,58].

Platelet-Leukocyte Crosstalk in Inflammation

There is increasing evidence that activated platelets adherent to the inflamed endothelium may enhance leukocyte recruitment, activation, and transmigration, thereby enhancing the inflammatory processes

(Fig 2), which are central to the genesis of clinically relevant thrombo-inflammatory diseases such as disseminated intravascular coagulation, transplant rejection, and ischemia-reperfusion injury. Leukocyte recruitment to endothelial-bound platelets is a coordinated multistep process involving selectin-mediated tethering and rolling, followed by β_2 integrin-mediated stable adhesion. Platelet P-selectin has a central role in mediating the initial capture and rolling of leukocytes through interactions with its key receptor, PSGL-1 [53,59]. Ligation of PSGL-1 induces signaling events culminating in the activation of the leukocyte β_2 integrins including macrophage antigen-1 (Mac-1; $\alpha_M\beta_2$) and lymphocyte function-associated antigen-1 (LFA-1; $\alpha_L\beta_2$), which are required to mediate stable leukocyte adhesion. Macrophage antigen-1 is the principal β_2 integrin that facilitates firm adhesion onto the surface of




Fig 2. Platelet-leukocyte interactions in inflammation. A, Platelet-mediated leukocyte recruitment and adhesion: leukocyte recruitment to endothelial-bound platelets occurs in a multi-step coordinated process. Initial tethering of leukocytes is mediated by the interaction of P-selectin expressed on the platelet surface with its cognate receptor PSGL-1. Ligation of PSGL-1 promotes activation of leukocyte β_2 integrins (MAC-1 and LFA-1), necessary for stable leukocyte adhesion. Note that Mac-1 can engage several platelet ligands including GPIb, ICAM-2, and JAM-3, as well as fibrinogen bound to $\alpha_{IIb}\beta_3$. Leukocytes subsequently undergo transendothelial migration to sites of infection and inflammation. B, Platelet leukocyte crosstalk: bioactive mediators derived from α -granules of activated platelets, including PAF, synergizes with RANTES to enhance leukocyte adhesion and monocyte differentiation. RANTES acting in concert with P-selectin results in MCP-1 and IL-8 secretion by monocytes. CTAP-III is another chemokine released from the platelet α -granules, which is subsequently converted into active NAP-2 by the neutrophil membrane associated serine-protease cathepsin G. NAP-2 enhances leukocyte adhesion and neutrophil transendothelial migration and, within the developing thrombus, provides a chemokine gradient guiding leukocytes toward the site of vascular injury. Up-regulation of COX-2 in monocytes via pathways involving IL-1 β and P-selectin results in increased production of proinflammatory mediators including PAF, leukotrienes, and TXA₂ via transcellular eicosanoid metabolism. Figure adapted from Kaplan and Jackson [85] with permission.

platelets via its binding to multiple ligands including GPIb α [60], ICAM-2 [61], and junctional adhesion molecule-3 (JAM-3) [62] as well as through fibrinogen bound to $\alpha_{IIb}\beta_3$ [56,63].

Platelet activation with consequent degranulation results in liberation of numerous chemokines stored in platelet α -granules (see Table). These chemokines can either be released into the circulation or displayed on the platelet surface where they exert numerous biological activities instrumental in mediating inflammatory and immune responses (Fig 2) [64]. Classically, chemokines are divided into 4 families (CXC, CC, CX3C, and XC) based on the arrangement of the N-terminal cysteine residues [65]. The CC and CXC bind exclusively to CC and CXC chemokine receptors, respectively; however, within families there is considerable promiscuity relating to binding of chemokines to chemokine receptors [64]. Platelet granules contain both CXC (PF4, growth-regulating oncogene- α [GRO- α], epithelial neutrophil-activating protein-78 [ENA-78], and CXCL7) and CC chemokines (RANTES, macrophage inflammatory protein-1 α [MIP-1 α], and monocyte chemoattractant protein [MCP]-3), with platelets being the most abundant and readily available source of CXC chemokines [66]. In addition to releasing prestored chemokines and cytokines, platelets also participate in transcellular metabolism of eicosanoids with endothelial cells and leukocytes, leading to the local synthesis of proinflammatory lipids including platelet-activating factor (PAF), leukotrienes, and TXA₂ [67].

In recent studies, we have identified a major role for the platelet chemokine neutrophil-activating peptide 2 (NAP-2; CXCL7) in regulating leukocyte polarization and motility [68]. CXCL7 is the most abundant platelet chemokine that appears in serum at micromolar concentration; this is 2-fold greater than PF4, the next most abundant, and several orders of magnitude greater than most other platelet chemokines [69]. CXCL7 has several molecular variants that are derived by sequential N-terminal proteolytic events [70]. Interestingly, NAP-2 is only generated in the presence of leukocytes through proteolytic cleavage by the neutrophil membrane-associated serine-protease cathepsin G [71] and mast cell chymase [72]. Furthermore, NAP-2 represents the only CXCL-7 derivative that exhibits typical chemokine properties [69]. Although NAP-2 has been demonstrated to induce neutrophil and monocyte adhesion in vitro as well as neutrophil transendothelial migration [64,73] and elastase release [74], our studies provide the first in vivo evidence that graded platelet activation within a developing thrombus results in the generation of a NAP-2 chemotactic gradient within the thrombus body, guiding leukocytes to sites of vascular injury.

Additional CXC chemokines expressed in platelets include CXCL4 (PF4), CXCL1 (GRO- α), CXCL5 (ENA-78), and CXCL8 (IL-8). PF4 is the second most abundant platelet chemokine; however, it does not exert classical chemokine functions at submicromolar concentrations [64,75] on its own, but rather acts together with other platelet-derived chemokines such as RANTES to modulate leukocyte responses. CXCL1 is not present exclusively in platelets, but it is also expressed by neutrophils, monocytes, and endothelial cells [64,76]. CXCL1 is able to promote both neutrophil recruitment [77] and arrest of monocytes under flow conditions [78]. CXCL5 has significant structural homology with NAP-2 [79] and shares its ability to induce neutrophil chemotaxis [80] with a similar potency to IL-8 [81]. In addition to being a chemoattractant for neutrophils, IL-8 is also able to stimulate neutrophil degranulation and enhance expression of Mac-1 [82]. Interleukin-8 is also a chemoattractant for T cells and basophils and also induces histamine release [83]. It, however, is not produced exclusively by platelets [84].

Platelets as Mediators of Inflammatory Disease

Atherosclerosis

The concept that platelets may have proinflammatory functions emerged from extensive experimental work highlighting their role in the pathogenesis of atherosclerosis. Consequently, the role of platelets in atherogenesis has been the focus of recent comprehensive reviews

[52,55,85,86] and therefore will only be discussed briefly. The adhesion of platelets to sites of injured and atherosclerosis prone endothelium may conspire to accelerate the progression of atherosclerotic lesions. Adherent platelets release a number of chemokines (CCL5), cytokines (IL-1 β), and other biologic response modulators (CD40L) that act to enhance the recruitment of proatherogenic monocytes. In addition, adherent platelets may release growth factors, such as platelet-derived growth factor (PDGF), which may enhance angiogenesis and smooth muscle proliferation in atherosclerotic plaques.

In addition to enhancing atherogenesis, platelets may also contribute to atherothrombosis by augmenting plaque instability via CD40L binding to smooth muscle cells, endothelial cells, and monocytes, which induce expression of matrix metalloproteinases [50]. Matrix metalloproteinase possesses collagenase activity and thus may cause degradation of the extracellular matrix collagen culminating in the thinning of the fibrous cap and consequent plaque instability. Consistent with this hypothesis, enhanced collagen content has been observed in the extracellular matrix of atherosclerotic lesion in murine models in which CD40L signaling has been interrupted [87,88]. Furthermore, though controversial, CD40L may be important in forming the necrotic core of the atheromatous plaque through induction of proapoptotic pathways [50]. Therefore, CD40L contingent mechanisms may enhance plaque, instability making them more susceptible to rupture.

Transfusion-Related Acute Lung Injury

Transfusion-related acute lung injury is defined as the new onset of acute lung injury within 6 hours of transfusion. Clinically, TRALI may manifest as hypoxia, fever, chills, tachycardia, and bilateral noncardiogenic pulmonary edema and is the leading cause of transfusion-associated mortality [89,90]. The central role of neutrophils in mediating TRALI has long been appreciated. Indeed, this has led to a “2-hit” model of neutrophil-induced acute lung injury, such that the first hit represents the susceptible host with underlying pulmonary pathology, with the second hit being a transfusion-related insult [89,90]. The “first hit” results in endothelial activation which facilitates neutrophil adhesion via the up-regulation of endothelial adhesion molecules and chemokine synthesis. The “second hit” then results in the activation of adherent neutrophils which leads to degranulation and release of mediators that cause endothelial damage and increased vascular permeability. In the context of TRALI, the second hit that causes neutrophil activation can include antineutrophil antibodies or bioactive lipids such as lysophosphatidylcholine. The source of these mediators that may induce the second hit is thought to either be donor derived (antineutrophil antibodies) or result from the breakdown of biological membranes, which may occur during the storage process of blood products. More recently, experimental data have demonstrated that soluble CD40L (also known as CD154) may be released by stored platelet concentrates and promote neutrophil activation via activation of neutrophil CD40 receptors [49]. Platelets have also been demonstrated to be directly involved in the pathogenesis of TRALI in murine models by way of their ability to bind to neutrophils and instigate NET formation [91]. The importance of platelets in this process was highlighted by the ability of platelet inhibition to protect from the development of TRALI [91]; however, the precise mechanisms leading to NET formation in TRALI are yet to be established.

Immune Thrombocytopenic Purpura (ITP)

Immune thrombocytopenic purpura (ITP) is perhaps the classical immune condition that one associates with platelets. It is associated with immune dysregulation and involves antigen-presenting cells, B cells, and T cells acting in concert to perpetuate an autoimmune response that hastens the destruction of platelets and/or perturbs megakaryopoiesis [92,93]. It has long been observed that ITP is associated with infections, and more recent data suggest that platelet TLR4

expression may play a central role in this process [32,94]. As discussed above, platelets express all of the TLRs and TLR4 is thought to be responsible for mediating thrombocytopenia and TNF- α production in response to LPS. In the context of ITP, recent evidence has demonstrated that gram-negative LPS can bind to platelet TLR4 and synergize with antiplatelet antibodies, thus enhancing phagocytosis in vitro and thrombocytopenia in vivo [32]. This may explain why some patients with ITP experience lower platelet counts with intercurrent infection, which may abate with antibacterial treatment.

Transfusion-Related Immunomodulation

The clinical observation that blood transfusion may be associated with an increase in infection risk or tumor growth or improved outcomes after renal allograft transplantation has given rise to the concept of transfusion-related immunomodulation [95]. Although there may be multiple ways the immune system could be modulated by blood transfusion, there are now data supporting a major role for platelets in this process. Although platelets are devoid of MHC class II molecules, they express a significant amount of MHC class I [96]. Surprisingly, there appears to be 2 distinct sources of platelet MHC class I molecules. Platelets contain both surface-derived, denatured class I MHC molecules that are adsorbed by platelets from the plasma. Furthermore, it has recently been demonstrated that platelets contain all the necessary machinery (such as MHC mRNA and proteasomes) for class I MHC expression and antigen presentation [97]. Interestingly, functional consequences of MHC class I molecule expression appear somewhat divergent. In the context of transfusion medicine, denatured MHC class I molecules can be shed from the surface of stored platelets; however, these shed allogeneic MHC class I molecules do not appear able to generate a cytotoxic T cell response and have been demonstrated to improve the survival of allogeneic skin grafts in mouse models [98]. On the other hand, the platelets' ability to process antigenic material and express it in a MHC class I dependent manner has recently been shown in a mouse model of cerebral malaria to play a central role in both the acquired immune response and in enhancing cytotoxic T-cell activation [97].

Sepsis

Thrombocytopenia has long been considered a manifestation of sepsis and is associated with increased mortality [9]. Recent experimental evidence now suggests that platelets play an active role in the pathogenesis of sepsis and multiorgan dysfunction. Elevated circulating levels of platelet-leukocyte aggregates have previously been reported in a broad range of inflammatory conditions including sepsis [15,67,99]. Platelet-neutrophil aggregates comprise activated platelets bound to leukocytes. These aggregates are supported by interactions between platelet P-selectin and GPIIb α and their respective leukocyte counter receptors, PSGL-1 and Mac-1 [100]. In addition, plasma from septic but not healthy patients can induce platelet-neutrophil interactions in vitro [36]. These interactions are likely highly contingent upon stimulation of the platelet TLR4 receptor given the observation that the use of the TLR4 antagonist eritoran resulted in a 50% reduction in platelet-neutrophil interactions [36].

The functional consequences of platelet-leukocyte interactions are elucidated by the recent observation that LPS-stimulated platelets bind to neutrophils and stimulate the formation of NETs. Neutrophil extracellular trap formation represents a mechanism used by neutrophils to entrap and possibly kill pathogens [101]. This neutrophil killing strategy involves the active extrusion of nuclear and cytoplasmic granular constituents that combine to form NETs [101]. Neutrophil extracellular traps are composed of chromatin (DNA and histones) fibers that are coated with proteins from azurophilic granules (neutrophil elastase, cathepsin G, and myeloperoxidase) as well as proteins from specific and tertiary granules, such as lactoferrin and gelatinase [101]. Neutrophil extracellular traps are highly efficient at trapping and killing gram-

negative and gram-positive bacteria as well as mycobacteria, yeast, fungi, and protozoan [102]. However, exaggerated NETs release can have deleterious consequences, resulting in inflammatory and autoimmune diseases [102], as well as induction of pathological thrombosis [103–105]. Induction of NET release from leukocytes by LPS-stimulated platelets is strikingly efficient and rapid, occurring within minutes of exposure [36]. This contrasts with other NETs inducing stimuli such as sepsis, cytokines, and reactive oxygen species [102,106,107], which promote NET formation by neutrophils typically after 2 to 4 hours [108]. Platelets promote NET release through the LPS receptor TLR4, suggesting that platelet TLR4 may represent “a threshold switch” for bacterial trapping by NETs in severe sepsis [36].

In addition to inducing NET formation, activated platelets from septic patients have enhanced microparticle formation [109,110]. Microparticles are small (0.1–1 μ m) fragments of membrane shed from different cell types including platelets, leukocytes, and endothelial cells [111]. Platelet microparticles constitute approximately 70% to 90% of circulating microparticles [112]. Microparticles shed from activated platelets can express adhesion molecules such as P-selectin and $\alpha_{IIb}\beta_3$ and therefore can adhere to and activate endothelial cells and leukocytes [110]. Importantly, microparticles may contain potent proinflammatory molecules such as IL-1 β and CCL5, which, when shed from activated platelets, may communicate proinflammatory signals to extravascular tissue [112,113]. Platelet-derived microparticles also contain small non-protein-coding (21–23 nucleotides) microRNAs which are able to modulate cellular function and protein expression via degradation of mRNA or repression of mRNA translation [114,115]. Platelet microparticles are able to transfer microRNAs to cells types including endothelial cells and leukocytes [116] and thus through this pathway may modulate cellular functions involved in immunity and inflammation as well as in hematopoiesis, developmental biology, angiogenesis, and malignancy [114,115].

Cancer

A relationship between thrombocytosis and malignant tumors was first observed more than 100 years ago by Reiss et al in 1872 [117]. More recently, clinical data have demonstrated that an elevated platelet count is associated with more advanced cancer in addition to being an adverse prognostic marker [118–120]. There is now a growing body of evidence demonstrating a key role for platelets in the pathogenesis of tumor progression and metastasis. Tumor cells can aggregate platelets in vitro (or tumor cell-induced platelet aggregation) [121,122]. This property of tumor cells correlates with their ability to metastasize in vivo. Furthermore, platelet depletion or inhibition of platelet function has been shown to inhibit tumor metastases in murine models [121,123]. It is proposed that the ability of tumor cells to aggregate platelets gives rise to several survival advantages in vivo. Platelet adhesion to tumor cells may form a “shield” around tumor cells entering the circulation and therefore confer protection from immune clearance [124,125]. In addition, interaction of metastasizing tumor cells in the circulation with endothelial cells is enhanced by the presence of bound platelets [126]. Furthermore, platelets may release growth factors contained in their granules such as PDGF, vascular endothelial growth factor, and angiopoietin 1, which can facilitate tumor growth [127,128]. Based on experimental models where deficiency or inhibition of the platelet $\alpha_{IIb}\beta_3$, GPVI, or P selectin resulted in a decreased frequency of tumor metastasis, these are thought to be the key platelet receptors facilitating platelet-tumor interactions [127,128]. In addition to playing a role in platelet-tumor interactions, $\alpha_{IIb}\beta_3$ activation is also believed to be essential for the tumor growth enhancing of platelets, as activation of this integrin appears necessary for the release of proangiogenic factors from the platelet which support tumorigenesis [129].

Rheumatoid Arthritis

Electron microscopic analysis of the synovium from patients with rheumatoid arthritis (RA) more than 30 years ago demonstrated platelet thrombi in synovial vessels [130]. Furthermore, these studies identified platelets in the vicinity of gap junctions of synovial vessels. More recently, considerable evidence has accumulated highlighting enhanced platelet activation in patients with RA. In samples from patients with RA, it has been shown that platelets express higher levels of P-selectin, display enhanced responsiveness to soluble agonists *in vitro*, and demonstrate elevated levels of circulating platelet microparticles and platelet-leukocyte aggregates [131–134]. Furthermore, platelet aggregates and platelet-adherent leukocytes have been observed in the synovial fluid of patients with RA [135,136]. These observations suggest that platelets may not merely be passive bystanders in RA pathogenesis but rather may be key players in its pathogenesis.

Fundamental new insights into the pathogenic role of platelets in RA were provided in a pivotal study by Boilard et al [137]. Surprisingly, examination of synovial fluid from patients with RA demonstrated that the synovial fluid in RA is enriched in platelet microparticles rather than whole platelets [137]. Importantly, the concentrations of platelet-derived microparticles are much higher in the synovial fluid than in the blood, suggesting that they form locally in the joint vasculature. The key stimulus for platelet activation and platelet microparticle formation in RA appears to be the interaction of platelet GPVI with collagen and laminin [137]. Activation of platelet GPVI induces the release of platelet-derived microparticles containing IL-1 which can induce fibroblast-like synoviocytes to release the proinflammatory cytokines IL-6 and IL-8 [137]. The central role for the GPVI-induced platelet activation axis in promoting inflammatory arthritis is evidenced by murine models where deletion of platelet GPVI or platelet depletion resulted in a marked reduction in the level of arthritis activity [137]. Although the precise mechanism by which microparticles enter the synovial space is yet to be fully elucidated, it is conceivable that neutrophils with adherent platelet-derived microparticles may act to shuttle microparticles into the synovial fluid. Alternatively, owing to their small size, microparticles may enter the synovial space by diffusing across dilated endothelial gap junctions—a process that itself is exaggerated by serotonin released by platelets in arthritic joints [138].

Summary and Perspectives

For more than a century, platelet research has focused primarily on defining the hemostatic and thrombotic roles of platelets. However, studies into the nonhemostatic properties of platelets are in relative terms, in their infancy. As a corollary, just as expanding scientific knowledge on the prothrombotic role of platelets has enabled the development of a broad range of antiplatelet drugs that are widely used in cardiovascular disease, it is envisaged that enhancing our understanding of the proinflammatory and immune modulating roles of platelets will facilitate the development of interventions to therapeutically target these nonhemostatic roles of platelets.

Conflict of Interest Statement

The authors declare no relevant conflict of interest.

Acknowledgments

We would like to thank Dr Andre Samson and Associate Professor Robert Andrews for helpful discussions and constructive advice. J.D.M. is a CSL Prof Heimburger Award recipient, and Z.S.K. is an NHMRC and National Heart Foundation postgraduate scholarship holder and CSL Prof Heimburger Award recipient.

References

- [1] Furie B, Furie BC. Mechanisms of thrombus formation. *N Engl J Med* 2008;359(9):938–49.
- [2] Jackson SP. The growing complexity of platelet aggregation. *Blood* 2007;109(12):5087–95.
- [3] Ruggeri ZM. Platelet adhesion under flow. *Microcirculation* 2009;16(1):58–83.
- [4] McFadyen JD, Jackson SP. Differentiating haemostasis from thrombosis for therapeutic benefit. *Thromb Haemost* 2013;110(5):859–67.
- [5] Lopez JA, Andrews RK, Afshar-Kharghan V, Berndt MC. Bernard-Soulier syndrome. *Blood* 1998;91(12):4397–418.
- [6] George JN, Caen JP, Nurden AT. Glanzmann's thrombasthenia: the spectrum of clinical disease. *Blood* 1990;75(7):1383–95.
- [7] Ruggeri ZM, Mendolicchio GL. Adhesion mechanisms in platelet function. *Circ Res* 2007;100(12):1673–85.
- [8] Andrews RK, Lopez JA, Berndt MC. Molecular mechanisms of platelet adhesion and activation. *Int J Biochem Cell Biol* 1997;29(1):91–105.
- [9] Semple JW, Italiano Jr JE, Freedman J. Platelets and the immune continuum. *Nat Rev Immunol* 2011;11(4):264–74.
- [10] Macaulay IC, Carr P, Gusnanto A, Ouweland WH, Fitzgerald D, Watkins NA. Platelet genomics and proteomics in human health and disease. *J Clin Invest* 2005;115(12):3370–7.
- [11] Coppinger JA, Cagney G, Toomey S, Kislinger T, Belton O, McRedmond JP, et al. Characterization of the proteins released from activated platelets leads to localization of novel platelet proteins in human atherosclerotic lesions. *Blood* 2004;103(6):2096–104.
- [12] Wijten P, van Holten T, Woo LL, Bleijerveld OB, Roest M, Heck AJ, et al. High precision platelet releasate definition by quantitative reversed protein profiling—brief report. *Arterioscler Thromb Vasc Biol* 2013;33(7):1635–8.
- [13] Vieira-de-Abreu A, Rondina MT, Weyrich AS, Zimmerman GA. Inflammation. In: Michelson AD, editor. *Platelets*. Amsterdam: Academic Press; 2013. p. 733–65.
- [14] Yeaman MR. Platelets in defense against bacterial pathogens. *Cell Mol Life Sci* 2010;67(4):525–44.
- [15] Vieira-de-Abreu A, Campbell RA, Weyrich AS, Zimmerman GA. Platelets: versatile effector cells in hemostasis, inflammation, and the immune continuum. *Semin Immunopathol* 2012;34(1):5–30.
- [16] Fitzgerald JR, Foster TJ, Cox D. The interaction of bacterial pathogens with platelets. *Nat Rev Microbiol* 2006;4(6):445–57.
- [17] Youssefian T, Drouin A, Masse JM, Guichard J, Cramer EM. Host defense role of platelets: engulfment of HIV and *Staphylococcus aureus* occurs in a specific subcellular compartment and is enhanced by platelet activation. *Blood* 2002;99(11):4021–9.
- [18] Yong EC, Chi EY, Fritsche TR, Henderson Jr WR. Human platelet-mediated cytotoxicity against *Toxoplasma gondii*: role of thromboxane. *J Exp Med* 1991;173(1):65–78.
- [19] McMorran BJ, Marshall VM, de Graaf C, Drysdale KE, Shabbar M, Smyth GK, et al. Platelets kill intraerythrocytic malarial parasites and mediate survival to infection. *Science* 2009;323(5915):797–800.
- [20] van der Heyde HC, Gramaglia I, Sun G, Woods C. Platelet depletion by anti-CD41 (alphaIIb) mAb injection early but not late in the course of disease protects against *Plasmodium berghei* pathogenesis by altering the levels of pathogenic cytokines. *Blood* 2005;105(5):1956–63.
- [21] Sun G, Chang WL, Li J, Berney SM, Kimpel D, van der Heyde HC. Inhibition of platelet adherence to brain microvasculature protects against severe *Plasmodium berghei* malaria. *Infect Immun* 2003;71(11):6553–61.
- [22] Faillle D, El-Assaad F, Alessi MC, Fusai T, Combes V, Grau GE. Platelet-endothelial cell interactions in cerebral malaria: the end of a cordial understanding. *Thromb Haemost* 2009;102(6):1093–102.
- [23] Cox D, McConkey S. The role of platelets in the pathogenesis of cerebral malaria. *Cell Mol Life Sci* 2010;67(4):557–68.
- [24] Kasirer-Friede A, Kahn ML, Shattil SJ. Platelet integrins and immunoreceptors. *Immunol Rev* 2007;218:247–64.
- [25] Warkentin TE. Heparin-induced thrombocytopenia: pathogenesis and management. *Br J Haematol* 2003;121(4):535–55.
- [26] Watson SP, Herbert JM, Pollitt AY. GPVI and CLEC-2 in hemostasis and vascular integrity. *J Thromb Haemost* 2010;8(7):1456–67.
- [27] Suzuki-Inoue K, Inoue O, Ozaki Y. Novel platelet activation receptor CLEC-2: from discovery to prospects. *J Thromb Haemost* 2011;9(Suppl. 1):44–55.
- [28] Suzuki-Inoue K, Fuller GL, Garcia A, Eble JA, Pohlmann S, Inoue O, et al. A novel Syk-dependent mechanism of platelet activation by the C-type lectin receptor CLEC-2. *Blood* 2006;107(2):542–9.
- [29] Suzuki-Inoue K, Inoue O, Ding G, Nishimura S, Hokamura K, Eto K, et al. Essential *in vivo* roles of the C-type lectin receptor CLEC-2: embryonic/neonatal lethality of CLEC-2-deficient mice by blood/lymphatic misconnections and impaired thrombus formation of CLEC-2-deficient platelets. *J Biol Chem* 2010;285(32):24494–507.
- [30] Kawai T, Akira S. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat Immunol* 2010;11(5):373–84.
- [31] Akira S, Uematsu S, Takeuchi O. Pathogen recognition and innate immunity. *Cell* 2006;124(4):783–801.
- [32] Semple JW, Aslam R, Kim M, Speck ER, Freedman J. Platelet-bound lipopolysaccharide enhances Fc receptor-mediated phagocytosis of IgG-opsonized platelets. *Blood* 2007;109(11):4803–5.
- [33] Aslam R, Speck ER, Kim M, Crow AR, Bang KW, Nestel FP, et al. Platelet Toll-like receptor expression modulates lipopolysaccharide-induced thrombocytopenia and tumor necrosis factor- α production *in vivo*. *Blood* 2006;107(2):637–41.
- [34] Stahl AL, Svensson M, Morgelin M, Svanborg C, Tarr PI, Mooney JC, et al. Lipopolysaccharide from enterohemorrhagic *Escherichia coli* binds to platelets through TLR4 and CD62 and is detected on circulating platelets in patients with hemolytic uremic syndrome. *Blood* 2006;108(1):167–76.

- [35] Andonegui G, Kerfoot SM, McNagny K, Ebbert KV, Patel KD, Kubes P. Platelets express functional Toll-like receptor-4. *Blood* 2005;106(7):2417–23.
- [36] Clark SR, Ma AC, Tavener SA, McDonald B, Goodarzi Z, Kelly MM, et al. Platelet TLR4 activates neutrophil extracellular traps to ensnare bacteria in septic blood. *Nat Med* 2007;13(4):463–9.
- [37] Blair P, Rex S, Vitseva O, Beaulieu L, Tanriverdi K, Chakrabarti S, et al. Stimulation of Toll-like receptor 2 in human platelets induces a thromboinflammatory response through activation of phosphoinositide 3-kinase. *Circ Res* 2009;104(3):346–54.
- [38] Keane C, Tilley D, Cunningham A, Smolenski A, Kadioglu A, Cox D, et al. Invasive *Streptococcus pneumoniae* trigger platelet activation via Toll-like receptor 2. *J Thromb Haemost* 2010;8(12):2757–65.
- [39] Assinger A, Kral JB, Yaiw KC, Schrottmaier WC, Kurzejamska E, Wang Y, et al. Human cytomegalovirus-platelet interaction triggers toll-like receptor 2-dependent proinflammatory and proangiogenic responses. *Arterioscler Thromb Vasc Biol* 2014;34(4):801–9.
- [40] Panigrahi S, Ma Y, Hong L, Gao D, West XZ, Salomon RG, et al. Engagement of platelet toll-like receptor 9 by novel endogenous ligands promotes platelet hyperreactivity and thrombosis. *Circ Res* 2013;112(1):103–12.
- [41] Thon JN, Peters CG, Machlus KR, Aslam R, Rowley J, Macleod H, et al. T granules in human platelets function in TLR9 organization and signaling. *J Cell Biol* 2012;198(4):561–74.
- [42] Antoniadis C, Bakogiannis C, Tousoulis D, Antonopoulos AS, Stefanadis C. The CD40/CD40 ligand system: linking inflammation with atherothrombosis. *J Am Coll Cardiol* 2009;54(8):669–77.
- [43] Henn V, Slupsky JR, Grafe M, Anagnostopoulos I, Forster R, Muller-Berghaus G, et al. CD40 ligand on activated platelets triggers an inflammatory reaction of endothelial cells. *Nature* 1998;391(6667):591–4.
- [44] Heeschen C, Dimmeler S, Hamm CV, van den Brand MJ, Boersma E, Zeiher AM, et al. Soluble CD40 ligand in acute coronary syndromes. *N Engl J Med* 2003;348(12):1104–11.
- [45] Schonbeck U, Libby P. The CD40/CD154 receptor/ligand dyad. *Cell Mol Life Sci* 2001;58(1):4–43.
- [46] Elzey BD, Ratliff TL, Sowa JM, Crist SA. Platelet CD40L at the interface of adaptive immunity. *Thromb Res* 2011;127(3):180–3.
- [47] Kroccek RA, Graf D, Brugnani D, Giliani S, Korthuer U, Ugazio A, et al. Defective expression of CD40 ligand on T cells causes "X-linked immunodeficiency with hyper-IgM (HIGM1)". *Immunol Rev* 1994;138:39–59.
- [48] Allen RC, Armitage RJ, Conley ME, Rosenblatt H, Jenkins NA, Copeland NG, et al. CD40 ligand gene defects responsible for X-linked hyper-IgM syndrome. *Science* 1993;259(5097):990–3.
- [49] Khan SY, Kelher MR, Heal JM, Blumberg N, Boshkov LK, Phipps R, et al. Soluble CD40 ligand accumulates in stored blood components, primes neutrophils through CD40, and is a potential cofactor in the development of transfusion-related acute lung injury. *Blood* 2006;108(7):2455–62.
- [50] Schonbeck U, Libby P. CD40 signaling and plaque instability. *Circ Res* 2001;89(12):1092–103.
- [51] Gawaz M, Neumann FJ, Ott I, Schiessler A, Schomig A. Platelet function in acute myocardial infarction treated with direct angioplasty. *Circulation* 1996;93(2):229–37.
- [52] Gawaz M, Langer H, May AE. Platelets in inflammation and atherogenesis. *J Clin Invest* 2005;115(12):3378–84.
- [53] Frenette PS, Denis CV, Weiss L, Jurk K, Subbarao S, Kehrel B, et al. P-Selectin glycoprotein ligand 1 (PSGL-1) is expressed on platelets and can mediate platelet-endothelial interactions in vivo. *J Exp Med* 2000;191(8):1413–22.
- [54] Romo GM, Dong JF, Schade AJ, Gardiner EE, Kansas GS, Li CQ, et al. The glycoprotein Ib-IX-V complex is a platelet counterreceptor for P-selectin. *J Exp Med* 1999;190(6):803–14.
- [55] Davi G, Patrono C. Platelet activation and atherothrombosis. *N Engl J Med* 2007;357(24):2482–94.
- [56] Gawaz M, Langer H, May A. Platelets in inflammation and atherogenesis. *J Clin Invest* 2005;115(12):3378–84.
- [57] Massberg S, Brand K, Gruner S, Page S, Muller E, Muller I, et al. A critical role of platelet adhesion in the initiation of atherosclerotic lesion formation. *J Exp Med* 2002;196(7):887–96.
- [58] Belton OA, Duffy A, Toomey S, Fitzgerald DJ. Cyclooxygenase isoforms and platelet vessel wall interactions in the apolipoprotein E knockout mouse model of atherosclerosis. *Circulation* 2003;108(24):3017–23.
- [59] McEver RP, Cummings RD. Perspectives series: cell adhesion in vascular biology. Role of PSGL-1 binding to selectins in leukocyte recruitment. *J Clin Invest* 1997;100(3):485–91.
- [60] Simon DI, Chen Z, Xu H, Li CQ, Dong J, McIntire LV, et al. Platelet glycoprotein Ib α is a counterreceptor for the leukocyte integrin Mac-1 (CD11b/CD18). *J Exp Med* 2000;192(2):193–204.
- [61] Diacovo TG, deFougerolles AR, Bainton DF, Springer TA. A functional integrin ligand on the surface of platelets: intercellular adhesion molecule-2. *J Clin Invest* 1994;94(3):1243–51.
- [62] Santos S, Sachs UJ, Kroll H, Linder M, Ruf A, Preissner KT, et al. The junctional adhesion molecule 3 (JAM-3) on human platelets is a counterreceptor for the leukocyte integrin Mac-1. *J Exp Med* 2002;196(5):679–91.
- [63] Weber C, Springer TA. Neutrophil accumulation on activated, surface-adherent platelets in flow is mediated by interaction of Mac-1 with fibrinogen bound to α IIb β 3 and stimulated by platelet-activating factor. *J Clin Invest* 1997;100(8):2085–93.
- [64] Gleissner CA, von Hundelshausen P, Ley K. Platelet chemokines in vascular disease. *Arterioscler Thromb Vasc Biol* 2008;28(11):1920–7.
- [65] Zlotnik A, Yoshie O. Chemokines: a new classification system and their role in immunity. *Immunity* 2000;12(2):121–7.
- [66] Klinger MH, Wilhelm D, Babel S, Sticherling M, Schroder JM, Kuehn W. Immunocytochemical localization of the chemokines RANTES and MIP-1 α within human platelets and their release during storage. *Int Arch Allergy Immunol* 1995;107(4):541–6.
- [67] Smyth SS, McEver RP, Weyrich AS, Morrell CN, Hoffman MR, Arepally GM, et al. Platelet functions beyond hemostasis. *J Thromb Haemost* 2009;7(11):1759–66.
- [68] Ghasemzadeh M, Kaplan ZS, Alwis I, Schoenwaelder SM, Ashworth KJ, Westein E, et al. The CXCR1/2 ligand NAP-2 promotes directed intravascular leukocyte migration through platelet thrombi. *Blood* 2013;121(22):4555–66.
- [69] von Hundelshausen P, Petersen F, Brandt E. Platelet-derived chemokines in vascular biology. *Thromb Haemost* 2007;97(5):704–13.
- [70] Holt JC, Rabellino EM, Gewirtz AM, Gunkel LM, Rucinski B, Niewiarowski S. Occurrence of platelet basic protein, a precursor of low affinity platelet factor 4 and beta-thromboglobulin, in human platelets and megakaryocytes. *Exp Hematol* 1988;16(4):302–6.
- [71] Brandt E, Van Damme J, Flad HD. Neutrophils can generate their activator neutrophil-activating peptide 2 by proteolytic cleavage of platelet-derived connective tissue-activating peptide III. *Cytokine* 1991;3(4):311–21.
- [72] Schiemann F, Grimm TA, Hoch J, Gross R, Lindner B, Petersen F, et al. Mast cells and neutrophils proteolytically activate chemokine precursor CTAP-III and are subject to counterregulation by PF-4 through inhibition of chymase and cathepsin G. *Blood* 2006;107(6):2234–42.
- [73] von Hundelshausen P, Koenen RR, Weber C. Platelet-mediated enhancement of leukocyte adhesion. *Microcirculation* 2009;16(1):84–96.
- [74] Walz A, Baggiolini M. A novel cleavage product of beta-thromboglobulin formed in cultures of stimulated mononuclear cells activates human neutrophils. *Biochem Biophys Res Commun* 1989;159(3):969–75.
- [75] Brandt E, Ludwig A, Petersen F, Flad HD. Platelet-derived CXC chemokines: old players in new games. *Immunol Rev* 2000;177:204–16.
- [76] Hagiwara H, Mitsumata M, Yamane T, Jin X, Yoshida Y. Laminar shear stress-induced GRO mRNA and protein expression in endothelial cells. *Circulation* 1998;98(23):2584–90.
- [77] Lira SA, Furtado GC. The biology of chemokines and their receptors. *Immunol Res* 2012;54(1–3):111–20.
- [78] Smith DF, Galkina E, Ley K, Huo Y. GRO family chemokines are specialized for monocyte arrest from flow. *Am J Physiol Heart Circ Physiol* 2005;289(5):H1976–84.
- [79] Power CA, Furness RB, Brawand C, Wells TN. Cloning of a full-length cDNA encoding the neutrophil-activating peptide ENA-78 from human platelets. *Gene* 1994;151(1–2):333–4.
- [80] Gear AR, Camerini D. Platelet chemokines and chemokine receptors: linking hemostasis, inflammation, and host defense. *Microcirculation* 2003;10(3–4):335–50.
- [81] Moser B, Clark-Lewis I, Zwahlen R, Baggiolini M. Neutrophil-activating properties of the melanoma growth-stimulatory activity. *J Exp Med* 1990;171(5):1797–802.
- [82] Holmes WE, Lee J, Kuang WJ, Rice GC, Wood WI. Structure and functional expression of a human interleukin-8 receptor. *Science* 1991;253(5025):1278–80.
- [83] Boehlen F, Clemetson KJ. Platelet chemokines and their receptors: what is their relevance to platelet storage and transfusion practice? *Transfus Med* 2001;11(6):403–17.
- [84] Rollins BJ. Chemokines. *Blood* 1997;90(3):909–28.
- [85] Kaplan ZS, Jackson SP. The role of platelets in atherothrombosis. *Hematology Am Soc Hematol Educ Program* 2011;2011:51–61.
- [86] Jennings LK. Role of platelets in atherothrombosis. *Am J Cardiol* 2009;103(3 Suppl):4A–10A.
- [87] Schonbeck U, Sukhova GK, Shimizu K, Mach F, Libby P. Inhibition of CD40 signaling limits evolution of established atherosclerosis in mice. *Proc Natl Acad Sci U S A* 2000;97(13):7458–63.
- [88] Lutgens E, Cleutjens KB, Heeneman S, Kotliansky VE, Burkly LC, Daemen MJ. Both early and delayed anti-CD40L antibody treatment induces a stable plaque phenotype. *Proc Natl Acad Sci U S A* 2000;97(13):7464–9.
- [89] Shaz BH, Stowell SR, Hillyer CD. Transfusion-related acute lung injury: from bedside to bench and back. *Blood* 2011;117(5):1463–71.
- [90] Bux J, Sachs UJ. The pathogenesis of transfusion-related acute lung injury (TRALI). *Br J Haematol* 2007;136(6):788–99.
- [91] Caudrillier A, Kessenbrock K, Gilliss BM, Nguyen JX, Marques MB, Monestier M, et al. Platelets induce neutrophil extracellular traps in transfusion-related acute lung injury. *J Clin Invest* 2012;122(7):2661–71.
- [92] Chang M, Nakagawa PA, Williams SA, et al. Immune thrombocytopenic purpura (ITP) plasma and purified ITP monoclonal autoantibodies inhibit megakaryocytopoiesis in vitro. *Blood* 2003;102(3):887–95.
- [93] Harrington WJ, Minnich V, Hollingsworth JW, Moore CV. Demonstration of a thrombocytopenic factor in the blood of patients with thrombocytopenic purpura. *J Lab Clin Med* 1951;38(1):1–10.
- [94] Asahi A, Nishimoto T, Okazaki Y, Suzuki H, Masaoka T, Kawakami Y, et al. *Helicobacter pylori* eradication shifts monocyte Fc γ receptor balance toward inhibitory Fc γ RIIB in immune thrombocytopenic purpura patients. *J Clin Invest* 2008;118(8):2939–49.
- [95] Blajchman MA. Transfusion-associated immunomodulation and universal white cell reduction: are we putting the cart before the horse? *Transfusion* 1999;39(7):665–70.
- [96] Gouttefangeas C, Diehl M, Keilholz W, Hornlein RF, Stevanovic S, Rammensee HG. Thrombocyte HLA molecules retain nonrenewable endogenous peptides of megakaryocyte lineage and do not stimulate direct alloreactivity in vitro. *Blood* 2000;95(10):3168–75.

- [97] Chapman LM, Aggrey AA, Field DJ, Srivastava K, Ture S, Yui K, et al. Platelets present antigen in the context of MHC class I. *J Immunol* 2012;189(2):916–23.
- [98] Aslam R, Speck ER, Kim M, Freedman J, Semple JW. Transfusion-related immunomodulation by platelets is dependent on their expression of MHC Class I molecules and is independent of white cells. *Transfusion* 2008;48(9):1778–86.
- [99] Freedman JE, Loscalzo J. Platelet-monocyte aggregates: bridging thrombosis and inflammation. *Circulation* 2002;105(18):2130–2.
- [100] van Gils JM, Zwaginga JJ, Hordijk PL. Molecular and functional interactions among monocytes, platelets, and endothelial cells and their relevance for cardiovascular diseases. *J Leukoc Biol* 2009;85(2):195–204.
- [101] Brinkmann V, Reichard U, Goosmann C, Fauler B, Uhlemann Y, Weiss DS, et al. Neutrophil extracellular traps kill bacteria. *Science* 2004;303(5663):1532–5.
- [102] Kaplan MJ, Radic M. Neutrophil extracellular traps: double-edged swords of innate immunity. *J Immunol* 2012;189(6):2689–95.
- [103] Brill A, Fuchs TA, Savchenko AS, Thomas GM, Martinod K, De Meyer SF, et al. Neutrophil extracellular traps promote deep vein thrombosis in mice. *J Thromb Haemost* 2012;10(1):136–44.
- [104] Fuchs TA, Brill A, Dueschmied D, Schatzberg D, Monestier M, Myers Jr DD, et al. Extracellular DNA traps promote thrombosis. *Proc Natl Acad Sci U S A* 2010;107(36):15880–5.
- [105] von Bruhl ML, Stark K, Steinhart A, Chandraratne S, Konrad I, Lorenz M, et al. Monocytes, neutrophils, and platelets cooperate to initiate and propagate venous thrombosis in mice in vivo. *J Exp Med* 2012;209(4):819–35.
- [106] Brinkmann V, Zychlinsky A. Beneficial suicide: why neutrophils die to make NETs. *Nat Rev Microbiol* 2007;5(8):577–82.
- [107] Ma AC, Kubes P. Platelets, neutrophils, and neutrophil extracellular traps (NETs) in sepsis. *J Thromb Haemost* 2008;6(3):415–20.
- [108] Phillipson M, Kubes P. The neutrophil in vascular inflammation. *Nat Med* 2011;17(11):1381–90.
- [109] Reid VL, Webster NR. Role of microparticles in sepsis. *Br J Anaesth* 2012;109(4):503–13.
- [110] Ogura H, Kawasaki T, Tanaka H, Koh T, Tanaka R, Ozeki Y, et al. Activated platelets enhance microparticle formation and platelet-leukocyte interaction in severe trauma and sepsis. *J Trauma* 2001;50(5):801–9.
- [111] Burnier L, Fontana P, Kwak BR, Angelillo-Scherrer A. Cell-derived microparticles in haemostasis and vascular medicine. *Thromb Haemost* 2009;101(3):439–51.
- [112] Italiano Jr JE, Mairuhu AT, Flaumenhaft R. Clinical relevance of microparticles from platelets and megakaryocytes. *Curr Opin Hematol* 2010;17(6):578–84.
- [113] Mause SF, von Hundelshausen P, Zernecke A, Koenen RR, Weber C. Platelet microparticles: a transcellular delivery system for RANTES promoting monocyte recruitment on endothelium. *Arterioscler Thromb Vasc Biol* 2005;25(7):1512–8.
- [114] Edelstein LC, McKenzie SE, Shaw C, Holinstat MA, Kunapuli SP, Bray PF. MicroRNAs in platelet production and activation. *J Thromb Haemost* 2013;11(Suppl 1):340–50.
- [115] Edelstein LC, Bray PF. MicroRNAs in platelet production and activation. *Blood* 2011;117(20):5289–96.
- [116] Risitano A, Beaulieu LM, Vitseva O, Freedman JE. Platelets and platelet-like particles mediate intercellular RNA transfer. *Blood* 2012;119(26):6288–95.
- [117] Tranum BL, Haut A. Thrombocytosis: platelet kinetics in neoplasia. *J Lab Clin Med* 1974;84(5):615–9.
- [118] Hernandez E, Lavine M, Dunton CJ, Gracely E, Parker J. Poor prognosis associated with thrombocytosis in patients with cervical cancer. *Cancer* 1992;69(12):2975–7.
- [119] Zeimet AG, Marth C, Muller-Holzner E, Daxenbichler G, Dapunt O. Significance of thrombocytosis in patients with epithelial ovarian cancer. *Am J Obstet Gynecol* 1994;170(2):549–54.
- [120] Ikeda M, Furukawa H, Imamura H, Shimizu J, Ishida H, Masutani S, et al. Poor prognosis associated with thrombocytosis in patients with gastric cancer. *Ann Surg Oncol* 2002;9(3):287–91.
- [121] Karparkin S, Pearlstein E, Ambrogio C, Collier BS. Role of adhesive proteins in platelet tumor interaction in vitro and metastasis formation in vivo. *J Clin Invest* 1988;81(4):1012–9.
- [122] Gasic GJ, Gasic TB, Stewart CC. Antimetastatic effects associated with platelet reduction. *Proc Natl Acad Sci U S A* 1968;61(1):46–52.
- [123] Karparkin S, Ambrogio C, Pearlstein E. The role of tumor-induced platelet aggregation, platelet adhesion and adhesive proteins in tumor metastasis. *Prog Clin Biol Res* 1988;283:585–606.
- [124] Shau H, Roth MD, Golub SH. Regulation of natural killer function by nonlymphoid cells. *Nat Immun* 1993;12(4–5):235–49.
- [125] Philippe C, Philippe B, Fouqueray B, Perez J, Lebret M, Baud L. Protection from tumor necrosis factor–mediated cytotoxicity by platelets. *Am J Pathol* 1993;143(6):1713–23.
- [126] Stone JP, Wagner DD. P-selectin mediates adhesion of platelets to neuroblastoma and small cell lung cancer. *J Clin Invest* 1993;92(2):804–13.
- [127] Bambace NM, Holmes CE. The platelet contribution to cancer progression. *J Thromb Haemost* 2011;9(2):237–49.
- [128] Erpenbeck L, Schon MP. Deadly allies: the fatal interplay between platelets and metastasizing cancer cells. *Blood* 2010;115(17):3427–36.
- [129] Trikha M, Zhou Z, Timar J, Raso E, Kennel M, Emmell E, et al. Multiple roles for platelet GPIIb/IIIa and alpha5beta3 integrins in tumor growth, angiogenesis, and metastasis. *Cancer Res* 2002;62(10):2824–33.
- [130] Schumacher Jr HR. Synovial membrane and fluid morphologic alterations in early rheumatoid arthritis: microvascular injury and virus-like particles. *Ann N Y Acad Sci* 1975;256:39–64.
- [131] Wang F, Wang NS, Yan CG, Li JH, Tang LQ. The significance of platelet activation in rheumatoid arthritis. *Clin Rheumatol* 2007;26(5):768–71.
- [132] Knijff-Dutmer EA, Koerts J, Nieuwland R, Kalsbeek-Batenburg EM, van de Laar MA. Elevated levels of platelet microparticles are associated with disease activity in rheumatoid arthritis. *Arthritis Rheum* 2002;46(6):1498–503.
- [133] Bunesco A, Seideman P, Lenkei R, Levin K, Egberg N. Enhanced Fc gamma receptor I, alpha5beta3 integrin receptor expression by monocytes and neutrophils in rheumatoid arthritis: interaction with platelets. *J Rheumatol* 2004;31(12):2347–55.
- [134] Sellam J, Proulle V, Jungel A, Ittah M, Miceli Richard C, Gottenberg JE, et al. Increased levels of circulating microparticles in primary Sjogren's syndrome, systemic lupus erythematosus and rheumatoid arthritis and relation with disease activity. *Arthritis Res Ther* 2009;11(5):R156.
- [135] Yaron M, Djalldetti M. Platelets in synovial fluid. *Arthritis Rheum* 1978;21(5):607–8.
- [136] Endresen GK, Forre O. Human platelets in synovial fluid. A focus on the effects of growth factors on the inflammatory responses in rheumatoid arthritis. *Clin Exp Rheumatol* 1992;10(2):181–7.
- [137] Boilard E, Nigrovic PA, Larabee K, Watts GF, Coblyn JS, Weinblatt ME, et al. Platelets amplify inflammation in arthritis via collagen-dependent microparticle production. *Science* 2010;327(5965):580–3.
- [138] Boilard E, Blanco P, Nigrovic PA. Platelets: active players in the pathogenesis of arthritis and SLE. *Nat Rev Rheumatol* 2012;8(9):534–42.
- [139] Flaumenhaft R. Platelet secretion. In: Michelson AD, editor. *Platelets*. Amsterdam: Academic Press; 2013. p. 343–66.
- [140] McNicol A, Israels SJ. Platelet dense granules: structure, function and implications for haemostasis. *Thromb Res* 1999;95(1):1–18.
- [141] Rendu F, Brohard-Bohn B. The platelet release reaction: granules' constituents, secretion and functions. *Platelets* 2001;12(5):261–73.
- [142] Blair P, Flaumenhaft R. Platelet alpha-granules: basic biology and clinical correlates. *Blood Rev* 2009;23(4):177–89.
- [143] Vasina E, Heemskerk JW, Weber C, Koenen RR. Platelets and platelet-derived microparticles in vascular inflammatory disease. *Inflamm Allergy Drug Targets* 2010;9(5):346–54.
- [144] Bizzozero J. Ueber einen neuen Formbestandteil des Blutes und dessen Rolle bei der Thrombose und Blutgerinnung. *Archiv für pathologische Anatomie und Physiologie und für klinische Medizin* 1882;90:261–332.