

A broad learning system for ^{18}F -FDG PET/MRI imaging diagnosis in temporal lobe epilepsy patients

Huanhua Wu^{1,1,1,*}, Bob Security^b, Junwei Duan^{b,2,*}, Hao Xu^{b,1,*}, Derek Zoolander^{a,2}

^aDepartment Street City State Zip

^bDepartment Street City State Zip

Abstract

This is the abstract.

It consists of two paragraphs.

Keywords: Epilepsy, Broad Learning System, Positron emission tomography, MRI

1. Abstract

China has about 10 million people with epilepsy(Beghi et al., 2019; Ding et al., 2021).

2. Introduction

To develop and validate a radiomics nomogram based on ^{18}F -FDG PET/MRI for individualised seizure prediction after temporal lobe epilepsy surgery.

Approximately half of patients who undergo resective brain surgery for drug-resistant epilepsy have recurrent seizures after surgery(Beghi et al., 2019). China has about 10 million people with epilepsy(Beghi et al., 2019; Ding et al., 2021). Several single predictors have been identified, however, no validated method has been developed to determine a patient's seizure outcome based on their complex clinical characteristics and ^{18}F -FDG PET/MRI radiomics signature(Jehi et al., 2015). The treatment of medically intractable focal epilepsy involves resection of the brain every year. There are probably thousands of others who would benefit from the procedure, but are not referred for surgery. Due to the unpredictability of seizure outcomes after epilepsy surgery, both groups of patients face difficulties.

Radiomics is an emerging method for imaging analysis that uses algorithms or statistical tools to determine phenotypic differences between lesions. Using radiomics features, morphology and function can be sensitively evaluated beyond the naked eye and can provide a great deal of additional information. The radiomics features can provide a great deal of information beyond analysis with the naked eye and can sensitively determine the subtle heterogeneity of the morphology and function on a cellular level, between different parts of the lesions.

The validity of these nomograms in prospective cohort studies could help predict seizure outcomes in patients who are eligible for epilepsy surgery(Cheong et al., 2021; Zhang et al., 2021).

*Corresponding author

Email addresses: wane199@outlook.com (Huanhua Wu), bob@example.com (Bob Security), cat@example.com (Junwei Duan), cat@example.com (Hao Xu), derek@example.com (Derek Zoolander)

¹This is the first author footnote.

²Another author footnote.




Figure 1: A meaningless scatterplot.

3. Equations

Here is another:

$$AI = c^2. \tag{1}$$

Inline equations: $\sum_{i=2}^{\infty} \{\alpha_i^\beta\}$

4. Figures and tables

Figure 1 is generated using an R chunk.

5. Tables coming from R

Tables can also be generated using R chunks, as shown in Table 1 for example.

```
knitr::kable(head(mtcars)[,1:4],
  caption = "\\label{tab1}Caption centered above table"
)
```

Table 1: Caption centered above table

| | mpg | cyl | disp | hp |
|----------------|------|-----|------|-----|
| Mazda RX4 | 21.0 | 6 | 160 | 110 |
| Mazda RX4 Wag | 21.0 | 6 | 160 | 110 |
| Datsun 710 | 22.8 | 4 | 108 | 93 |
| Hornet 4 Drive | 21.4 | 6 | 258 | 110 |

| | mpg | cyl | disp | hp |
|-------------------|------|-----|------|-----|
| Hornet Sportabout | 18.7 | 8 | 360 | 175 |
| Valiant | 18.1 | 6 | 225 | 105 |

References

- Ettore Beghi, Giorgia Giussani, Emma Nichols, Foad Abd-Allah, Jemal Abdela, Ahmed Abdelalim, Haftom Niguse Abraha, Mina G Adib, Sutapa Agrawal, Fares Alahdab, et al. Global, regional, and national burden of epilepsy, 1990–2016: a systematic analysis for the global burden of disease study 2016. *The Lancet Neurology*, 18(4):357–375, 2019.
- E-Nae Cheong, Ji Eun Park, Da Eun Jung, and Woo Hyun Shim. Extrahippocampal radiomics analysis can potentially identify laterality in patients with mri-negative temporal lobe epilepsy. *Frontiers in neurology*, 12:1277, 2021.
- Ding Ding, Dong Zhou, Josemir W Sander, Wenzhi Wang, Shichuo Li, and Zhen Hong. Epilepsy in china: major progress in the past two decades. *The Lancet Neurology*, 20(4):316–326, 2021.
- Lara Jehi, Ruta Yardi, Kevin Chagin, Laura Tassi, Giorgio Lo Russo, Gregory Worrell, Wei Hu, Fernando Cendes, Marcia Morita, Fabrice Bartolomei, et al. Development and validation of nomograms to provide individualised predictions of seizure outcomes after epilepsy surgery: a retrospective analysis. *The Lancet Neurology*, 14(3):283–290, 2015.
- Qinming Zhang, Yi Liao, Xiawan Wang, Teng Zhang, Jianhua Feng, Jianing Deng, Kexin Shi, Lin Chen, Liu Feng, Mindi Ma, et al. A deep learning framework for 18f-fdg pet imaging diagnosis in pediatric patients with temporal lobe epilepsy. *European journal of nuclear medicine and molecular imaging*, 48(8):2476–2485, 2021.