

Stein-Weiss inequality on product spaces

Zipeng Wang

In communication with Charles Fefferman

Abstract

We give the characterization between weighted norm inequalities of strong fractional integral operators and their associated multi parameter Muckenhoupt characteristics, by considering the weights to be power functions. As a result, we extend the classical Stein-Weiss theorem to product spaces.

1 Introduction

Let $0 < \alpha < \mathbf{N}$. A fractional integral operator I_α is defined by

$$I_\alpha f(x) = \int_{\mathbb{R}^{\mathbf{N}}} f(y) \left[\frac{1}{|x-y|} \right]^{\mathbf{N}-\alpha} dy. \quad (1.1)$$

In 1928, Hardy and Littlewood [1] first established a two-weight $L^p \rightarrow L^q$ -norm inequality for I_α in the one dimensional space, by considering the *weights* as suitable powers. This result has been extended to higher dimensions by Stein and Weiss [3]. Today, it bears the name of Stein-Weiss inequality.

◇ Throughout, $\mathfrak{B} > 0$ is regarded as a generic constant depending on its subindices.

Theorem A: Stein and Weiss (1958) *Let I_α defined in (1.1) for $0 < \alpha < \mathbf{N}$. Suppose that $\omega(x) = |x|^{-\gamma}$, $\sigma(x) = |x|^\delta$ for $x \neq 0$ and $\gamma, \delta \in \mathbb{R}$. We have*

$$\|\omega I_\alpha f\|_{L^q(\mathbb{R}^{\mathbf{N}})} \leq \mathfrak{B}_{p,q,\alpha,\gamma,\delta} \|f\sigma\|_{L^p(\mathbb{R}^{\mathbf{N}})} \quad (1.2)$$

for $1 < p \leq q < \infty$ if

$$\gamma < \frac{\mathbf{N}}{q}, \quad \delta < \mathbf{N} \left(\frac{p-1}{p} \right), \quad \gamma + \delta \geq 0 \quad (1.3)$$

and

$$\frac{\alpha}{\mathbf{N}} = \frac{1}{p} - \frac{1}{q} + \frac{\gamma + \delta}{\mathbf{N}}. \quad (1.4)$$

In the case of $\gamma = \delta = 0$, **Theorem A** was proved in $\mathbb{R}^{\mathbf{N}}$ by Sobolev [2]. This is also known as Hardy-Littlewood-Sobolev inequality.

Fractional integration in weighted norms have been systematically studied over the several past decades. For instance, see Muckenhoupt and Wheeden [5], Coifman and Fefferman [9], Fefferman and Muckenhoupt [8], Pérez [10] and Sawyer and Wheeden [6].

Denote Q to be a cube in \mathbb{R}^N parallel to coordinates. As a well known estimate, (1. 2) implies

$$\sup_{Q \subset \mathbb{R}^N} |Q|^{\frac{\alpha}{N} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|Q|} \int_Q \omega^q(x) dx \right\}^{\frac{1}{q}} \left\{ \frac{1}{|Q|} \int_Q \left(\frac{1}{\sigma} \right)^{\frac{p}{p-1}}(x) dx \right\}^{\frac{p-1}{p}} < \infty. \quad (1. 5)$$

The supremum in (1. 5) is called Muckenhoupt characteristic, as was first introduced by Muckenhoupt for which ω^q and $\sigma^{-\frac{p}{p-1}}$ are nonnegative and locally integrable.

Take into account $\omega(x) = |x|^{-\gamma}$, $\sigma(x) = |x|^\delta$ for $x \neq 0$ and $\gamma, \delta \in \mathbb{R}$. We find the two-weight $\mathbf{L}^p \rightarrow \mathbf{L}^q$ -norm inequality in (1. 5) implying the constraints in (1. 3)-(1. 4). Hence that (1. 2), (1. 3)-(1. 4) and (1. 5) are in fact equivalent conditions.

Now, consider $\mathbb{R}^N = \mathbb{R}^{N_1} \times \mathbb{R}^{N_2} \times \dots \times \mathbb{R}^{N_n}$ as a product space. Let

$$0 < \alpha_i < N_i, \quad i = 1, 2, \dots, n \quad \text{and} \quad \alpha = \alpha_1 + \alpha_2 + \dots + \alpha_n. \quad (1. 6)$$

In this paper, we give an extension of **Theorem A** to multi-parameter settings by asserting the *strong* fractional integral operator \mathbf{I}_α defined by

$$\mathbf{I}_\alpha f(x) = \int_{\mathbb{R}^N} f(y) \prod_{i=1}^n \left[\frac{1}{|x_i - y_i|} \right]^{N_i - \alpha_i} dy. \quad (1. 7)$$

Observe that the kernel of \mathbf{I}_α has singularity appeared on every coordinate subspace.

The study of certain operators that commute with a multi-parameter family of dilations, dates back to the time of Jessen, Marcinkiewicz and Zygmund. A number of pioneering results have been accomplished, for example by Cordoba and Fefferman [11], Fefferman [12], Fefferman and Stein [13], Müller, Ricci and Stein [14], Journé [15] and Pipher [16]. This area remains largely open for fractional integration.

2 Statement of the main result

Denote $\mathbf{Q} = \mathbf{Q}_1 \times \mathbf{Q}_2 \times \dots \times \mathbf{Q}_n \subset \mathbb{R}^{N_1} \times \mathbb{R}^{N_2} \times \dots \times \mathbb{R}^{N_n} = \mathbb{R}^N$ where \mathbf{Q}_i is a cube in \mathbb{R}^{N_i} parallel to coordinates for every $i = 1, 2, \dots, n$.

Theorem A*: Let \mathbf{I}_α defined in (1. 6)-(1. 7). Suppose $\omega(x) = |x|^{-\gamma}$, $\sigma(x) = |x|^\delta$, $\gamma, \delta \in \mathbb{R}$ at $x \neq 0$. For $1 < p \leq q < \infty$, the following conditions are equivalent:

1.

$$\sup_{\mathbf{Q} \subset \mathbb{R}^N} \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{N_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \omega^q(x) dx \right\}^{\frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{\sigma} \right)^{\frac{p}{p-1}}(x) dx \right\}^{\frac{p-1}{p}} < \infty. \quad (2. 1)$$

2.

$$\gamma < \frac{N}{q}, \quad \delta < N \left(\frac{p-1}{p} \right), \quad \gamma + \delta \geq 0 \quad (2. 2)$$

and

$$\frac{\alpha}{N} = \frac{1}{p} - \frac{1}{q} + \frac{\gamma + \delta}{N}. \quad (2. 3)$$

For $\gamma \geq 0, \delta \leq 0$,

$$\alpha_i - \frac{\mathbf{N}_i}{p} < \delta, \quad i = 1, 2, \dots, n. \quad (2.4)$$

For $\gamma \leq 0, \delta \geq 0$,

$$\alpha_i - \mathbf{N}_i \left(\frac{q-1}{q} \right) < \gamma, \quad i = 1, 2, \dots, n. \quad (2.5)$$

For $\gamma > 0, \delta > 0$,

$$\begin{aligned} \sum_{i \in \mathcal{U}} \alpha_i - \frac{\mathbf{N}_i}{p} < \delta, \quad \mathcal{U} &= \left\{ i \in \{1, 2, \dots, n\} : \alpha_i - \frac{\mathbf{N}_i}{p} \geq 0 \right\}, \\ \sum_{i \in \mathcal{V}} \alpha_i - \left(\frac{q-1}{q} \right) \mathbf{N}_i < \gamma, \quad \mathcal{V} &= \left\{ i \in \{1, 2, \dots, n\} : \alpha_i - \mathbf{N}_i \left(\frac{q-1}{q} \right) \geq 0 \right\}. \end{aligned} \quad (2.6)$$

3.

$$\|\omega \mathbf{I}_\alpha f\|_{\mathbf{L}^q(\mathbb{R}^N)} \leq \mathfrak{B}_{p,q,\alpha,\gamma,\delta} \|f\sigma\|_{\mathbf{L}^p(\mathbb{R}^N)}, \quad 1 < p \leq q < \infty. \quad (2.7)$$

Remark 2.1 In the 2-parameter setting ($n = 2$), **Theorem A*** is first proved in the joint work of Sawyer and Wang [7]. For $\gamma \geq 0, \delta \leq 0$ or $\gamma \leq 0, \delta \geq 0$, the "sandwiching" idea introduced there applies to the general multi-parameter situation. However, the difficult case occurs at $\gamma > 0, \delta > 0$. The method used in [7] relies on solving a system of algebraic equations, which is no longer solvable for $n > 2$.

The remaining paper is organized as follows. In section 3, we introduce a new framework, where the product space is decomposed into an infinitely many of dyadic cones. Every partial sum operator defined on a dyadic cone is essentially an one-parameter fractional integral operator, satisfying the desired regularity.

In section 4, we prove that the multi-parameter Muckenhoupt characteristic in (2.1) implies the constraints in (2.2)-(2.6).

In section 5, by using (2.2)-(2.6), we show that

$$\prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{\mathbf{N}_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \omega^{qr}(x) dx \right\}^{\frac{1}{qr}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{\sigma} \right)^{\frac{pr}{p-1}}(x) dx \right\}^{\frac{p-1}{pr}}, \quad r > 1 \quad (2.8)$$

decays exponentially as the eccentricity of \mathbf{Q} getting large, whenever

$$\alpha_i > \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right), \quad i = 1, 2, \dots, n.$$

On the other hand, we handle the case $\alpha_i = \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)$, $i = 1, 2, \dots, n$ in section 6.

We prove **Theorem A*** in the last section, by decomposing \mathbf{I}_α so that the resulting estimates can be reduced to either of the above two cases.

Because \mathbf{I}_α is positive definite, we assume $f \geq 0$ from now on.

3 Cone decomposition on product spaces

Denote \mathbf{t} as an n -tuple $(2^{-t_1}, 2^{-t_2}, \dots, 2^{-t_n})$ where $t_i, i = 1, 2, \dots, n$ are non-negative integers. We require $t_v \doteq \min\{t_i: i = 1, 2, \dots, n\} = 0$. Define

$$\Delta_{\mathbf{t}} \mathbf{I}_{\alpha} f(x) = \int_{\Gamma_{\mathbf{t}}(x)} f(y) \prod_{i=1}^n \left[\frac{1}{|x_i - y_i|} \right]^{\mathbf{N}_i - \alpha_i} dy \quad (3.1)$$

where

$$\Gamma_{\mathbf{t}}(x) \doteq \bigotimes_{i=1}^n \left\{ y_i \in \mathbb{R}^{\mathbf{N}_i}: 2^{-t_i} \leq \frac{|x_i - y_i|}{|x_v - y_v|} < 2^{-t_i+1} \right\}. \quad (3.2)$$

Observe that $\Gamma_{\mathbf{t}}(x)$ in (3.2) is a dyadic cone with vertex on x whose eccentricity depends on \mathbf{t} . In particular, we write

$$\Gamma_o(x) \doteq \Gamma_{\mathbf{t}}(x), \quad t_1 = t_2 = \dots = t_n = 0. \quad (3.3)$$




Figure 1: dyadic cones in a 2-parameter setting.

Let $\mathbf{Q}^{\mathbf{t}}$ be a dilated of \mathbf{Q} such that $|\mathbf{Q}_i^{\mathbf{t}}|^{\frac{1}{\mathbf{N}_i}} = 2^{-t_i} |\mathbf{Q}_i|^{\frac{1}{\mathbf{N}_i}}, i = 1, 2, \dots, n$.

Write $\mathbf{t}x = (2^{-t_1}x_1, 2^{-t_2}x_2, \dots, 2^{-t_n}x_n)$. We have

$$\begin{aligned}
& \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{N_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \omega^{qr}(\mathbf{t}x) dx \right\}^{\frac{1}{qr}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{\sigma} \right)^{\frac{pr}{p-1}}(\mathbf{t}x) dx \right\}^{\frac{p-1}{pr}} \\
&= \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{N_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}^{\mathbf{t}}|} \int_{\mathbf{Q}^{\mathbf{t}}} \omega^{qr}(x) dx \right\}^{\frac{1}{qr}} \left\{ \frac{1}{|\mathbf{Q}^{\mathbf{t}}|} \int_{\mathbf{Q}^{\mathbf{t}}} \left(\frac{1}{\sigma} \right)^{\frac{pr}{p-1}}(x) dx \right\}^{\frac{p-1}{pr}} \\
&= \prod_{i=1}^n 2^{t_i [\alpha_i - \frac{N_i}{p} + \frac{N_i}{q}]} \prod_{i=1}^n |\mathbf{Q}_i^{\mathbf{t}}|^{\frac{\alpha_i}{N_i} - (\frac{1}{p} - \frac{1}{q})} \left\{ \frac{1}{|\mathbf{Q}^{\mathbf{t}}|} \int_{\mathbf{Q}^{\mathbf{t}}} \omega^{qr}(x) dx \right\}^{\frac{1}{qr}} \left\{ \frac{1}{|\mathbf{Q}^{\mathbf{t}}|} \int_{\mathbf{Q}^{\mathbf{t}}} \left(\frac{1}{\sigma} \right)^{\frac{pr}{p-1}}(x) dx \right\}^{\frac{p-1}{pr}}
\end{aligned} \tag{3.4}$$

for every $\mathbf{Q} \subset \mathbb{R}^N$.

Given \mathbf{t} , consider

$$\mathbf{Q} \subset \mathbb{R}^N : |\mathbf{Q}_i|^{\frac{1}{N_i}} / |\mathbf{Q}_v|^{\frac{1}{N_v}} = 2^{-t_i}, \quad i = 1, 2, \dots, n. \tag{3.5}$$

For $r > 1$, we define

$$\mathbf{A}_{pqr}^{\alpha}(\mathbf{t} : \omega, \sigma) = \sup_{\mathbf{Q} \text{ in (3.5)}} \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{N_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \omega^{qr}(x) dx \right\}^{\frac{1}{qr}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{\sigma} \right)^{\frac{pr}{p-1}}(x) dx \right\}^{\frac{p-1}{pr}}. \tag{3.6}$$

Suppose that \mathbf{Q} is a cube satisfying

$$|\mathbf{Q}_1|^{\frac{1}{N_1}} = |\mathbf{Q}_2|^{\frac{1}{N_2}} = \dots = |\mathbf{Q}_n|^{\frac{1}{N_n}}. \tag{3.7}$$

We find

$$\begin{aligned}
& |\mathbf{Q}|^{\frac{\alpha}{N} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \omega^{qr}(\mathbf{t}x) dx \right\}^{\frac{1}{qr}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{\sigma} \right)^{\frac{pr}{p-1}}(\mathbf{t}x) dx \right\}^{\frac{p-1}{pr}} \\
&= \prod_{i=1}^n 2^{t_i [\alpha_i - \frac{N_i}{p} + \frac{N_i}{q}]} \prod_{i=1}^n |\mathbf{Q}_i^{\mathbf{t}}|^{\frac{\alpha_i}{N_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}^{\mathbf{t}}|} \int_{\mathbf{Q}^{\mathbf{t}}} \omega^{qr}(x) dx \right\}^{\frac{1}{qr}} \left\{ \frac{1}{|\mathbf{Q}^{\mathbf{t}}|} \int_{\mathbf{Q}^{\mathbf{t}}} \left(\frac{1}{\sigma} \right)^{\frac{pr}{p-1}}(x) dx \right\}^{\frac{p-1}{pr}} \quad \text{by (3.4)} \\
&\leq \prod_{i=1}^n 2^{t_i [\alpha_i - \frac{N_i}{p} + \frac{N_i}{q}]} \mathbf{A}_{pqr}^{\alpha}(\mathbf{t} : \omega, \sigma) \quad \text{by (3.5)-(3.6)}.
\end{aligned} \tag{3.8}$$

Now, we recall Sawyer-Wheeden theorem for one-parameter fractional integral operators, stated as Theorem 1 in [6]:

$$\left\{ \int_{\mathbb{R}^N} \left\{ \int_{\mathbb{R}^N} f(y) \left[\frac{1}{|x-y|} \right]^{N-\alpha} dy \right\}^q \omega^q(x) dx \right\}^{\frac{1}{q}} \leq \mathfrak{B}_{pqr\alpha} A_{pqr}^{\alpha}(\omega, \sigma) \left\{ \int_{\mathbb{R}^N} (f\sigma)^p(x) dx \right\}^{\frac{1}{p}} \tag{3.9}$$

for $1 < p \leq q < \infty$, if

$$A_{pqr}^\alpha(\omega, \sigma) = \sup_{\mathbf{Q} : |\mathbf{Q}_1|^{\frac{1}{N_1}} = \dots = |\mathbf{Q}_n|^{\frac{1}{N_n}}} |\mathbf{Q}|^{\frac{\alpha}{N} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \omega^{qr}(x) dx \right\}^{\frac{1}{qr}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{\sigma} \right)^{\frac{pr}{p-1}} (x) dx \right\}^{\frac{p-1}{pr}} < \infty \quad (3.10)$$

for some $r > 1$.

Remark 3.1 The constant $\mathfrak{B}_{pqr\alpha} A_{pqr}^\alpha(\omega, \sigma)$ in (3.9) is not written explicitly in the statement of Theorem 1 by Sawyer and Wheeden [6]. But it can be computed directly by carrying out the proof given in section 2 of [6].

By applying (3.9)-(3.10) and using the estimate in (3.8), we have

$$\begin{aligned} & \left\{ \int_{\mathbb{R}^N} \left\{ \int_{\mathbb{R}^N} f(\mathbf{t}y) \left[\frac{1}{|x-y|} \right]^{N-\alpha} dy \right\}^q \omega^q(\mathbf{t}x) dx \right\}^{\frac{1}{q}} \\ & \leq \mathfrak{B}_{pqr\alpha} \prod_{i=1}^n 2^{t_i[\alpha_i - \frac{N_i}{p} + \frac{N_i}{q}]} \mathbf{A}_{pqr}^\alpha(\mathbf{t} : \omega, \sigma) \left\{ \int_{\mathbb{R}^N} (f\sigma)^p(\mathbf{t}x) dx \right\}^{\frac{1}{p}}. \end{aligned} \quad (3.11)$$

Recall from (3.1)-(3.2). By changing dilations $x \rightarrow \mathbf{t}x, y \rightarrow \mathbf{t}y$, we have

$$\begin{aligned} & \left\{ \int_{\mathbb{R}^N} (\Delta_{\mathbf{t}} \mathbf{I}_\alpha f)^q(x) \omega^q(x) dx \right\}^{\frac{1}{q}} \\ & = \left\{ \int_{\mathbb{R}^N} \left\{ \int_{\Gamma_{\mathbf{t}}(x)} f(y) \prod_{i=1}^n \left[\frac{1}{|x_i - y_i|} \right]^{N_i - \alpha_i} dy \right\}^q \omega^q(x) dx \right\}^{\frac{1}{q}} \\ & = \left\{ \int_{\mathbb{R}^N} \left\{ \int_{\Gamma_o(x)} f(\mathbf{t}y) \left\{ \prod_{i=1}^n 2^{-t_i N_i} \left[\frac{1}{2^{-t_i} |x_i - y_i|} \right]^{N_i - \alpha_i} \right\} dy \right\}^q \omega^q(\mathbf{t}x) \prod_{i=1}^n 2^{-t_i N_i} dx \right\}^{\frac{1}{q}} \\ & \leq \mathfrak{B}_\alpha \prod_{i=1}^n 2^{-t_i(\alpha_i + \frac{N_i}{q})} \left\{ \int_{\mathbb{R}^N} \left\{ \int_{\mathbb{R}^N} f(\mathbf{t}y) \left[\frac{1}{|x-y|} \right]^{N-\alpha} dy \right\}^q \omega^q(\mathbf{t}x) dx \right\}^{\frac{1}{q}} \\ & \leq \mathfrak{B}_{pqr\alpha} \prod_{i=1}^n 2^{-t_i(\alpha_i + \frac{N_i}{q})} 2^{t_i[\alpha_i - \frac{N_i}{p} + \frac{N_i}{q}]} \mathbf{A}_{pqr}^\alpha(\mathbf{t} : \omega, \sigma) \left\{ \int_{\mathbb{R}^N} (f\sigma)^p(\mathbf{t}x) dx \right\}^{\frac{1}{p}} \quad \text{by (3.11)} \\ & = \mathfrak{B}_{pqr\alpha} \mathbf{A}_{pqr}^\alpha(\mathbf{t} : \omega, \sigma) \prod_{i=1}^n 2^{-t_i(\alpha_i + \frac{N_i}{q})} 2^{t_i[\alpha_i - \frac{N_i}{p} + \frac{N_i}{q}]} \left\{ \int_{\mathbb{R}^N} (f\sigma)^p(x) \prod_{i=1}^n 2^{t_i N_i} dx \right\}^{\frac{1}{p}} \\ & = \mathfrak{B}_{pqr\alpha} \mathbf{A}_{pqr}^\alpha(\mathbf{t} : \omega, \sigma) \left\{ \int_{\mathbb{R}^N} (f\sigma)^p(x) dx \right\}^{\frac{1}{p}}. \end{aligned} \quad (3.12)$$

Observe that $\Delta_t \mathbf{I}_\alpha$ is essentially an one-parameter fractional integral operator, satisfying

$$\left\| (\Delta_t \mathbf{I}_\alpha f) \omega \right\|_{\mathbf{L}^q(\mathbb{R}^N)} \leq \mathfrak{B}_{p,q,r,\alpha} \mathbf{A}_{pqr}^\alpha(\mathbf{t} : \omega, \sigma) \|f\sigma\|_{\mathbf{L}^p(\mathbb{R}^N)}, \quad 1 < p \leq q < \infty. \quad (3.13)$$

By applying Minkowski inequality, we obtain (2.7) provided that

$$\sum_{\mathbf{t}} \mathbf{A}_{pqr}^\alpha(\mathbf{t} : \omega, \sigma) < \infty. \quad (3.14)$$

4 Necessary constraints

First, as investigated by Muckenhoupt, we choose $f = \sigma^{-\frac{p}{p-1}} \chi_{\mathbf{Q}}$ in (2.7). This two-weight $\mathbf{L}^p \rightarrow \mathbf{L}^q$ -norm inequality implies

$$\mathbf{A}_{pq}^\alpha(\omega, \sigma) = \sup_{\mathbf{Q} \subset \mathbb{R}^N} \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{N_i} - (\frac{1}{p} - \frac{1}{q})} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \omega^q(x) dx \right\}^{\frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{\sigma} \right)^{\frac{p}{p-1}}(x) dx \right\}^{\frac{p-1}{p}} < \infty. \quad (4.1)$$

Let $\omega(x) = |x|^{-\gamma}$, $\sigma(x) = |x|^\delta$ for $x \neq 0$ and $\gamma, \delta \in \mathbb{R}$. We aim to show (4.1) implying (2.2)-(2.6).

For the remaining section, assume \mathbf{Q} centered on the origin of \mathbb{R}^N . Denote \mathbf{Q}^λ to be a dilated variant of \mathbf{Q} for $\lambda > 0$ such that $\mathbf{Q}^\lambda = \mathbf{Q}_1^\lambda \times \mathbf{Q}_2^\lambda \times \cdots \times \mathbf{Q}_n^\lambda$ and $|\mathbf{Q}_i^\lambda|^{\frac{1}{N_i}} = \lambda |\mathbf{Q}_i|^{\frac{1}{N_i}}$, $i = 1, 2, \dots, n$. From (4.1) we find

$$\begin{aligned} & \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{N_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q} dx \right\}^{\frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\frac{\delta p}{p-1}} dx \right\}^{\frac{p-1}{p}} \\ &= \lambda^{\gamma + \delta - \alpha + \mathbf{N}(\frac{1}{p} - \frac{1}{q})} \prod_{i=1}^n |\mathbf{Q}_i^\lambda|^{\frac{\alpha_i}{N_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}^\lambda|} \int_{\mathbf{Q}^\lambda} \left(\frac{1}{|x|} \right)^{\gamma q} dx \right\}^{\frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}^\lambda|} \int_{\mathbf{Q}^\lambda} \left(\frac{1}{|x|} \right)^{\frac{\delta p}{p-1}} dx \right\}^{\frac{p-1}{p}} \\ &\leq \lambda^{\gamma + \delta - \alpha + \mathbf{N}(\frac{1}{p} - \frac{1}{q})} \mathbf{A}_{pq}^\alpha(|x|^{-\gamma}, |x|^\delta). \end{aligned} \quad (4.2)$$

Consider $|\mathbf{Q}_1|^{\frac{1}{N_1}} = |\mathbf{Q}_2|^{\frac{1}{N_2}} = \cdots = |\mathbf{Q}_n|^{\frac{1}{N_n}} = 1$. The first line of (4.2) is bounded from below. Suppose $\gamma + \delta - \alpha + \mathbf{N}(\frac{1}{p} - \frac{1}{q}) \neq 0$. By either taking $\lambda \rightarrow 0$ or $\lambda \rightarrow \infty$, the last line of (4.2) is vanished. We must have $\gamma + \delta - \alpha + \mathbf{N}(\frac{1}{p} - \frac{1}{q}) = 0$ which is (2.3).

Write $x = (x_i, x_i^\dagger) \in \mathbb{R}^{N_i} \times \mathbb{R}^{N-N_i}$, $i = 1, 2, \dots, n$ and $\mathbf{Q}_i^\dagger = \bigotimes_{j \neq i} \mathbf{Q}_j$. Suppose \mathbf{Q}_i centered on the origin of \mathbb{R}^{N_i} . Let \mathbf{Q}_i shrink to 0 and $|\mathbf{Q}_j|^{\frac{1}{N_j}} = 1$, $j \neq i$ in (4.1). By applying Lebesgue Differentiation Theorem, we have

$$\begin{aligned} & \lim_{|\mathbf{Q}_i| \rightarrow 0} |\mathbf{Q}_i|^{\frac{\alpha_i}{N_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}_i^\dagger|} \int_{\mathbf{Q}_i^\dagger} \left(\frac{1}{|x_i^\dagger|} \right)^{\gamma q} dx_i^\dagger \right\}^{\frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}_i^\dagger|} \int_{\mathbf{Q}_i^\dagger} \left(\frac{1}{|x_i^\dagger|} \right)^{\frac{\delta p}{p-1}} dx_i^\dagger \right\}^{\frac{p-1}{p}} \\ &\leq \mathbf{A}_{pq}^\alpha(|x|^{-\gamma}, |x|^\delta), \quad i = 1, 2, \dots, n. \end{aligned} \quad (4.3)$$

The boundedness of $\mathbf{A}_{pq}^\alpha(|x|^{-\gamma}, |x|^\delta)$ requires

$$\frac{\alpha_i}{\mathbf{N}_i} \geq \frac{1}{p} - \frac{1}{q}, \quad i = 1, 2, \dots, n. \quad (4.4)$$

By putting together (2.3) and (4.4), we find $\gamma + \delta \geq 0$. On the other and, it is essential to require $\gamma q < \mathbf{N}$ and $\delta(\frac{p}{p-1}) < \mathbf{N}$ for the local integrability of $|x|^{-\gamma q}$ and $|x|^{-\delta(\frac{p}{p-1})}$ respectively. These are the constraints in (2.2).

Let \mathcal{S} to be a proper subset of $\{1, 2, \dots, n\}$. We define ${}^t\mathbf{Q}_i = \mathbf{Q}_i \cap \{2^{-t_i} \leq |x_i| \leq 2^{-t_i+1}\}$ for every $t_i > 0$ and $i \in \mathcal{S}$. Denote ${}^t\mathbf{Q} = \bigotimes_{i \in \mathcal{S}} {}^t\mathbf{Q}_i \times \bigotimes_{i \in \mathcal{S}^c} \mathbf{Q}_i$ and $\mathbf{Q}_{\mathcal{S}} = \bigotimes_{i \in \mathcal{S}} \mathbf{Q}_i$, $\mathbf{Q}_{\mathcal{S}^c} = \bigotimes_{i \in \mathcal{S}^c} \mathbf{Q}_i$. Moreover, we write $x = (x_{\mathcal{S}}, x_{\mathcal{S}^c}) \in \mathbb{R}^{\mathbf{N}_{\mathcal{S}}} \times \mathbb{R}^{\mathbf{N}-\mathbf{N}_{\mathcal{S}}}$ for which $\mathbf{N}_{\mathcal{S}} = \sum_{i \in \mathcal{S}} \mathbf{N}_i$.

Suppose that there exists at least one $i \in \mathcal{S}^c$ such that $\alpha_i - \mathbf{N}_i(\frac{1}{p} - \frac{1}{q}) > 0$. Let $0 < \lambda < 1$.

Consider $|\mathbf{Q}_i|^{\frac{1}{\mathbf{N}_i}} = 1$ for $i \in \mathcal{S}$ and $|\mathbf{Q}_i|^{\frac{1}{\mathbf{N}_i}} = \lambda$ for $i \in \mathcal{S}^c$. We have

$$\begin{aligned} & \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{\mathbf{N}_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q} dx \right\}^{\frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\frac{\delta p}{p-1}} dx \right\}^{\frac{p-1}{p}} \\ &= \lambda^{\sum_{i \in \mathcal{S}^c} \alpha_i - \mathbf{N}_i(\frac{1}{p} - \frac{1}{q})} \left\{ \sum_{\mathbf{t}} \frac{1}{|\mathbf{Q}|} \int_{{}^t\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q} dx \right\}^{\frac{1}{q}} \left\{ \sum_{\mathbf{t}} \frac{1}{|\mathbf{Q}|} \int_{{}^t\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\frac{\delta p}{p-1}} dx \right\}^{\frac{p-1}{p}} \\ &= \left\{ \lambda^{[\sum_{i \in \mathcal{S}^c} \alpha_i - \mathbf{N}_i(\frac{1}{p} - \frac{1}{q})] \frac{q}{2}} \sum_{\mathbf{t}} \frac{1}{|\mathbf{Q}|} \int_{{}^t\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q} dx \right\}^{\frac{1}{q}} \\ & \quad \left\{ \lambda^{[\sum_{i \in \mathcal{S}^c} \alpha_i - \mathbf{N}_i(\frac{1}{p} - \frac{1}{q})] (\frac{p}{p-1}) \frac{1}{2}} \sum_{\mathbf{t}} \frac{1}{|\mathbf{Q}|} \int_{{}^t\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\frac{\delta p}{p-1}} dx \right\}^{\frac{p-1}{p}} \\ &\doteq \left\{ \sum_{\mathbf{t}} \mathbf{A}_{\mathbf{t}}(\lambda) \right\}^{\frac{1}{q}} \left\{ \sum_{\mathbf{t}} \mathbf{B}_{\mathbf{t}}(\lambda) \right\}^{\frac{p-1}{p}}. \end{aligned} \quad (4.5)$$

By applying Lebesgue differentiation theorem, we find

$$\begin{aligned} \lim_{\lambda \rightarrow 0} \frac{1}{|\mathbf{Q}|} \int_{{}^t\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q} dx &= \int \cdots \int_{\bigotimes_{i \in \mathcal{S}} {}^t\mathbf{Q}_i} \left[\frac{1}{\sum_{i \in \mathcal{S}} |x_i|^2} \right]^{\frac{\gamma q}{2}} \prod_{i \in \mathcal{S}} dx_i, \\ \lim_{\lambda \rightarrow 0} \frac{1}{|\mathbf{Q}|} \int_{{}^t\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\frac{\delta p}{p-1}} dx &= \int \cdots \int_{\bigotimes_{i \in \mathcal{S}} {}^t\mathbf{Q}_i} \left[\frac{1}{\sum_{i \in \mathcal{S}} |x_i|^2} \right]^{\frac{1}{2} \frac{\delta p}{p-1}} \prod_{i \in \mathcal{S}} dx_i. \end{aligned} \quad (4.6)$$

Because $\sum_{i \in \mathcal{S}^c} \alpha_i - \mathbf{N}_i(\frac{1}{p} - \frac{1}{q}) > 0$, we have $\mathbf{A}_{\mathbf{t}}(0) = \mathbf{B}_{\mathbf{t}}(0) = 0$ for every \mathbf{t} . This remains to be true if $\sum_{i \in \mathcal{S}^c} \alpha_i - \mathbf{N}_i(\frac{1}{p} - \frac{1}{q})$ is replaced by any smaller positive number. Therefore, each $\mathbf{A}_{\mathbf{t}}(\lambda)$ and $\mathbf{B}_{\mathbf{t}}(\lambda)$ is Hölder continuous for $\lambda \geq 0$ whose exponent is strictly positive. Furthermore, $\sum_{\mathbf{t}} \mathbf{A}_{\mathbf{t}}(\lambda)$ and $\sum_{\mathbf{t}} \mathbf{B}_{\mathbf{t}}(\lambda)$ are uniformly bounded for every $\lambda > 0$. Consequently, both summations are continuous at $\lambda = 0$ and

$$\lim_{\lambda \rightarrow 0} \sum_{\mathbf{t}} \mathbf{A}_{\mathbf{t}}(\lambda) = 0, \quad \lim_{\lambda \rightarrow 0} \sum_{\mathbf{t}} \mathbf{B}_{\mathbf{t}}(\lambda) = 0. \quad (4.7)$$

Suppose $\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right) = 0$ for every $i \in \mathcal{S}^c$. Shrink \mathbf{Q}_i to the origin of $\mathbb{R}^{\mathbf{N}_i}$ for $i \in \mathcal{S}^c$ in (4. 1). By applying Lebesgue differentiation theorem, we have

$$\mathbf{A}_{pq}^\alpha \left(|x|^{-\gamma}, |x|^\delta \right) \geq \prod_{i \in \mathcal{S}} |\mathbf{Q}_i|^{\frac{\alpha_i}{\mathbf{N}_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}_S|} \int_{\mathbf{Q}_S} \left(\frac{1}{|x_S|} \right)^{\gamma q} dx_S \right\}^{\frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}_S|} \int_{\mathbf{Q}_S} \left(\frac{1}{|x_S|} \right)^{\frac{\delta p}{p-1}} dx_S \right\}^{\frac{p-1}{p}} \quad (4. 8)$$

where $\gamma q < \mathbf{N}_S$ and $\delta \left(\frac{p}{p-1} \right) < \mathbf{N}_S$ become necessities.

Case One: Consider $\gamma \geq 0, \delta \leq 0$. Let $|\mathbf{Q}_i|^{\frac{1}{\mathbf{N}_i}} = 1$ for $i \in \{1, 2, \dots, n\}$ and $|\mathbf{Q}_j|^{\frac{1}{\mathbf{N}_j}} = \lambda$ for all $j \neq i$. Suppose $\alpha_j - \mathbf{N}_j \left(\frac{1}{p} - \frac{1}{q} \right) = 0$ for every $j \neq i$. We have

$$\begin{aligned} & \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{\mathbf{N}_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q} dx \right\}^{\frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\frac{\delta p}{p-1}} dx \right\}^{\frac{p-1}{p}} \\ & \geq \mathfrak{B}_{q \gamma} \left\{ \int_{\mathbf{Q}_i} \left(\frac{1}{\lambda + |x_i|} \right)^{\gamma q} dx_i \right\}^{\frac{1}{q}} \left\{ \int_{\mathbf{Q}_i} \left(\frac{1}{|x_i|} \right)^{\frac{\delta p}{p-1}} dx_i \right\}^{\frac{p-1}{p}} \quad (\delta \leq 0) \\ & \geq \mathfrak{B}_{p \gamma} \left\{ \int_{\lambda < |x_i| \leq 1} \left(\frac{1}{\lambda + |x_i|} \right)^{\gamma q} dx_i \right\}^{\frac{1}{q}} \end{aligned} \quad (4. 9)$$

where

$$\begin{aligned} & \int_{\lambda < |x_i| \leq 1} \left(\frac{1}{\lambda + |x_i|} \right)^{\gamma q} dx_i = \mathfrak{B} \ln \left(\frac{1 + \lambda}{2\lambda} \right) \quad \text{if } \gamma = \frac{\mathbf{N}_i}{q}, \\ & \int_{\lambda < |x_i| \leq 1} \left(\frac{1}{\lambda + |x_i|} \right)^{\gamma q} dx_i = \mathfrak{B} \frac{1}{\gamma q - \mathbf{N}_i} \left[\left(\frac{1}{2\lambda} \right)^{\gamma q - \mathbf{N}_i} - \left(\frac{1}{\lambda + 1} \right)^{\gamma q - \mathbf{N}_i} \right] \quad \text{if } \gamma > \frac{\mathbf{N}_i}{q}. \end{aligned} \quad (4. 10)$$

From (4. 9)-(4. 10), to satisfy the inequality in (4. 2) as $\lambda \rightarrow 0$, we need

$$\gamma < \frac{\mathbf{N}_i}{q}, \quad i = 1, 2, \dots, n. \quad (4. 11)$$

Suppose that there exists $j \neq i$ such that $\alpha_j - \mathbf{N}_j \left(\frac{1}{p} - \frac{1}{q} \right) > 0$. We have

$$\begin{aligned} & \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{\mathbf{N}_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q} dx \right\}^{\frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\frac{\delta p}{p-1}} dx \right\}^{\frac{p-1}{p}} \\ & \geq \mathfrak{B}_{q \gamma} \prod_{j \neq i} \lambda^{\alpha_j - \mathbf{N}_j \left(\frac{1}{p} - \frac{1}{q} \right)} \left\{ \int_{\mathbf{Q}_i} \left(\frac{1}{\lambda + |x_i|} \right)^{\gamma q} dx_i \right\}^{\frac{1}{q}} \left\{ \int_{\mathbf{Q}_i} \left(\frac{1}{|x_i|} \right)^{\frac{\delta p}{p-1}} dx_i \right\}^{\frac{p-1}{p}} \quad (\delta \leq 0) \\ & \geq \mathfrak{B}_{p \gamma} \prod_{j \neq i} \lambda^{\alpha_j - \mathbf{N}_j \left(\frac{1}{p} - \frac{1}{q} \right)} \left\{ \int_{0 < |x_i| \leq \lambda} \left(\frac{1}{\lambda} \right)^{\gamma q} dx_i \right\}^{\frac{1}{q}} = \mathfrak{B}_{p \gamma} \lambda^{\frac{\mathbf{N}_i}{q} - \gamma + \sum_{j \neq i} \alpha_j - \mathbf{N}_j \left(\frac{1}{p} - \frac{1}{q} \right)}. \end{aligned} \quad (4. 12)$$

Recall the estimates in (4. 5)-(4. 7). Take $\mathcal{S} = \{i\}$. The first line of (4. 12) equals to zero as $\lambda \rightarrow 0$. Together with (4. 11), we find

$$\gamma < \frac{\mathbf{N}_i}{q} + \sum_{j \neq i} \alpha_j - \mathbf{N}_j \left(\frac{1}{p} - \frac{1}{q} \right), \quad i = 1, 2, \dots, n. \quad (4. 13)$$

Case Two: Consider $\gamma \leq 0, \delta \geq 0$. Let $|\mathbf{Q}_i|^{\frac{1}{\mathbf{N}_i}} = 1$ for $i \in \{1, 2, \dots, n\}$ and $|\mathbf{Q}_j|^{\frac{1}{\mathbf{N}_j}} = \lambda$ for all $j \neq i$. Suppose $\alpha_j - \mathbf{N}_j \left(\frac{1}{p} - \frac{1}{q} \right) = 0$ for every $j \neq i$. We have

$$\begin{aligned} & \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{\mathbf{N}_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q} dx \right\}^{\frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\frac{\delta p}{p-1}} dx \right\}^{\frac{p-1}{p}} \\ & \geq \mathfrak{B}_p \delta \left\{ \int_{\mathbf{Q}_i} \left(\frac{1}{|x_i|} \right)^{\gamma q} dx_i \right\}^{\frac{1}{q}} \left\{ \int_{\mathbf{Q}_i} \left(\frac{1}{\lambda + |x_i|} \right)^{\frac{\delta p}{p-1}} dx_i \right\}^{\frac{p-1}{p}} \quad (\gamma \leq 0) \\ & \geq \mathfrak{B}_{p \ q \ \gamma \ \delta} \left\{ \int_{\lambda < |x_i| \leq 1} \left(\frac{1}{\lambda + |x_i|} \right)^{\frac{\delta p}{p-1}} dx_i \right\}^{\frac{p-1}{p}} \end{aligned} \quad (4. 14)$$

where

$$\begin{aligned} & \int_{\lambda < |x_i| \leq 1} \left(\frac{1}{\lambda + |x_i|} \right)^{\delta \left(\frac{p}{p-1} \right)} dx_i = \mathfrak{B}_{\mathbf{N}} \ln \left(\frac{1 + \lambda}{2\lambda} \right) \quad \text{if } \delta = \mathbf{N}_i \left(\frac{p-1}{p} \right), \\ & \int_{\lambda < |x_i| \leq 1} \left(\frac{1}{\lambda + |x_i|} \right)^{\delta \left(\frac{p}{p-1} \right)} dx_i = \mathfrak{B}_{\mathbf{N}} \frac{1}{\delta \left(\frac{p}{p-1} \right) - \mathbf{N}_i} \left[\left(\frac{1}{2\lambda} \right)^{\delta \left(\frac{p}{p-1} \right) - \mathbf{N}_i} - \left(\frac{1}{\lambda + 1} \right)^{\delta \left(\frac{p}{p-1} \right) - \mathbf{N}_i} \right] \\ & \quad \text{if } \delta > \mathbf{N}_i \left(\frac{p-1}{p} \right). \end{aligned} \quad (4. 15)$$

From (4. 14)-(4. 15), to satisfy the inequality in (4. 2) as $\lambda \rightarrow 0$, we need

$$\delta < \mathbf{N}_i \left(\frac{p-1}{p} \right), \quad i = 1, 2, \dots, n. \quad (4. 16)$$

Suppose that there exists $j \neq i$ such that $\alpha_j - \mathbf{N}_j \left(\frac{1}{p} - \frac{1}{q} \right) > 0$. We have

$$\begin{aligned} & \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{\mathbf{N}_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q} dx \right\}^{\frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\frac{\delta p}{p-1}} dx \right\}^{\frac{p-1}{p}} \\ & \geq \mathfrak{B}_p \delta \prod_{j \neq i} \lambda^{\alpha_j - \mathbf{N}_j \left(\frac{1}{p} - \frac{1}{q} \right)} \left\{ \int_{\mathbf{Q}_i} \left(\frac{1}{|x_i|} \right)^{\gamma q} dx_i \right\}^{\frac{1}{q}} \left\{ \int_{\mathbf{Q}_i} \left(\frac{1}{\lambda + |x_i|} \right)^{\frac{\delta p}{p-1}} dx_i \right\}^{\frac{p-1}{p}} \quad (\gamma \leq 0) \\ & \geq \mathfrak{B}_{p \ q \ \gamma \ \delta} \prod_{j \neq i} \lambda^{\alpha_j - \mathbf{N}_j \left(\frac{1}{p} - \frac{1}{q} \right)} \left\{ \int_{0 < |x_i| \leq \lambda} \left(\frac{1}{\lambda} \right)^{\frac{\delta p}{p-1}} dx_i \right\}^{\frac{p-1}{p}} \\ & = \mathfrak{B}_{p \ q \ \gamma \ \delta} \lambda^{\left(\frac{p-1}{p} \right) \mathbf{N}_i - \delta + \sum_{j \neq i} \alpha_j - \mathbf{N}_j \left(\frac{1}{p} - \frac{1}{q} \right)}. \end{aligned} \quad (4. 17)$$

Recall the estimates in (4. 5)-(4. 7). Take $\mathcal{S} = \{i\}$. The first line of (4. 17) equals to zero as $\lambda \rightarrow 0$. Together with (4. 16), we find

$$\delta < \mathbf{N}_i \left(\frac{p-1}{p} \right) + \sum_{j \neq i} \alpha_j - \mathbf{N}_j \left(\frac{1}{p} - \frac{1}{q} \right), \quad i = 1, 2, \dots, n. \quad (4. 18)$$

Case Three: Consider $\gamma > 0, \delta > 0$. Note that (4. 1) is invariant by changing dilation in one-parameter as shown in (4. 2) because of (2. 3).

Recall the definition of \mathcal{U} and \mathcal{V} from (2. 6). We write $x_{\mathcal{U}} \in \mathbb{R}^{\mathbf{N}_{\mathcal{U}}}$ and $x_{\mathcal{V}} \in \mathbb{R}^{\mathbf{N}_{\mathcal{V}}}$ where $\mathbb{R}^{\mathbf{N}_{\mathcal{U}}} = \bigotimes_{i \in \mathcal{U}} \mathbb{R}^{\mathbf{N}_i}$ and $\mathbb{R}^{\mathbf{N}_{\mathcal{V}}} = \bigotimes_{i \in \mathcal{V}} \mathbb{R}^{\mathbf{N}_i}$.

Let $|\mathbf{Q}_i|^{\frac{1}{\mathbf{N}_i}} = \lambda^{-1}$ for every $i \in \mathcal{U}$ and $|\mathbf{Q}_i|^{\frac{1}{\mathbf{N}_i}} = 1$ for all other $i \notin \mathcal{U}$. We have

$$\begin{aligned} & \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{\mathbf{N}_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q} dx \right\}^{\frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\frac{\delta p}{p-1}} dx \right\}^{\frac{p-1}{p}} \\ & \geq \mathfrak{B}_{p \ q \ \gamma \ \delta} \prod_{i \in \mathcal{U}} \left(\frac{1}{\lambda} \right)^{\alpha_i - \frac{\mathbf{N}_i}{p}} \left\{ \int \cdots \int_{\bigotimes_{i \in \mathcal{U}} \mathbf{Q}_i} \left[\frac{1}{1 + \sum_{i \in \mathcal{U}} |x_i|} \right]^{\gamma q} \prod_{i \in \mathcal{U}} dx_i \right\}^{\frac{1}{q}} \\ & \quad \left\{ \prod_{i \in \mathcal{U}} \lambda^{\mathbf{N}_i} \int \cdots \int_{\bigotimes_{i \in \mathcal{U}} \mathbf{Q}_i} \lambda^{\frac{\delta p}{p-1}} \prod_{i \in \mathcal{U}} dx_i \right\}^{\frac{p-1}{p}} \quad (0 < \lambda < 1) \\ & \geq \mathfrak{B}_{p \ q \ \gamma \ \delta} \prod_{i \in \mathcal{U}} \left(\frac{1}{\lambda} \right)^{\alpha_i - \frac{\mathbf{N}_i}{p}} \left\{ \int \cdots \int_{\bigotimes_{i \in \mathcal{U}} 0 < |x_i| \leq 1} \prod_{i \in \mathcal{U}} dx_i \right\}^{\frac{1}{q}} \\ & \quad \left\{ \prod_{i \in \mathcal{U}} \lambda^{\mathbf{N}_i} \int \cdots \int_{\bigotimes_{i \in \mathcal{U}} \mathbf{Q}_i} \lambda^{\frac{\delta p}{p-1}} \prod_{i \in \mathcal{U}} dx_i \right\}^{\frac{p-1}{p}} \\ & \geq \mathfrak{B}_{p \ q \ \gamma \ \delta} \left(\frac{1}{\lambda} \right)^{\sum_{i \in \mathcal{U}} \alpha_i - \frac{\mathbf{N}_i}{p} - \delta}. \end{aligned} \quad (4. 19)$$

In the case of $\mathcal{U} = \{1, 2, \dots, n\}$, as $\gamma < \frac{\mathbf{N}}{q}$, we find

$$\begin{aligned} \delta &= \frac{\mathbf{N}}{q} - \gamma + \sum_{i=1}^n \alpha_i - \frac{\mathbf{N}_i}{p} \quad \text{by (2. 3)} \\ &> \sum_{i=1}^n \alpha_i - \frac{\mathbf{N}_i}{p} = \sum_{i \in \mathcal{U}} \alpha_i - \frac{\mathbf{N}_i}{p}. \end{aligned} \quad (4. 20)$$

Suppose that \mathcal{U} is a proper subset of $\{1, 2, \dots, n\}$. There exists at least one $i \in \mathcal{U}^c$ such that $\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right) > 0$. Recall the estimates in (4. 5)-(4. 7). Take $\mathcal{S} = \mathcal{U}$. The first line of (4. 19) equals to zero as $\lambda \rightarrow 0$. We must have

$$\sum_{i \in \mathcal{U}} \alpha_i - \frac{\mathbf{N}_i}{p} < \delta. \quad (4. 21)$$

Suppose that \mathcal{U} is a proper subset of $\{1, 2, \dots, n\}$ where $\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right) = 0$ for every $i \in \mathcal{U}^c$.

Let $\mathcal{S} = \mathcal{U}$. We have $|x_{\mathcal{U}}|^{-\gamma}$ and $|x_{\mathcal{U}}|^{\delta}$ satisfying the Muckenhoupt characteristic in (4. 8) on $\mathbb{R}^{\mathbf{N}_{\mathcal{U}}} \doteq \bigotimes_{i \in \mathcal{U}} \mathbb{R}^{\mathbf{N}_i}$. Denote

$$\alpha_{\mathcal{U}} = \sum_{i \in \mathcal{U}} \alpha_i.$$

By carrying out the same estimate in (4. 2), we find

$$\gamma < \frac{\mathbf{N}_{\mathcal{U}}}{q}, \quad \delta < \mathbf{N}_{\mathcal{U}} \left(\frac{p-1}{p} \right), \quad \frac{\alpha_{\mathcal{U}}}{\mathbf{N}_{\mathcal{U}}} = \frac{1}{p} - \frac{1}{q} + \frac{\gamma + \delta}{\mathbf{N}_{\mathcal{U}}}. \quad (4. 22)$$

This further implies

$$\begin{aligned} \delta &= \frac{\mathbf{N}_{\mathcal{U}}}{q} - \gamma + \sum_{i \in \mathcal{U}} \alpha_i - \frac{\mathbf{N}_i}{p} \\ &> \sum_{i \in \mathcal{U}} \alpha_i - \frac{\mathbf{N}_i}{p}. \end{aligned} \quad (4. 23)$$

Let $|\mathbf{Q}_i|^{\frac{1}{\mathbf{N}_i}} = \lambda^{-1}$ for every $i \in \mathcal{V}$ and $|\mathbf{Q}_i|^{\frac{1}{\mathbf{N}_i}} = 1$ for all other $i \notin \mathcal{V}$. We have

$$\begin{aligned} &\prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{\mathbf{N}_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q} dx \right\}^{\frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\frac{\delta p}{p-1}} dx \right\}^{\frac{p-1}{p}} \\ &\geq \mathfrak{B}_{p \ q \ \gamma \ \delta} \left(\frac{1}{\lambda} \right)^{\sum_{i \in \mathcal{V}} \alpha_i - \left(\frac{q-1}{q} \right) \mathbf{N}_i} \left\{ \prod_{i \in \mathcal{V}} \lambda^{\mathbf{N}_i} \int \cdots \int_{\bigotimes_{i \in \mathcal{V}} \mathbf{Q}_i} \lambda^{\gamma q} \prod_{i \in \mathcal{V}} dx_i \right\}^{\frac{1}{q}} \\ &\quad \left\{ \int \cdots \int_{\bigotimes_{i \in \mathcal{V}} \mathbf{Q}_i} \left[\frac{1}{1 + \sum_{i \in \mathcal{V}} |x_i|} \right]^{\frac{\delta p}{p-1}} \prod_{i \in \mathcal{V}} dx_i \right\}^{\frac{p-1}{p}} \\ &\geq \mathfrak{B}_{p \ q \ \gamma \ \delta} \left(\frac{1}{\lambda} \right)^{\sum_{i \in \mathcal{V}} \alpha_i - \left(\frac{q-1}{q} \right) \mathbf{N}_i} \left\{ \prod_{i \in \mathcal{V}} \lambda^{\mathbf{N}_i} \int \cdots \int_{\bigotimes_{i \in \mathcal{V}} \mathbf{Q}_i} \lambda^{\gamma q} \prod_{i \in \mathcal{V}} dx_i \right\}^{\frac{1}{q}} \\ &\quad \left\{ \int \cdots \int_{\bigotimes_{i \in \mathcal{V}} 0 < |x_i| \leq 1} \prod_{i \in \mathcal{V}} dx_i \right\}^{\frac{p-1}{p}} \\ &\geq \mathfrak{B}_{p \ q \ \gamma \ \delta} \left(\frac{1}{\lambda} \right)^{\sum_{i \in \mathcal{V}} \alpha_i - \left(\frac{q-1}{q} \right) \mathbf{N}_i - \gamma}. \end{aligned} \quad (4. 24)$$

In the case of $\mathcal{V} = \{1, 2, \dots, n\}$, as $\delta < \mathbf{N} \left(\frac{p-1}{p} \right)$, we find

$$\begin{aligned} \gamma &= \left(\frac{p-1}{p} \right) \mathbf{N} - \delta + \sum_{i=1}^n \alpha_i - \mathbf{N}_i \left(\frac{q-1}{q} \right) \quad \text{by (2. 3)} \\ &> \sum_{i=1}^n \alpha_i - \mathbf{N}_i \left(\frac{q-1}{q} \right) = \sum_{i \in \mathcal{V}} \alpha_i - \mathbf{N}_i \left(\frac{q-1}{q} \right). \end{aligned} \quad (4. 25)$$

Suppose that \mathcal{V} is a proper subset of $\{1, 2, \dots, n\}$. There exists at least one $i \in \mathcal{V}^c$ such that $\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right) > 0$. Recall the estimates in (4. 5)-(4. 7). Take $\mathcal{S} = \mathcal{V}$. The first line of (4. 24) equals to zero at $\lambda \rightarrow 0$. We must have

$$\sum_{i \in \mathcal{V}} \alpha_i - \left(\frac{q-1}{q} \right) \mathbf{N}_i < \gamma. \quad (4. 26)$$

Suppose that \mathcal{V} is a proper subset of $\{1, 2, \dots, n\}$ where $\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right) = 0$ for every $i \in \mathcal{V}^c$. Let $\mathcal{S} = \mathcal{V}$. We have $|x_{\mathcal{V}}|^{-\gamma}$ and $|x_{\mathcal{V}}|^{\delta}$ satisfying the Muckenhoupt characteristic in (4. 8) on $\mathbb{R}^{\mathbf{N}_{\mathcal{V}}} \doteq \bigotimes_{i \in \mathcal{V}} \mathbb{R}^{\mathbf{N}_i}$. Denote

$$\alpha_{\mathcal{V}} = \sum_{i \in \mathcal{V}} \alpha_i.$$

By carrying out the same estimate in (4. 2), we find

$$\gamma < \frac{\mathbf{N}_{\mathcal{V}}}{q}, \quad \delta < \mathbf{N}_{\mathcal{V}} \left(\frac{p-1}{p} \right), \quad \frac{\alpha_{\mathcal{V}}}{\mathbf{N}_{\mathcal{V}}} = \frac{1}{p} - \frac{1}{q} + \frac{\gamma + \delta}{\mathbf{N}_{\mathcal{V}}}. \quad (4. 27)$$

This further implies

$$\begin{aligned} \gamma &= \left(\frac{p-1}{p} \right) \mathbf{N}_{\mathcal{V}} - \delta + \sum_{i \in \mathcal{V}} \alpha_i - \mathbf{N}_i \left(\frac{q-1}{q} \right) \\ &> \sum_{i \in \mathcal{V}} \alpha_i - \mathbf{N}_i \left(\frac{q-1}{q} \right). \end{aligned} \quad (4. 28)$$

Remark 4.1 By using the formula in (2. 3), we can verify that the constraints in (4. 13) and (4. 18) are equivalent to (2. 4) and (2. 5) respectively: For $\gamma \geq 0, \delta \leq 0$, we have

$$\gamma < \frac{\mathbf{N}_i}{q} + \sum_{j \neq i} \alpha_j - \mathbf{N}_j \left(\frac{1}{p} - \frac{1}{q} \right) \iff \alpha_i - \frac{\mathbf{N}_i}{p} < \delta, \quad i = 1, 2, \dots, n. \quad (4. 29)$$

For $\gamma \leq 0, \delta \geq 0$, we find

$$\begin{aligned} \delta &< \mathbf{N}_i \left(\frac{p-1}{p} \right) + \sum_{j \neq i} \alpha_j - \mathbf{N}_j \left(\frac{1}{p} - \frac{1}{q} \right) \iff \\ \alpha_i - \mathbf{N}_i \left(\frac{q-1}{q} \right) &< \gamma, \quad i = 1, 2, \dots, n. \end{aligned} \quad (4. 30)$$

5 Decay estimate on varying eccentricities

Recall $\mathbf{A}_{pqr}^{\alpha}(\mathbf{t} : \omega, \sigma)$ defined in (3. 6). We aim to show

$$\mathbf{A}_{pqr}^{\alpha}(\mathbf{t} : \omega, \sigma) \leq \mathfrak{B}_{pqr\alpha\gamma\delta} \prod_{i=1}^n 2^{-\varepsilon |t_i|}$$

for some $\varepsilon = \varepsilon(p, q, r, \alpha, \gamma, \delta) > 0$. This further implies the summability in (3. 14).

Principal Lemma: Let γ, δ satisfying (2. 2)-(2. 6). Suppose

$$\frac{\alpha_i}{N_i} > \frac{1}{p} - \frac{1}{q}, \quad i = 1, 2, \dots, n. \quad (5. 1)$$

For $0 < \lambda_i \leq 1, i = 1, 2, \dots, n$, define

$${}^\lambda \mathbf{Q} \subset \mathbb{R}^N : |\mathbf{Q}_i|^{\frac{1}{N_i}} / |\mathbf{Q}_v|^{\frac{1}{N_v}} = \lambda_i. \quad (5. 2)$$

There exists an $\varepsilon > 0$ such that

$$\begin{aligned} \sup_{{}^\lambda \mathbf{Q}} \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{N_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q r} dx \right\}^{\frac{1}{q r}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\frac{\delta p r}{p-1}} dx \right\}^{\frac{p-1}{p r}} \\ \leq \mathfrak{B}_{p \ q \ r \ \alpha \ \gamma \ \delta} \prod_{i=1}^n (\lambda_i)^\varepsilon \end{aligned} \quad (5. 3)$$

for some $r > 1$. The values of ε and r depend only on $p, q, \alpha, \gamma, \delta$.

Proof: By carrying out the same estimate in (4. 2) and using the formula (2. 3), we find that the r -bump characteristic in (5. 3) is invariant by changing one-parameter dilation. Therefore, it is suffice to consider $|\mathbf{Q}_v|^{\frac{1}{N_v}} = 1$.

Let \mathbf{Q}_i^o and $\mathbf{Q}_i^* \subset \mathbb{R}^{N_i}$ to be centered on the origin of \mathbb{R}^{N_i} and

$$\begin{aligned} |\mathbf{Q}_i^o|^{\frac{1}{N_i}} &= |\mathbf{Q}_i|^{\frac{1}{N_i}}, \quad i = 1, 2, \dots, n; \\ |\mathbf{Q}_i^*|^{\frac{1}{N_i}} &= 3|\mathbf{Q}_i|^{\frac{1}{N_i}} = 3\lambda_i, \quad i = 1, 2, \dots, n. \end{aligned} \quad (5. 4)$$

Remark 5.1 Suppose $\mathbf{Q}_i \cap \mathbf{Q}_i^o = \emptyset$. We must have $|x_i| \geq |x_i^o| / \sqrt{n}$ for every $x_i \in \mathbf{Q}_i$ and every $x_i^o \in \mathbf{Q}_i^o$. Otherwise, if \mathbf{Q}_i intersects \mathbf{Q}_i^o , then $\mathbf{Q}_i \subset \mathbf{Q}_i^*$.

After a permutation on indices $i = 1, 2, \dots, n$, we can assume $v = 1$ and

$$1 = \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n. \quad (5. 5)$$

Case One: Let $\gamma \geq 0, \delta \leq 0$ satisfying (2. 2)-(2. 4). By adjusting the value of r , we assume

$$\sum_{i=1}^{m-1} N_i < \gamma q r < \sum_{i=1}^m N_i, \quad 1 \leq m \leq n. \quad (5. 6)$$

Suppose that \mathbf{Q} is centered on some $z \in \mathbb{R}^N$ with $|z| \leq 3$. We have

$$\begin{aligned}
& \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{N_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q r} dx \right\}^{\frac{1}{q r}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\delta \frac{p r}{p-1}} dx \right\}^{\frac{p-1}{p r}} \\
& \leq \mathfrak{B}_{p \ q \ r \ \gamma \ \delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})} \left\{ \prod_{i=1}^n \left(\frac{1}{\lambda_i} \right)^{N_i} \int \cdots \int_{\bigotimes_{i=1}^n \mathbf{Q}_i} \left[\frac{1}{|x_1| + \cdots + |x_n|} \right]^{\gamma q r} dx_1 \cdots dx_n \right\}^{\frac{1}{q r}} \\
& \quad (\delta \leq 0) \\
& \leq \mathfrak{B}_{p \ q \ r \ \gamma \ \delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=1}^n \left(\frac{1}{\lambda_i} \right)^{\frac{N_i}{q r}} \\
& \quad \left\{ \int \cdots \int_{\bigotimes_{i=m}^n \mathbf{Q}_i} \left\{ \int \cdots \int_{\bigotimes_{i=1}^{m-1} \mathbb{R}^{N_i}} \left[\frac{1}{|x_1| + \cdots + |x_n|} \right]^{\gamma q r} dx_1 \cdots dx_{m-1} \right\} dx_m \cdots dx_n \right\}^{\frac{1}{q r}} \\
& \leq \mathfrak{B}_{p \ q \ r \ \gamma \ \delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=1}^n \left(\frac{1}{\lambda_i} \right)^{\frac{N_i}{q r}} \left\{ \int \cdots \int_{\bigotimes_{i=m}^n \mathbf{Q}_i} \left[\frac{1}{|x_m| + \cdots + |x_n|} \right]^{\gamma q r - \sum_{i=1}^{m-1} N_i} dx_m \cdots dx_n \right\}^{\frac{1}{q r}} \\
& \leq \mathfrak{B}_{p \ q \ r \ \gamma \ \delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=1}^n \left(\frac{1}{\lambda_i} \right)^{\frac{N_i}{q r}} \left\{ \int \cdots \int_{\bigotimes_{i=m}^n \mathbf{Q}_i} \left(\frac{1}{|x_m|} \right)^{\gamma q r - \sum_{i=1}^{m-1} N_i} dx_m \cdots dx_n \right\}^{\frac{1}{q r}} \\
& \leq \mathfrak{B}_{p \ q \ r \ \gamma \ \delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=1}^m \left(\frac{1}{\lambda_i} \right)^{\frac{N_i}{q r}} \left\{ \int_{\mathbf{Q}_m^*} \left(\frac{1}{|x_m|} \right)^{\gamma q r - \sum_{i=1}^{m-1} N_i} dx_m \right\}^{\frac{1}{q r}} \quad \text{by Remark 5.1} \\
& \leq \mathfrak{B}_{p \ q \ r \ \gamma \ \delta} (\lambda_m)^{\frac{1}{q r} \sum_{i=1}^m N_i - \gamma} \prod_{i=1}^m \left(\frac{1}{\lambda_i} \right)^{\frac{N_i}{q r}} \prod_{i=1}^n (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})}.
\end{aligned} \tag{5.7}$$

From direct computation, the formula in the last line of (5.7) can be rewritten as

$$\begin{aligned}
& (\lambda_m)^{\frac{1}{q r} \sum_{i=1}^m N_i - \gamma} \prod_{i=1}^m (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q}) - \frac{N_i}{q r}} \prod_{i=m+1}^n (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})} \\
& = (\lambda_m)^{\frac{1}{q r} \sum_{i=1}^m N_i - \gamma} \prod_{i=2}^m (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q}) - \frac{N_i}{q r}} \prod_{i=m+1}^n (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})} \quad (\lambda_1 = 1) \\
& = (\lambda_m)^{\frac{N_1}{q r} + \sum_{i=2}^m \alpha_i - N_i (\frac{1}{p} - \frac{1}{q}) - \gamma} \prod_{i=2}^m \left(\frac{\lambda_i}{\lambda_m} \right)^{\alpha_i - \frac{N_i}{p} + (1 - \frac{1}{r}) \frac{N_i}{q}} \prod_{i=m+1}^n \left(\frac{\lambda_i}{\lambda_m} \right)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})}.
\end{aligned} \tag{5.8}$$

Recall **Remark 4.1**. $\gamma \geq 0, \delta \leq 0$ satisfy the two equivalent strict inequalities in (4.29)-(4.30). Define $0 \leq \vartheta \leq 1$ implicitly by requiring $\lambda_m = (\lambda_n)^\vartheta$. For r sufficiently close to 1, we find

$$\vartheta \left[\frac{N_1}{q r} + \sum_{i=2}^n \alpha_i - N_i \left(\frac{1}{p} - \frac{1}{q} \right) - \gamma \right] + (1 - \vartheta) \left[\alpha_n - N_n \left(\frac{1}{p} - \frac{1}{q} \right) \right] > 0 \tag{5.9}$$

and

$$\alpha_i - \frac{\mathbf{N}_i}{p} + \left(1 - \frac{1}{r}\right) \frac{\mathbf{N}_i}{q} < 0, \quad i = 1, 2, \dots, n. \quad (5.10)$$

Note that $\alpha_n - \mathbf{N}_n \left(\frac{1}{p} - \frac{1}{q}\right) > 0$. By using (5.9)-(5.10), we conclude that (5.8) is bounded by $\mathfrak{B}_{p,q,r,\gamma,\delta}(\lambda_n)^\varepsilon$ for some $\varepsilon = \varepsilon(p, q, r, \alpha, \gamma, \delta) > 0$.

Case Two: Let $\gamma \leq 0, \delta \geq 0$ satisfying (2.2)-(2.3) and (2.5). By adjusting the value of r , assume

$$\sum_{i=1}^{m-1} \mathbf{N}_i < \delta \left(\frac{pr}{p-1} \right) < \sum_{i=1}^m \mathbf{N}_i, \quad 1 \leq m \leq n. \quad (5.11)$$

Suppose that \mathbf{Q} is centered on some $z \in \mathbb{R}^{\mathbf{N}}$ with $|z| \leq 3$. We have

$$\begin{aligned} & \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{\mathbf{N}_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q r} dx \right\}^{\frac{1}{qr}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\delta \frac{pr}{p-1}} dx \right\}^{\frac{p-1}{pr}} \\ & \leq \mathfrak{B}_{p,q,r,\gamma,\delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)} \left\{ \prod_{i=1}^n \left(\frac{1}{\lambda_i} \right)^{\mathbf{N}_i} \int \cdots \int_{\bigotimes_{i=1}^n \mathbf{Q}_i} \left[\frac{1}{|x_1| + \cdots + |x_n|} \right]^{\delta \frac{pr}{p-1}} dx_1 \cdots dx_n \right\}^{\frac{p-1}{pr}} \\ & \quad (\gamma \leq 0) \\ & \leq \mathfrak{B}_{p,q,r,\gamma,\delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)} \prod_{i=1}^n \left(\frac{1}{\lambda_i} \right)^{\frac{p-1}{pr} \mathbf{N}_i} \\ & \quad \left\{ \int \cdots \int_{\bigotimes_{i=m}^n \mathbf{Q}_i} \left\{ \int \cdots \int_{\bigotimes_{i=1}^{m-1} \mathbb{R}^{\mathbf{N}_i}} \left[\frac{1}{|x_1| + \cdots + |x_n|} \right]^{\delta \frac{pr}{p-1}} dx_1 \cdots dx_{m-1} \right\} dx_m \cdots dx_n \right\}^{\frac{p-1}{pr}} \\ & \leq \mathfrak{B}_{p,q,r,\gamma,\delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)} \prod_{i=1}^n \left(\frac{1}{\lambda_i} \right)^{\frac{p-1}{pr} \mathbf{N}_i} \\ & \quad \left\{ \int \cdots \int_{\bigotimes_{i=m}^n \mathbf{Q}_i} \left[\frac{1}{|x_m| + \cdots + |x_n|} \right]^{\delta \frac{pr}{p-1} - \sum_{i=1}^{m-1} \mathbf{N}_i} dx_m \cdots dx_n \right\}^{\frac{p-1}{pr}} \\ & \leq \mathfrak{B}_{p,q,r,\gamma,\delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)} \prod_{i=1}^n \left(\frac{1}{\lambda_i} \right)^{\frac{p-1}{pr} \mathbf{N}_i} \left\{ \int \cdots \int_{\bigotimes_{i=m}^n \mathbf{Q}_i} \left(\frac{1}{|x_m|} \right)^{\delta \frac{pr}{p-1} - \sum_{i=1}^{m-1} \mathbf{N}_i} dx_m \cdots dx_n \right\}^{\frac{p-1}{pr}} \\ & \leq \mathfrak{B}_{p,q,r,\gamma,\delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)} \prod_{i=1}^m \left(\frac{1}{\lambda_i} \right)^{\frac{p-1}{pr} \mathbf{N}_i} \left\{ \int_{\mathbf{Q}_m^*} \left(\frac{1}{|x_m|} \right)^{\delta \frac{pr}{p-1} - \sum_{i=1}^{m-1} \mathbf{N}_i} dx_m \right\}^{\frac{p-1}{pr}} \quad \text{by Remark 5.1} \\ & \leq \mathfrak{B}_{p,q,r,\gamma,\delta} (\lambda_m)^{\frac{p-1}{pr} \sum_{i=1}^m \mathbf{N}_i - \delta} \prod_{i=1}^m \left(\frac{1}{\lambda_i} \right)^{\frac{p-1}{pr} \mathbf{N}_i} \prod_{i=1}^n (\lambda_i)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)}. \end{aligned} \quad (5.12)$$

From direct computation, the formula in the last line of (5. 12) can be rewritten as

$$\begin{aligned}
& (\lambda_m)^{\frac{p-1}{pr} \sum_{i=1}^m \mathbf{N}_i - \delta} \prod_{i=1}^m (\lambda_i)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right) - \frac{p-1}{pr} \mathbf{N}_i} \prod_{i=m+1}^n (\lambda_i)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)} \\
&= (\lambda_m)^{\frac{p-1}{pr} \sum_{i=1}^m \mathbf{N}_i - \delta} \prod_{i=2}^m (\lambda_i)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right) - \frac{p-1}{pr} \mathbf{N}_i} \prod_{i=m+1}^n (\lambda_i)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)} \quad (\lambda_1 = 1) \quad (5. 13) \\
&= (\lambda_m)^{\frac{p-1}{pr} \mathbf{N}_1 + \sum_{i=2}^n \alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right) - \delta} \prod_{i=2}^m \left(\frac{\lambda_i}{\lambda_m} \right)^{\alpha_i - \mathbf{N}_i \frac{q-1}{q} + \left(1 - \frac{1}{r} \right) \frac{p-1}{p} \mathbf{N}_i} \prod_{i=m+1}^n \left(\frac{\lambda_i}{\lambda_m} \right)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)}.
\end{aligned}$$

Recall **Remark 4.1**. $\gamma \leq 0, \delta \geq 0$ satisfy the two equivalent strict inequalities in (4. 30). Define $0 \leq \vartheta \leq 1$ implicitly by requiring $\lambda_m = (\lambda_n)^\vartheta$. For r sufficiently close to 1, we find

$$\vartheta \left[\mathbf{N}_1 \left(\frac{p-1}{pr} \right) + \sum_{i=2}^n \alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right) - \delta \right] + (1 - \vartheta) \left[\alpha_n - \mathbf{N}_n \left(\frac{1}{p} - \frac{1}{q} \right) \right] > 0 \quad (5. 14)$$

and

$$\alpha_i - \mathbf{N}_i \left(\frac{q-1}{q} \right) + \left(1 - \frac{1}{r} \right) \left(\frac{p-1}{p} \right) \mathbf{N}_i < 0, \quad i = 1, 2, \dots, n. \quad (5. 15)$$

Note that $\alpha_n - \mathbf{N}_n \left(\frac{1}{p} - \frac{1}{q} \right) > 0$. By using (5. 14)-(5. 15), we conclude that (5. 13) is bounded by $\mathfrak{B}_{p,q,r,\gamma,\delta}(\lambda_n)^\varepsilon$ for some $\varepsilon = \varepsilon(p, q, r, \alpha, \gamma, \delta) > 0$.

Suppose that \mathbf{Q} is centered on some $z \in \mathbb{R}^N$ with $|z| > 3$. Because \mathbf{Q} has a diameter 1, we have

$$\frac{1}{2}|z| \leq |x| \leq 2|z| \quad (5. 16)$$

whenever $x \in \mathbf{Q}$. By using (5. 16), we obtain

$$\begin{aligned}
& \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{\mathbf{N}_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q r} dx \right\}^{\frac{1}{qr}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\delta \left(\frac{pr}{p-1} \right)} dx \right\}^{\frac{p-1}{pr}} \\
& \leq \mathfrak{B}_{\gamma,\delta} \left(\frac{1}{|z|} \right)^{\gamma + \delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)} \leq \mathfrak{B}_{\gamma,\delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)}. \quad (\gamma + \delta \geq 0)
\end{aligned} \quad (5. 17)$$

Case Three: Let $\gamma > 0, \delta > 0$ satisfying (2. 2)-(2. 3) and (2. 6). By adjusting the value of r , assume

$$\sum_{i=1}^{m-1} \mathbf{N}_i < \gamma q r < \sum_{i=1}^m \mathbf{N}_i, \quad 1 \leq m \leq n, \quad (5. 18)$$

$$\sum_{i=1}^{\ell-1} \mathbf{N}_i < \delta \left(\frac{pr}{p-1} \right) < \sum_{i=1}^{\ell} \mathbf{N}_i, \quad 1 \leq \ell \leq n. \quad (5. 19)$$

We have

$$\begin{aligned}
& \prod_{i=1}^n |\mathbf{Q}_i|^{\frac{\alpha_i}{N_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\gamma q r} dx \right\}^{\frac{1}{q r}} \left\{ \frac{1}{|\mathbf{Q}|} \int_{\mathbf{Q}} \left(\frac{1}{|x|} \right)^{\delta \frac{p r}{p-1}} dx \right\}^{\frac{p-1}{p r}} \\
& \leq \mathfrak{B}_{p \ q \ r \ \gamma \ \delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=1}^n \left(\frac{1}{\lambda_i} \right)^{\frac{N_i}{q r}} \prod_{i=1}^n \left(\frac{1}{\lambda_i} \right)^{N_i \frac{p-1}{p r}} \\
& \quad \left\{ \int \cdots \int_{\otimes_{i=m}^n \mathbf{Q}_i} \left\{ \int \cdots \int_{\otimes_{i=1}^{m-1} \mathbb{R}^{N_i}} \left[\frac{1}{|x_1| + \cdots + |x_n|} \right]^{\gamma q r} dx_1 \cdots dx_{m-1} \right\} dx_m \cdots dx_n \right\}^{\frac{1}{q r}} \\
& \quad \left\{ \int \cdots \int_{\otimes_{i=\ell}^n \mathbf{Q}_i} \left\{ \int \cdots \int_{\otimes_{i=1}^{\ell-1} \mathbb{R}^{N_i}} \left[\frac{1}{|x_1| + \cdots + |x_n|} \right]^{\delta \frac{p r}{p-1}} dx_1 \cdots dx_{\ell-1} \right\} dx_{\ell} \cdots dx_n \right\}^{\frac{p-1}{p r}} \\
& \leq \mathfrak{B}_{p \ q \ r \ \gamma \ \delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=1}^n \left(\frac{1}{\lambda_i} \right)^{\frac{N_i}{q r}} \prod_{i=1}^n \left(\frac{1}{\lambda_i} \right)^{N_i \frac{p-1}{p r}} \\
& \quad \left\{ \int \cdots \int_{\otimes_{i=m}^n \mathbf{Q}_i} \left[\frac{1}{|x_m| + \cdots + |x_n|} \right]^{\gamma q r - \sum_{i=1}^{m-1} N_i} dx_m \cdots dx_n \right\}^{\frac{1}{q r}} \\
& \quad \left\{ \int \cdots \int_{\otimes_{i=\ell}^n \mathbf{Q}_i} \left[\frac{1}{|x_{\ell}| + \cdots + |x_n|} \right]^{\delta \frac{p r}{p-1} - \sum_{i=1}^{\ell-1} N_i} dx_{\ell} \cdots dx_n \right\}^{\frac{p-1}{p r}} \\
& \leq \mathfrak{B}_{p \ q \ r \ \gamma \ \delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=1}^n \left(\frac{1}{\lambda_i} \right)^{\frac{N_i}{q r}} \prod_{i=1}^n \left(\frac{1}{\lambda_i} \right)^{N_i \frac{p-1}{p r}} \\
& \quad \left\{ \int \cdots \int_{\otimes_{i=m}^n \mathbf{Q}_i} \left(\frac{1}{|x_m|} \right)^{\gamma q r - \sum_{i=1}^{m-1} N_i} dx_m \cdots dx_n \right\}^{\frac{1}{q r}} \\
& \quad \left\{ \int \cdots \int_{\otimes_{i=\ell}^n \mathbf{Q}_i} \left(\frac{1}{|x_{\ell}|} \right)^{\delta \frac{p r}{p-1} - \sum_{i=1}^{\ell-1} N_i} dx_{\ell} \cdots dx_n \right\}^{\frac{p-1}{p r}} \\
& \leq \mathfrak{B}_{p \ q \ r \ \gamma \ \delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=1}^m \left(\frac{1}{\lambda_i} \right)^{\frac{N_i}{q r}} \prod_{i=1}^{\ell} \left(\frac{1}{\lambda_i} \right)^{N_i \frac{p-1}{p r}} \\
& \quad \left\{ \int_{\mathbf{Q}_m^*} \left(\frac{1}{|x_m|} \right)^{\gamma q r - \sum_{i=1}^{m-1} N_i} dx_m \right\}^{\frac{1}{q r}} \left\{ \int_{\mathbf{Q}_{\ell}^*} \left(\frac{1}{|x_{\ell}|} \right)^{\delta \frac{p r}{p-1} - \sum_{i=1}^{\ell-1} N_i} dx_{\ell} \right\}^{\frac{p-1}{p r}} \quad \text{by Remark 5.1} \\
& \leq \mathfrak{B}_{p \ q \ r \ \gamma \ \delta} (\lambda_m)^{\frac{1}{q r} \sum_{i=1}^m N_i - \gamma} (\lambda_{\ell})^{\left(\frac{p-1}{p r} \right) \sum_{i=1}^{\ell} N_i - \delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=1}^m \left(\frac{1}{\lambda_i} \right)^{\frac{N_i}{q r}} \prod_{i=1}^{\ell} \left(\frac{1}{\lambda_i} \right)^{\frac{p-1}{p r} N_i}.
\end{aligned}$$

(5. 20)

Let $0 \leq k \leq n-1$. From direct computation, we find

$$\begin{aligned}
& \frac{1}{r} \left[\frac{1}{q} + \frac{p-1}{p} \right] \sum_{i=1}^k \mathbf{N}_i - (\gamma + \delta) + \sum_{i=k+1}^n \alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right) \\
&= \frac{\mathbf{N}}{r} - \frac{1}{r} \left(\frac{1}{p} - \frac{1}{q} \right) \mathbf{N} - (\gamma + \delta) + \sum_{i=k+1}^n \alpha_i - \frac{\mathbf{N}_i}{r} - \mathbf{N}_i \left(1 - \frac{1}{r} \right) \left(\frac{1}{p} - \frac{1}{q} \right) \\
&= \frac{\mathbf{N}}{r} - \frac{1}{r} \left(\frac{1}{p} - \frac{1}{q} \right) \mathbf{N} - \alpha + \mathbf{N} \left(\frac{1}{p} - \frac{1}{q} \right) + \sum_{i=k+1}^n \alpha_i - \frac{\mathbf{N}_i}{r} - \mathbf{N}_i \left(1 - \frac{1}{r} \right) \left(\frac{1}{p} - \frac{1}{q} \right) \quad \text{by (2. 3)} \\
&= \left(\frac{\mathbf{N}}{r} - \alpha \right) + \mathbf{N} \left(1 - \frac{1}{r} \right) \left(\frac{1}{p} - \frac{1}{q} \right) + \sum_{i=k+1}^n \alpha_i - \frac{\mathbf{N}_i}{r} - \mathbf{N}_i \left(1 - \frac{1}{r} \right) \left(\frac{1}{p} - \frac{1}{q} \right) \\
&= \sum_{i=1}^k \frac{\mathbf{N}_i}{r} - \alpha_i + \mathbf{N}_i \left(1 - \frac{1}{r} \right) \left(\frac{1}{p} - \frac{1}{q} \right).
\end{aligned} \tag{5. 21}$$

Suppose $\ell \leq m$. The formula in the last line of (5. 20) can be rewritten as

$$\begin{aligned}
& (\lambda_m)^{\frac{1}{r} \left[\frac{1}{q} + \frac{p-1}{p} \right] \sum_{i=1}^{\ell-1} \mathbf{N}_i - (\gamma + \delta)} \left(\frac{\lambda_\ell}{\lambda_m} \right)^{\frac{p-1}{pr} \sum_{i=1}^{\ell-1} \mathbf{N}_i - \delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)} \prod_{i=\ell}^m \left(\frac{\lambda_m}{\lambda_i} \right)^{\frac{\mathbf{N}_i}{qr}} \prod_{i=1}^{\ell-1} \left(\frac{1}{\lambda_i} \right)^{\frac{1}{r} \left[\frac{1}{q} + \frac{p-1}{p} \right] \mathbf{N}_i} \\
&= (\lambda_m)^{\frac{1}{r} \left[\frac{1}{q} + \frac{p-1}{p} \right] \sum_{i=1}^{\ell-1} \mathbf{N}_i + \sum_{i=\ell}^n \alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right) - (\gamma + \delta)} \left(\frac{\lambda_\ell}{\lambda_m} \right)^{\frac{p-1}{pr} \sum_{i=1}^{\ell-1} \mathbf{N}_i - \delta} \\
& \quad \prod_{i=\ell}^n \left(\frac{\lambda_i}{\lambda_m} \right)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)} \prod_{i=\ell}^m \left(\frac{\lambda_m}{\lambda_i} \right)^{\frac{\mathbf{N}_i}{qr}} \prod_{i=1}^{\ell-1} \left(\frac{1}{\lambda_i} \right)^{\frac{\mathbf{N}_i}{r} - \alpha_i + \left(1 - \frac{1}{r} \right) \left(\frac{1}{p} - \frac{1}{q} \right) \mathbf{N}_i} \\
&= \left(\frac{\lambda_\ell}{\lambda_m} \right)^{\frac{p-1}{pr} \sum_{i=1}^{\ell-1} \mathbf{N}_i - \delta} \prod_{i=\ell}^n \left(\frac{\lambda_i}{\lambda_m} \right)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)} \prod_{i=\ell}^m \left(\frac{\lambda_m}{\lambda_i} \right)^{\frac{\mathbf{N}_i}{qr}} \prod_{i=1}^{\ell-1} \left(\frac{\lambda_m}{\lambda_i} \right)^{\frac{\mathbf{N}_i}{r} - \alpha_i + \left(1 - \frac{1}{r} \right) \left(\frac{1}{p} - \frac{1}{q} \right) \mathbf{N}_i} \quad \text{by (5. 21)} \\
&= \left(\frac{\lambda_\ell}{\lambda_m} \right)^{\frac{p-1}{pr} \sum_{i=1}^{\ell-1} \mathbf{N}_i - \delta} \prod_{i=m+1}^n \left(\frac{\lambda_i}{\lambda_m} \right)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)} \prod_{i=\ell}^m \left(\frac{\lambda_i}{\lambda_m} \right)^{\alpha_i - \frac{\mathbf{N}_i}{p} + \left(1 - \frac{1}{r} \right) \frac{\mathbf{N}_i}{q}} \prod_{i=1}^{\ell-1} \left(\frac{\lambda_m}{\lambda_i} \right)^{\frac{\mathbf{N}_i}{r} - \alpha_i + \left(1 - \frac{1}{r} \right) \left(\frac{1}{p} - \frac{1}{q} \right) \mathbf{N}_i} \\
&= \prod_{i=m+1}^n \left(\frac{\lambda_i}{\lambda_m} \right)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)} \prod_{i=1}^{\ell-1} \left(\frac{\lambda_\ell}{\lambda_i} \right)^{\frac{\mathbf{N}_i}{r} - \alpha_i + \left(1 - \frac{1}{r} \right) \left(\frac{1}{p} - \frac{1}{q} \right) \mathbf{N}_i} \\
& \quad \left(\frac{\lambda_m}{\lambda_\ell} \right)^{\sum_{i=1}^{\ell-1} \frac{\mathbf{N}_i}{p} - \alpha_i - \left(1 - \frac{1}{r} \right) \frac{\mathbf{N}_i}{q} + \delta} \prod_{i=\ell}^m \left(\frac{\lambda_m}{\lambda_i} \right)^{\frac{\mathbf{N}_i}{p} - \alpha_i - \left(1 - \frac{1}{r} \right) \frac{\mathbf{N}_i}{q}}.
\end{aligned} \tag{5. 22}$$

Recall the subset \mathcal{U} defined in (2. 6) where $\alpha_i - \mathbf{N}_i/p < 0$ for every $i \notin \mathcal{U}$.

Note that $\lambda_m \leq \lambda_\ell$ when $\ell \leq m$. For r sufficiently close to 1, we have

$$\begin{aligned}
& \left(\frac{\lambda_m}{\lambda_\ell} \right)^{\sum_{i=1}^{\ell-1} \frac{N_i}{p} - \alpha_i - (1 - \frac{1}{r}) \frac{N_i}{q} + \delta} \prod_{i=\ell}^m \left(\frac{\lambda_m}{\lambda_i} \right)^{\frac{N_i}{p} - \alpha_i - (1 - \frac{1}{r}) \frac{N_i}{q}} \\
& \leq \left(\frac{\lambda_m}{\lambda_\ell} \right)^{\sum_{i \in \mathcal{U} \cap \{1, \dots, \ell-1\}} \frac{N_i}{p} - \alpha_i - (1 - \frac{1}{r}) \frac{N_i}{q} + \delta} \left(\frac{\lambda_m}{\lambda_\ell} \right)^{\sum_{i \in \mathcal{U} \cap \{\ell, \dots, m\}} \frac{N_i}{p} - \alpha_i - (1 - \frac{1}{r}) \frac{N_i}{q}} \\
& \quad \left(\frac{\lambda_m}{\lambda_\ell} \right)^{\sum_{i \in \mathcal{U}^c \cap \{1, \dots, \ell-1\}} \frac{N_i}{p} - \alpha_i - (1 - \frac{1}{r}) \frac{N_i}{q}} \prod_{i \in \mathcal{U}^c \cap \{\ell, \dots, m\}} \left(\frac{\lambda_m}{\lambda_i} \right)^{\frac{N_i}{p} - \alpha_i - (1 - \frac{1}{r}) \frac{N_i}{q}} \\
& \leq \left(\frac{\lambda_m}{\lambda_\ell} \right)^{\sum_{i \in \mathcal{U} \cap \{1, \dots, m\}} \frac{N_i}{p} - \alpha_i - (1 - \frac{1}{r}) \frac{N_i}{q} + \delta}.
\end{aligned} \tag{5. 23}$$

By bringing the estimates in (5. 22)-(5. 23) back to (5. 20), we find

$$\begin{aligned}
& (\lambda_m)^{\frac{1}{qr} \sum_{i=1}^m N_i - \gamma} (\lambda_\ell)^{\frac{p-1}{pr} \sum_{i=1}^\ell N_i - \delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=1}^m \left(\frac{1}{\lambda_i} \right)^{\frac{N_i}{qr}} \prod_{i=1}^\ell \left(\frac{1}{\lambda_i} \right)^{\frac{p-1}{pr} N_i} \\
& = \left(\frac{\lambda_m}{\lambda_\ell} \right)^{\sum_{i=1}^{\ell-1} \frac{N_i}{p} - \alpha_i - (1 - \frac{1}{r}) \frac{N_i}{q} + \delta} \prod_{i=\ell}^m \left(\frac{\lambda_m}{\lambda_i} \right)^{\frac{N_i}{p} - \alpha_i - (1 - \frac{1}{r}) \frac{N_i}{q}} \\
& \quad \prod_{i=m+1}^n \left(\frac{\lambda_i}{\lambda_m} \right)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=1}^{\ell-1} \left(\frac{\lambda_\ell}{\lambda_i} \right)^{\frac{N_i}{r} - \alpha_i + (1 - \frac{1}{r}) (\frac{1}{p} - \frac{1}{q}) N_i} \\
& \leq \left(\frac{\lambda_m}{\lambda_\ell} \right)^{\sum_{i \in \mathcal{U} \cap \{1, 2, \dots, m\}} \frac{N_i}{p} - \alpha_i - (1 - \frac{1}{r}) \frac{N_i}{q} + \delta} \prod_{i=1}^\ell \left(\frac{\lambda_\ell}{\lambda_i} \right)^{\frac{N_i}{r} - \alpha_i + (1 - \frac{1}{r}) (\frac{1}{p} - \frac{1}{q}) N_i} \prod_{i=m+1}^n \left(\frac{\lambda_i}{\lambda_m} \right)^{\alpha_i - N_i (\frac{1}{p} - \frac{1}{q})}.
\end{aligned} \tag{5. 24}$$

Recall that $\delta > 0$ satisfies the first strict inequality in (2. 6). From (5. 1) and (1. 6), we also have $(\frac{1}{p} - \frac{1}{q}) N_i < \alpha_i < N_i$ for every $i = 1, 2, \dots, n$. Define implicitly $0 \leq \vartheta_1 \leq \vartheta_2 \leq 1$ by requiring $\lambda_\ell = (\lambda_n)^{\vartheta_1}$ and $\lambda_m = (\lambda_n)^{\vartheta_2}$.

For r sufficiently close to 1, we have

$$\begin{aligned}
& \vartheta_1 \left[\frac{N_1}{r} - \alpha_1 + \left(1 - \frac{1}{r} \right) \left(\frac{1}{p} - \frac{1}{q} \right) N_1 \right] + (1 - \vartheta_2) \left(\alpha_n - N_n \left(\frac{1}{p} - \frac{1}{q} \right) \right) \\
& + (\vartheta_2 - \vartheta_1) \left[\sum_{i \in \mathcal{U} \cup \{1, 2, \dots, m\}} \frac{N_i}{p} - \alpha_i - \left(1 - \frac{1}{r} \right) \frac{N_i}{q} + \delta \right] > 0.
\end{aligned} \tag{5. 25}$$

The estimate in (5. 25) implies that (5. 24) is bounded by $\mathfrak{B}_{pqr\alpha\gamma\delta}(\lambda_n)^\varepsilon$ for some $\varepsilon = \varepsilon(pqr\alpha\gamma\delta) > 0$.

On the other hand, suppose $m \leq \ell$. The last line of (5. 20) can be rewritten as

$$\begin{aligned}
& (\lambda_m)^{\frac{1}{qr} \sum_{i=1}^{m-1} \mathbf{N}_i - \gamma} (\lambda_\ell)^{\frac{p-1}{pr} \sum_{i=1}^\ell \mathbf{N}_i - \delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - \mathbf{N}_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=1}^{m-1} \left(\frac{1}{\lambda_i} \right)^{\frac{\mathbf{N}_i}{qr}} \prod_{i=1}^\ell \left(\frac{1}{\lambda_i} \right)^{\frac{p-1}{pr} \mathbf{N}_i} \\
&= (\lambda_\ell)^{\frac{1}{r} [\frac{1}{q} + \frac{p-1}{p}] \sum_{i=1}^{m-1} \mathbf{N}_i - (\gamma + \delta)} \left(\frac{\lambda_m}{\lambda_\ell} \right)^{\frac{1}{qr} \sum_{i=1}^{m-1} \mathbf{N}_i - \gamma} \prod_{i=1}^n (\lambda_i)^{\alpha_i - \mathbf{N}_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=m}^\ell \left(\frac{\lambda_\ell}{\lambda_i} \right)^{\frac{p-1}{pr} \mathbf{N}_i} \prod_{i=1}^{m-1} \left(\frac{1}{\lambda_i} \right)^{\frac{1}{r} [\frac{1}{q} + \frac{p-1}{p}] \mathbf{N}_i} \\
&= (\lambda_\ell)^{\frac{1}{r} [\frac{1}{q} + \frac{p-1}{p}] \sum_{i=1}^{m-1} \mathbf{N}_i + \sum_{i=m}^n \alpha_i - \mathbf{N}_i (\frac{1}{p} - \frac{1}{q}) - (\gamma + \delta)} \left(\frac{\lambda_m}{\lambda_\ell} \right)^{\frac{1}{qr} \sum_{i=1}^{m-1} \mathbf{N}_i - \gamma} \\
&\quad \prod_{i=m}^n \left(\frac{\lambda_i}{\lambda_\ell} \right)^{\alpha_i - \mathbf{N}_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=m}^\ell \left(\frac{\lambda_\ell}{\lambda_i} \right)^{\frac{p-1}{pr} \mathbf{N}_i} \prod_{i=1}^{m-1} \left(\frac{1}{\lambda_i} \right)^{\frac{\mathbf{N}_i}{r} - \alpha_i + (1 - \frac{1}{r}) (\frac{1}{p} - \frac{1}{q}) \mathbf{N}_i} \\
&= \left(\frac{\lambda_m}{\lambda_\ell} \right)^{\frac{1}{qr} \sum_{i=1}^{m-1} \mathbf{N}_i - \gamma} \prod_{i=m}^n \left(\frac{\lambda_i}{\lambda_\ell} \right)^{\alpha_i - \mathbf{N}_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=m}^\ell \left(\frac{\lambda_\ell}{\lambda_i} \right)^{\frac{p-1}{pr} \mathbf{N}_i} \prod_{i=1}^{m-1} \left(\frac{\lambda_\ell}{\lambda_i} \right)^{\frac{\mathbf{N}_i}{r} - \alpha_i + (1 - \frac{1}{r}) (\frac{1}{p} - \frac{1}{q}) \mathbf{N}_i} \quad \text{by (5. 21)} \\
&= \left(\frac{\lambda_m}{\lambda_\ell} \right)^{\frac{1}{qr} \sum_{i=1}^{m-1} \mathbf{N}_i - \gamma} \prod_{i=\ell+1}^n \left(\frac{\lambda_i}{\lambda_\ell} \right)^{\alpha_i - \mathbf{N}_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=m}^\ell \left(\frac{\lambda_i}{\lambda_\ell} \right)^{\alpha_i - \frac{q-1}{q} \mathbf{N}_i + (1 - \frac{1}{r}) \frac{p-1}{p} \mathbf{N}_i} \prod_{i=1}^{m-1} \left(\frac{\lambda_\ell}{\lambda_i} \right)^{\frac{\mathbf{N}_i}{r} - \alpha_i + (1 - \frac{1}{r}) (\frac{1}{p} - \frac{1}{q}) \mathbf{N}_i} \\
&= \prod_{i=\ell+1}^n \left(\frac{\lambda_i}{\lambda_\ell} \right)^{\alpha_i - \mathbf{N}_i (\frac{1}{p} - \frac{1}{q})} \prod_{i=1}^m \left(\frac{\lambda_m}{\lambda_i} \right)^{\frac{\mathbf{N}_i}{r} - \alpha_i + (1 - \frac{1}{r}) (\frac{1}{p} - \frac{1}{q}) \mathbf{N}_i} \\
&\quad \left(\frac{\lambda_\ell}{\lambda_m} \right)^{\sum_{i=1}^{m-1} \frac{q-1}{q} \mathbf{N}_i - \alpha_i - (1 - \frac{1}{r}) \frac{p-1}{p} \mathbf{N}_i + \gamma} \prod_{i=m}^\ell \left(\frac{\lambda_\ell}{\lambda_i} \right)^{\frac{q-1}{q} \mathbf{N}_i - \alpha_i - (1 - \frac{1}{r}) \frac{p-1}{p} \mathbf{N}_i}.
\end{aligned} \tag{5. 26}$$

Recall the subset \mathcal{V} defined in (2. 6) where $\alpha_i - \mathbf{N}_i (\frac{q-1}{q}) < 0$ for every $i \notin \mathcal{V}$.

Note that $\lambda_\ell \leq \lambda_m$ when $m \leq \ell$. For r sufficiently close to 1, we find

$$\begin{aligned}
& \left(\frac{\lambda_\ell}{\lambda_m} \right)^{\sum_{i=1}^{m-1} \frac{q-1}{q} \mathbf{N}_i - \alpha_i - (1 - \frac{1}{r}) \frac{p-1}{p} \mathbf{N}_i + \gamma} \prod_{i=m}^\ell \left(\frac{\lambda_\ell}{\lambda_i} \right)^{\frac{q-1}{q} \mathbf{N}_i - \alpha_i - (1 - \frac{1}{r}) \frac{p-1}{p} \mathbf{N}_i} \\
&\leq \left(\frac{\lambda_\ell}{\lambda_m} \right)^{\sum_{i \in \mathcal{V} \cap \{1, \dots, m-1\}} \frac{q-1}{q} \mathbf{N}_i - \alpha_i - (1 - \frac{1}{r}) \frac{p-1}{p} \mathbf{N}_i + \gamma} \left(\frac{\lambda_\ell}{\lambda_m} \right)^{\sum_{i \in \mathcal{V} \cap \{m, \dots, \ell\}} \frac{q-1}{q} \mathbf{N}_i - \alpha_i - (1 - \frac{1}{r}) \frac{p-1}{p} \mathbf{N}_i} \\
&\quad \left(\frac{\lambda_\ell}{\lambda_m} \right)^{\sum_{i \in \mathcal{V}^c \cap \{1, \dots, m-1\}} \frac{q-1}{q} \mathbf{N}_i - \alpha_i - (1 - \frac{1}{r}) \frac{p-1}{p} \mathbf{N}_i} \prod_{i \in \mathcal{V}^c \cap \{m, \dots, \ell\}} \left(\frac{\lambda_\ell}{\lambda_i} \right)^{\frac{q-1}{q} \mathbf{N}_i - \alpha_i - (1 - \frac{1}{r}) \frac{p-1}{p} \mathbf{N}_i} \\
&\leq \left(\frac{\lambda_\ell}{\lambda_m} \right)^{\sum_{i \in \mathcal{V} \cap \{1, \dots, \ell\}} \frac{q-1}{q} \mathbf{N}_i - \alpha_i - (1 - \frac{1}{r}) \frac{p-1}{p} \mathbf{N}_i + \gamma}.
\end{aligned} \tag{5. 27}$$

By bringing the estimates in (5. 26)-(5. 27) back to (5. 20), we find

$$\begin{aligned}
& (\lambda_m)^{\frac{1}{qr} \sum_{i=1}^m \mathbf{N}_i - \gamma} (\lambda_\ell)^{\frac{p-1}{pr} \sum_{i=1}^\ell \mathbf{N}_i - \delta} \prod_{i=1}^n (\lambda_i)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q}\right)} \prod_{i=1}^m \left(\frac{1}{\lambda_i}\right)^{\frac{\mathbf{N}_i}{qr}} \prod_{i=1}^\ell \left(\frac{1}{\lambda_i}\right)^{\frac{p-1}{pr} \mathbf{N}_i} \\
&= \prod_{i=\ell+1}^n \left(\frac{\lambda_i}{\lambda_\ell}\right)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q}\right)} \prod_{i=1}^m \left(\frac{\lambda_m}{\lambda_i}\right)^{\frac{\mathbf{N}_i}{r} - \alpha_i + \left(1 - \frac{1}{r}\right) \left(\frac{1}{p} - \frac{1}{q}\right) \mathbf{N}_i} \\
&\quad \left(\frac{\lambda_\ell}{\lambda_m}\right)^{\sum_{i=1}^{m-1} \frac{q-1}{q} \mathbf{N}_i - \alpha_i - \left(1 - \frac{1}{r}\right) \frac{p-1}{p} \mathbf{N}_i + \gamma} \prod_{i=m}^\ell \left(\frac{\lambda_\ell}{\lambda_i}\right)^{\frac{q-1}{q} \mathbf{N}_i - \alpha_i - \left(1 - \frac{1}{r}\right) \frac{p-1}{p} \mathbf{N}_i} \\
&\leq \left(\frac{\lambda_\ell}{\lambda_m}\right)^{\sum_{i \in \mathcal{V} \cap \{1, \dots, \ell\}} \frac{q-1}{q} \mathbf{N}_i - \alpha_i - \left(1 - \frac{1}{r}\right) \frac{p-1}{p} \mathbf{N}_i + \gamma} \prod_{i=1}^m \left(\frac{\lambda_m}{\lambda_i}\right)^{\frac{\mathbf{N}_i}{r} - \alpha_i + \left(1 - \frac{1}{r}\right) \left(\frac{1}{p} - \frac{1}{q}\right) \mathbf{N}_i} \prod_{i=\ell+1}^n \left(\frac{\lambda_i}{\lambda_\ell}\right)^{\alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q}\right)}. \tag{5. 28}
\end{aligned}$$

Recall that $\gamma > 0$ satisfies the second strict inequality in (2. 6). From (5. 1) and (1. 6), we also have $\left(\frac{1}{p} - \frac{1}{q}\right) \mathbf{N}_i < \alpha_i < \mathbf{N}_i$ for every $i = 1, 2, \dots, n$. Define implicitly $0 \leq \vartheta_1 \leq \vartheta_2 \leq 1$ by requiring $\lambda_m = (\lambda_n)^{\vartheta_1}$ and $\lambda_\ell = (\lambda_n)^{\vartheta_2}$. For r sufficiently close to 1, we have

$$\begin{aligned}
& \vartheta_1 \left[\frac{\mathbf{N}_1}{r} - \alpha_1 + \left(1 - \frac{1}{r}\right) \left(\frac{1}{p} - \frac{1}{q}\right) \mathbf{N}_1 \right] + (1 - \vartheta_2) \left(\alpha_n - \mathbf{N}_n \left(\frac{1}{p} - \frac{1}{q}\right) \right) \\
&+ (\vartheta_2 - \vartheta_1) \left[\sum_{i \in \mathcal{V} \cap \{1, \dots, \ell\}} \left(\frac{q-1}{q}\right) \mathbf{N}_i - \alpha_i - \left(1 - \frac{1}{r}\right) \left(\frac{p-1}{p}\right) \mathbf{N}_i + \gamma \right] > 0. \tag{5. 29}
\end{aligned}$$

The estimate in (5. 29) implies that (5. 28) is bounded by $\mathfrak{B}_{p,q,r,\alpha,\gamma,\delta}(\lambda_n)^\varepsilon$ for some $\varepsilon = \varepsilon(p, q, r, \alpha, \gamma, \delta) > 0$. \square

6 One-weight inequality on product spaces

In this section, we consider the special case of balance indices:

$$\frac{\alpha_i}{\mathbf{N}_i} = \frac{1}{p} - \frac{1}{q}, \quad i = 1, 2, \dots, n. \tag{6. 1}$$

Consequently, we must have $\frac{\alpha_i}{\mathbf{N}_i} = \frac{\alpha}{\mathbf{N}}$ for every $i = 1, 2, \dots, n$. Recall (2. 3). $\frac{\alpha}{\mathbf{N}} = \frac{1}{p} - \frac{1}{q}$ implies $\gamma + \delta = 0$. Consider $\omega(x) = |x|^{-\gamma}$, $\sigma(x) = |x|^\delta$ for $x \neq 0$ and $\gamma, \delta \in \mathbb{R}$. Our assertion reduces to the one-weight situation: $\omega = \sigma$.

Write $x = (x_i, x_i^\dagger) \in \mathbb{R}^{\mathbf{N}_i} \times \mathbb{R}^{\mathbf{N} - \mathbf{N}_i}$ and $\mathbf{Q}_i^\dagger = \bigotimes_{i \neq j} \mathbf{Q}_i$ for $i = 1, 2, \dots, n$. Shrink \mathbf{Q}_i^\dagger to x_i^\dagger in (2. 1). By applying the Lebesgue Differentiation Theorem, we find

$$\left\{ \frac{1}{|\mathbf{Q}_i|} \int_{\mathbf{Q}_i} \omega^q(x_i, x_i^\dagger) dx_i \right\}^{\frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}_i|} \int_{\mathbf{Q}_i} \left(\frac{1}{\omega}\right)^{\frac{p}{p-1}}(x_i, x_i^\dagger) dx_i \right\}^{\frac{p-1}{p}} < \infty \tag{6. 2}$$

for every $\mathbf{Q}_i \subset \mathbb{R}^{\mathbf{N}_i}$ and $x_i^\dagger \in \mathbb{R}^{\mathbf{N} - \mathbf{N}_i}$, $i = 1, 2, \dots, n$.

Observe that (6. 1)-(6. 2) are sufficient conditions of **Muckenhoupt-Wheeden theorem** [5] which implies

$$\begin{aligned} & \left\{ \int_{\mathbb{R}^{N_i}} \left\{ \int_{\mathbb{R}^{N_i}} f(y_i, x_i^\dagger) \left[\frac{1}{|x_i - y_i|} \right]^{N_i - \alpha_i} dy_i \right\}^q \omega^q(x_i, x_i^\dagger) dx_i \right\}^{\frac{1}{q}} \\ & \leq \mathfrak{B}_{p \ q \ \omega} \left\{ \int_{\mathbb{R}^{N_i}} (f\omega)^p(x_i, x_i^\dagger) dx_i \right\}^{\frac{1}{p}} \end{aligned} \quad (6. 3)$$

for $1 < p < q < \infty$ and $x_i^\dagger \in \mathbb{R}^{N-N_i}$, $i = 1, 2, \dots, n$.

By using (6. 3), we have

$$\begin{aligned} & \left\{ \int_{\mathbb{R}^N} (\omega \mathbf{I}_\alpha f)^q(x) dx \right\}^{\frac{1}{q}} = \left\{ \int_{\mathbb{R}^N} \left\{ \int_{\mathbb{R}^N} f(y) \prod_{i=1}^n \left[\frac{1}{|x_i - y_i|} \right]^{N_i - \alpha_i} dy \right\}^q \omega^q(x) dx \right\}^{\frac{1}{q}} \\ & \leq \mathfrak{B}_{p \ q \ \omega} \left\{ \int_{\mathbb{R}^{N-N_i}} \left\{ \int_{\mathbb{R}^{N_i}} \left\{ \int_{\mathbb{R}^{N-N_i}} f(x_i, y_i^\dagger) \prod_{j \neq i} \left[\frac{1}{|x_j - y_j|} \right]^{N_j - \alpha_j} dy_i^\dagger \right\}^p \omega^p(x_i, x_i^\dagger) dx_i \right\}^{\frac{q}{p}} dx_i^\dagger \right\}^{\frac{1}{q}} \\ & \leq \mathfrak{B}_{p \ q \ \omega} \left\{ \int_{\mathbb{R}^{N_i}} \left\{ \int_{\mathbb{R}^{N-N_i}} \left\{ \int_{\mathbb{R}^{N-N_i}} f(x_i, y_i^\dagger) \prod_{j \neq i} \left[\frac{1}{|x_j - y_j|} \right]^{N_j - \alpha_j} dy_i^\dagger \right\}^q \omega^q(x_i, x_i^\dagger) dx_i^\dagger \right\}^{\frac{p}{q}} dx_i \right\}^{\frac{1}{p}} \\ & \quad \vdots \quad \text{by Minkowski integral inequality} \\ & \leq \mathfrak{B}_{p \ q \ \omega} \left\{ \int_{\mathbb{R}^N} (f\omega)^p(x) dx \right\}^{\frac{1}{p}}, \quad 1 < p < q < \infty. \end{aligned} \quad (6. 4)$$

7 Proof of Theorem A*

Suppose that $\omega(x) = |x|^{-\gamma}$, $\sigma(x) = |x|^\delta$ for $x \neq 0$ and $\gamma, \delta \in \mathbb{R}$ satisfy the multi-parameter Muckenhoupt characteristic in (2. 1). Let $\{1, 2, \dots, n\} = \mathbf{U} \cup \mathbf{V}$ where

$$\mathbf{U} = \left\{ i \in \{1, 2, \dots, n\} : \frac{\alpha_i}{N_i} = \frac{1}{p} - \frac{1}{q} \right\}, \quad \mathbf{V} = \left\{ i \in \{1, 2, \dots, n\} : \frac{\alpha_i}{N_i} > \frac{1}{p} - \frac{1}{q} \right\}. \quad (7. 1)$$

Define

$$\begin{aligned} \alpha_{\mathbf{U}} &= \sum_{i \in \mathbf{U}} \alpha_i, \quad \mathbf{Q}_{\mathbf{U}} = \bigotimes_{i \in \mathbf{U}} \mathbf{Q}_i, \quad \mathbb{R}^{N_{\mathbf{U}}} = \bigotimes_{i \in \mathbf{U}} \mathbb{R}^{N_i}, \\ \alpha_{\mathbf{V}} &= \sum_{i \in \mathbf{V}} \alpha_i, \quad \mathbf{Q}_{\mathbf{V}} = \bigotimes_{i \in \mathbf{V}} \mathbf{Q}_i, \quad \mathbb{R}^{N_{\mathbf{V}}} = \bigotimes_{i \in \mathbf{V}} \mathbb{R}^{N_i}. \end{aligned} \quad (7. 2)$$

Write $x = (x_{\mathbf{U}}, x_{\mathbf{V}}) \in \mathbb{R}^{N_{\mathbf{U}}} \times \mathbb{R}^{N_{\mathbf{V}}}$. Denote the cardinality of \mathbf{U} and \mathbf{V} by $|\mathbf{U}|$ and $|\mathbf{V}|$.

Consider \mathbf{Q}_i centered on the origin of \mathbb{R}^{N_i} for $i \in \mathbf{U}$. Shrink $\mathbf{Q}_i, i \in \mathbf{U}$ to 0 in (2. 1). By applying Lebesgue Differentiation Theorem, we have

$$\begin{aligned} & \sup_{\mathbf{Q}_V \subset \mathbb{R}^{N_V}} \prod_{i \in \mathbf{V}} |\mathbf{Q}_i|^{\frac{\alpha_i}{N_i} - \frac{1}{p} + \frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}_V|} \int_{\mathbf{Q}_V} \left(\frac{1}{|x_V|} \right)^{q\gamma} dx_V \right\}^{\frac{1}{q}} \left\{ \frac{1}{|\mathbf{Q}_V|} \int_{\mathbf{Q}_V} \left(\frac{1}{|x_V|} \right)^{\frac{\delta p}{p-1}} dx_V \right\}^{\frac{p-1}{p}} \\ & \leq \mathbf{A}_{pq}^\alpha (|x|^{-\gamma}, |x|^\delta) < \infty. \end{aligned} \quad (7. 3)$$

Remark 7.1 By carrying out the same estimates in section 4, we find (7. 3) implying (2. 2)-(2. 6) with α, n, \mathbf{N} replaced by $\alpha_V, |\mathbf{V}|, \mathbf{N}_V$ respectively.

Proposition 7.1 Suppose $\omega(x_V) = |x_V|^{-\gamma}, \sigma(x_V) = |x_V|^\delta$ for $x_V \neq 0$ and $\gamma, \delta \in \mathbb{R}$ satisfying (7. 3). For a.e $x_U \in \mathbb{R}^{N_U}$, we have

$$\begin{aligned} & \left\{ \int_{\mathbb{R}^{N_V}} \left\{ \int_{\mathbb{R}^{N_V}} f(x_U, y_V) \prod_{i \in \mathbf{V}} \left[\frac{1}{|x_i - y_i|} \right]^{N_i - \alpha_i} dy_V \right\}^q \omega^q(x_V) dx_V \right\}^{\frac{1}{q}} \\ & \leq \mathfrak{B}_{p,q,\alpha,\gamma,\delta} \left\{ \int_{\mathbb{R}^{N_V}} (f(x_U, x_V))^p \sigma^p(x_V) dx_V \right\}^{\frac{1}{p}}, \quad 1 < p \leq q < \infty. \end{aligned} \quad (7. 4)$$

Proof: Let $|\mathbf{V}| = 1$. **Theorem A** by Stein and Weiss [3] together with **Remark 7.1** imply (7. 4).

Consider $|\mathbf{V}| \geq 2$. By applying **Principal Lemma** in the beginning of section 5, we have $\omega(x_V) = |x_V|^{-\gamma}, \sigma(x_V) = |x_V|^\delta$ satisfying the estimate in (5. 2)-(5. 3) for every $\mathbf{Q}_V \subset \mathbb{R}^{N_V}$ whereas $\alpha_i > N_i \left(\frac{1}{p} - \frac{1}{q} \right), i \in \mathbf{V}$.

Denote \mathbf{t} as a $|\mathbf{V}|$ -tuple $(2^{-t_1}, 2^{-t_2}, \dots, 2^{-t_{|\mathbf{V}|}})$. Recall $\mathbf{A}_{pqr}^\alpha(\mathbf{t}; \omega, \sigma)$ defined in (3. 6). We find

$$\sum_{\mathbf{t}} \mathbf{A}_{pqr}^{\alpha_V}(\mathbf{t}; |x_V|^{-\gamma}, |x_V|^\delta) < \infty \quad (7. 5)$$

for some $r = r(p, q, \alpha, \gamma, \delta) > 1$. This is the desired summability in (3. 14). \square

Remark 7.2 Note that $\mathbf{V} = \{1, 2, \dots, n\}$ when $1 < p = q < \infty$. **Proposition 7.1** already implies **Theorem A***.

For the remaining section, we assert $1 < p < q < \infty$ only.

In the case of $\gamma > 0, \delta > 0$, we observe

$$\omega(x) = |x|^{-\gamma} \leq |x_V|^{-\gamma} = \omega(x_V), \quad \sigma(x_V) = |x_V|^\delta \leq |x|^\delta = \sigma(x). \quad (7. 6)$$

From (7. 6), we have

$$\begin{aligned}
& \left\{ \int_{\mathbb{R}^N} (\omega \mathbf{I}_\alpha f)^q(x) dx \right\}^{\frac{1}{q}} \leq \left\{ \int_{\mathbb{R}^N} \left\{ \int_{\mathbb{R}^N} f(y) \prod_{i=1}^n \left[\frac{1}{|x_i - y_i|} \right]^{\mathbf{N}_i - \alpha_i} dy \right\}^q \omega^q(x_{\mathbf{V}}) dx \right\}^{\frac{1}{q}} \\
& \leq \mathfrak{B}_{p \ q \ \alpha \ \gamma \ \delta} \left\{ \int_{\mathbb{R}^{N_{\mathbf{U}}}} \left\{ \int_{\mathbb{R}^{N_{\mathbf{V}}}} \left\{ \int_{\mathbb{R}^{N_{\mathbf{U}}}} f(y_{\mathbf{U}}, x_{\mathbf{V}}) \prod_{i \in \mathbf{U}} \left[\frac{1}{|x_i - y_i|} \right]^{\mathbf{N}_i - \alpha_i} dy_{\mathbf{U}} \right\}^p \sigma^p(x_{\mathbf{V}}) dx_{\mathbf{V}} \right\}^{\frac{q}{p}} dx_{\mathbf{U}} \right\}^{\frac{1}{q}} \\
& \quad \text{by Proposition 7.1} \\
& \leq \mathfrak{B}_{p \ q \ \alpha \ \gamma \ \delta} \left\{ \int_{\mathbb{R}^{N_{\mathbf{V}}}} \left\{ \int_{\mathbb{R}^{N_{\mathbf{U}}}} \left\{ \int_{\mathbb{R}^{N_{\mathbf{U}}}} f(y_{\mathbf{U}}, x_{\mathbf{V}}) \prod_{i \in \mathbf{U}} \left[\frac{1}{|x_i - y_i|} \right]^{\mathbf{N}_i - \alpha_i} dy_{\mathbf{U}} \right\}^q dx_{\mathbf{U}} \right\}^{\frac{p}{q}} \sigma^p(x_{\mathbf{V}}) dx_{\mathbf{V}} \right\}^{\frac{1}{p}} \\
& \quad \text{by Minkowski integral inequality} \\
& \leq \mathfrak{B}_{p \ q \ \alpha \ \gamma \ \delta} \left\{ \iint_{\mathbb{R}^{N_{\mathbf{U}}} \times \mathbb{R}^{N_{\mathbf{V}}}} (f(x_{\mathbf{U}}, x_{\mathbf{V}}))^p \sigma^p(x_{\mathbf{V}}) dx_{\mathbf{U}} dx_{\mathbf{V}} \right\}^{\frac{1}{p}} \quad \text{by (6. 2)-(6. 4) with } \omega = 1 \\
& \leq \mathfrak{B}_{p \ q \ \alpha \ \gamma \ \delta} \left\{ \int_{\mathbb{R}^N} (f\sigma)^p(x) dx \right\}^{\frac{1}{p}}, \quad 1 < p < q < \infty.
\end{aligned} \tag{7. 7}$$

Consider $\gamma \geq 0, \delta \leq 0$ or $\gamma \leq 0, \delta \geq 0$. Note that it is suffice to study one of these two cases because \mathbf{I}_α is self-adjoint and

$$\|\omega \mathbf{I}_\alpha f\|_{\mathbf{L}^q(\mathbb{R}^N)} \lesssim \|f\sigma\|_{\mathbf{L}^p(\mathbb{R}^N)} \quad \text{if and only if} \quad \|\sigma^{-1} \mathbf{I}_\alpha g\|_{\mathbf{L}^{\frac{p}{p-1}}(\mathbb{R}^N)} \lesssim \|g\omega^{-1}\|_{\mathbf{L}^{\frac{q}{q-1}}(\mathbb{R}^N)}. \tag{7. 8}$$

Let $\gamma \geq 0, \delta \leq 0$. Suppose f supported in the region where $|x_{\mathbf{U}}| \leq |x_{\mathbf{V}}|$. By using (7. 6) and a repeat estimate in (7. 7), we obtain

$$\begin{aligned}
& \left\{ \int_{\mathbb{R}^N} \left\{ \int_{|y_{\mathbf{U}}| \leq |y_{\mathbf{V}}|} f(y) \prod_{i=1}^n \left[\frac{1}{|x_i - y_i|} \right]^{\mathbf{N}_i - \alpha_i} dy \right\}^q \omega^q(x) dx \right\}^{\frac{1}{q}} \\
& \leq \left\{ \iint_{\mathbb{R}^{N_{\mathbf{U}}} \times \mathbb{R}^{N_{\mathbf{V}}}} \left\{ \iint_{|y_{\mathbf{U}}| \leq |y_{\mathbf{V}}|} f(y_{\mathbf{U}}, y_{\mathbf{V}}) \prod_{i \in \mathbf{U} \cup \mathbf{V}} \left[\frac{1}{|x_i - y_i|} \right]^{\mathbf{N}_i - \alpha_i} dy_{\mathbf{U}} dy_{\mathbf{V}} \right\}^q \omega^q(x_{\mathbf{V}}) dx \right\}^{\frac{1}{q}} \\
& \quad \text{(7. 9)} \\
& \leq \mathfrak{B}_{p \ q \ \alpha \ \gamma \ \delta} \left\{ \iint_{|x_{\mathbf{U}}| \leq |x_{\mathbf{V}}|} (f(x_{\mathbf{U}}, x_{\mathbf{V}}))^p \sigma^p(x_{\mathbf{V}}) dx_{\mathbf{U}} dx_{\mathbf{V}} \right\}^{\frac{1}{p}} \\
& \leq \mathfrak{B}_{p \ q \ \alpha \ \gamma \ \delta} \left\{ \int_{\mathbb{R}^N} (f\sigma)^p(x) dx \right\}^{\frac{1}{p}}.
\end{aligned}$$

The last inequality holds in (7. 9) because $\sigma(x_{\mathbf{V}}) = |x_{\mathbf{V}}|^\delta \approx |x|^\delta = \sigma(x)$ for $|x_{\mathbf{U}}| \leq |x_{\mathbf{V}}|$.

On the other hand, suppose f supported in the region $|x_U| > |x_V|$. Recall that $\gamma \geq 0, \delta \leq 0$ satisfy (2. 2)-(2. 4). In particular,

$$\gamma + \delta = \sum_{i=1}^n \alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right) = \sum_{i \in \mathbf{V}} \alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right) \quad \text{by (7. 1)} \quad (7. 10)$$

and

$$\alpha_i - \frac{\mathbf{N}_i}{p} < \delta \leq 0 \quad \text{for every } i \in \{1, 2, \dots, n\} = \mathbf{U} \cup \mathbf{V}. \quad (7. 11)$$

By putting together (7. 10) and (7. 11), we find

$$0 \leq \gamma + \delta = \sum_{i \in \mathbf{V}} \alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right) < \frac{\mathbf{N}_V}{q}, \quad 0 < \mathbf{N}_V \left(\frac{p-1}{p} \right). \quad (7. 12)$$

Proposition 7.2 Let $\rho(x_V) = |x_V|^{-(\gamma+\delta)}$, $\eta(x_V) \equiv 1$. For a.e $x_U \in \mathbb{R}^{\mathbf{N}_U}$, we have

$$\begin{aligned} & \left\{ \int_{\mathbb{R}^{\mathbf{N}_V}} \left\{ \int_{\mathbb{R}^{\mathbf{N}_V}} f(x_U, y_V) \prod_{i \in \mathbf{V}} \left[\frac{1}{|x_i - y_i|} \right]^{\mathbf{N}_i - \alpha_i} dy_V \right\}^q \rho^q(x_V) dx_V \right\}^{\frac{1}{q}} \\ & \leq \mathfrak{B}_{p,q,\alpha,\gamma,\delta} \left\{ \int_{\mathbb{R}^{\mathbf{N}_V}} (f(x_U, x_V))^p \eta^p(x_V) dx_V \right\}^{\frac{1}{p}}, \quad 1 < p \leq q < \infty. \end{aligned} \quad (7. 13)$$

Proof: Observe that (7. 10)-(7. 12) imply the constraints in (2. 2)-(2. 4) with $\gamma, \delta, \alpha, n, \mathbf{N}$ replaced by $\gamma + \delta, 0, \alpha_V, |\mathbf{V}|, \mathbf{N}_V$ respectively.

Let $|\mathbf{V}| = 1$. **Theorem A** by Stein and Weiss [3] shows that these constraints are sufficient conditions to imply (7. 13).

Consider $|\mathbf{V}| \geq 2$. By applying **Principal Lemma**, $\rho(x_V) = |x_V|^{-(\gamma+\delta)}, \eta(x_V) \equiv 1$ satisfy the estimate in (5. 2)-(5. 3) for every $\mathbf{Q}_V \subset \mathbb{R}^{\mathbf{N}_V}$ whereas $\alpha_i > \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right)$, $i \in \mathbf{V}$.

Denote \mathbf{t} as a $|\mathbf{V}|$ -tuple $(2^{-t_1}, 2^{-t_2}, \dots, 2^{-t_{|\mathbf{V}|}})$. We find

$$\sum_{\mathbf{t}} \mathbf{A}_{pqr}^{\alpha_V}(\mathbf{t} : |x_V|^{-(\gamma+\delta)}, 1) < \infty \quad (7. 14)$$

for some $r = r(p, q, \alpha, \gamma, \delta) > 1$. Again, this is the desired summability in (3. 14). \square

Proposition 7.3 Let $\omega(x_U) = \sigma(x_U) = |x_U|^\delta$. For a.e $x_V \in \mathbb{R}^{\mathbf{N}_V}$, we have

$$\begin{aligned} & \left\{ \int_{\mathbb{R}^{\mathbf{N}_U}} \left\{ \int_{\mathbb{R}^{\mathbf{N}_U}} f(y_U, x_V) \prod_{i \in \mathbf{U}} \left[\frac{1}{|x_i - y_i|} \right]^{\mathbf{N}_i - \alpha_i} dy_U \right\}^q \omega^q(x_U) dx_U \right\}^{\frac{1}{q}} \\ & \leq \mathfrak{B}_{p,q,\alpha,\gamma,\delta} \left\{ \int_{\mathbb{R}^{\mathbf{N}_U}} (f(x_U, x_V))^p \omega^p(x_U) dx_U \right\}^{\frac{1}{p}}, \quad 1 < p < q < \infty. \end{aligned} \quad (7. 15)$$

Proof: From (7. 1) and (7. 11), we find

$$-\delta + \delta = 0 = \sum_{i \in \mathbf{U}} \alpha_i - \mathbf{N}_i \left(\frac{1}{p} - \frac{1}{q} \right), \quad -\delta < \frac{\mathbf{N}_i}{p} - \alpha = \frac{\mathbf{N}_i}{q} \quad \text{for } i \in \mathbf{U}. \quad (7. 16)$$

These constraints are sufficient conditions for **Theorem A** on every subspace $\mathbb{R}^{\mathbf{N}_i}$ for $i \in \mathbf{U}$. The one-weight $\mathbf{L}^p \rightarrow \mathbf{L}^q$ -norm inequality in (7. 15) follows the iteration argument given in section 6. \square

Let $\rho(x_{\mathbf{V}}) = |x_{\mathbf{V}}|^{-(\gamma+\delta)}$ and $\sigma(x_{\mathbf{U}}) = |x_{\mathbf{U}}|^{\delta}$ where $\gamma + \delta \geq 0$ and $\delta \leq 0$. Observe that

$$\omega(x) = |x|^{-\gamma} \leq \rho(x_{\mathbf{V}}) \sigma(x_{\mathbf{U}}). \quad (7. 17)$$

We have

$$\begin{aligned} & \left\{ \int_{\mathbb{R}^{\mathbf{N}}} \left\{ \int_{|y_{\mathbf{U}}| > |y_{\mathbf{V}}|} f(y) \prod_{i=1}^n \left[\frac{1}{|x_i - y_i|} \right]^{\mathbf{N}_i - \alpha_i} dy \right\}^q \omega^q(x) dx \right\}^{\frac{1}{q}} \\ & \leq \left\{ \int_{\mathbb{R}^{\mathbf{N}}} \left\{ \int_{|y_{\mathbf{U}}| > |y_{\mathbf{V}}|} f(y) \prod_{i=1}^n \left[\frac{1}{|x_i - y_i|} \right]^{\mathbf{N}_i - \alpha_i} dy \right\}^q \rho^q(x_{\mathbf{V}}) \sigma^q(x_{\mathbf{U}}) dx \right\}^{\frac{1}{q}} \quad \text{by (7. 17)} \\ & = \left\{ \iint_{\mathbb{R}^{\mathbf{N}_{\mathbf{U}}} \times \mathbb{R}^{\mathbf{N}_{\mathbf{V}}}} \left\{ \iint_{|y_{\mathbf{U}}| > |y_{\mathbf{V}}|} f(y_{\mathbf{U}}, y_{\mathbf{V}}) \prod_{i \in \mathbf{U} \cup \mathbf{V}} \left[\frac{1}{|x_i - y_i|} \right]^{\mathbf{N}_i - \alpha_i} dy_{\mathbf{U}} dy_{\mathbf{V}} \right\}^q \rho^q(x_{\mathbf{V}}) \sigma^q(x_{\mathbf{U}}) dx_{\mathbf{U}} dx_{\mathbf{V}} \right\}^{\frac{1}{q}} \\ & \leq \mathfrak{B}_{p \ q \ \alpha \ \gamma \ \delta} \left\{ \int_{\mathbb{R}^{\mathbf{N}_{\mathbf{U}}}} \left\{ \int_{\mathbb{R}^{\mathbf{N}_{\mathbf{V}}}} \left\{ \int_{|y_{\mathbf{U}}| > |x_{\mathbf{V}}|} f(y_{\mathbf{U}}, x_{\mathbf{V}}) \prod_{i \in \mathbf{U}} \left[\frac{1}{|x_i - y_i|} \right]^{\mathbf{N}_i - \alpha_i} dy_{\mathbf{U}} \right\}^p dx_{\mathbf{V}} \right\}^{\frac{q}{p}} \sigma^q(x_{\mathbf{U}}) dx_{\mathbf{U}} \right\}^{\frac{1}{q}} \\ & \quad \text{by Proposition 7.2} \\ & \leq \mathfrak{B}_{p \ q \ \alpha \ \gamma \ \delta} \left\{ \int_{\mathbb{R}^{\mathbf{N}_{\mathbf{V}}}} \left\{ \int_{\mathbb{R}^{\mathbf{N}_{\mathbf{U}}}} \left\{ \int_{|y_{\mathbf{U}}| > |x_{\mathbf{V}}|} f(y_{\mathbf{U}}, x_{\mathbf{V}}) \prod_{i \in \mathbf{U}} \left[\frac{1}{|x_i - y_i|} \right]^{\mathbf{N}_i - \alpha_i} dy_{\mathbf{U}} \right\}^q \sigma^q(x_{\mathbf{U}}) dx_{\mathbf{U}} \right\}^{\frac{p}{q}} dx_{\mathbf{V}} \right\}^{\frac{1}{p}} \\ & \quad \text{by Minkowski integral inequality} \\ & \leq \mathfrak{B}_{p \ q \ \alpha \ \gamma \ \delta} \left\{ \iint_{|x_{\mathbf{U}}| > |x_{\mathbf{V}}|} (f(x_{\mathbf{U}}, x_{\mathbf{V}}))^p \sigma^p(x_{\mathbf{U}}) dx_{\mathbf{U}} dx_{\mathbf{V}} \right\}^{\frac{1}{p}} \quad \text{by Proposition 7.3} \\ & \leq \mathfrak{B}_{p \ q \ \alpha \ \gamma \ \delta} \left\{ \int_{\mathbb{R}^{\mathbf{N}}} (f\sigma)^p(x) dx \right\}^{\frac{1}{p}}, \quad 1 < p < q < \infty. \end{aligned} \quad (7. 18)$$

The last inequality holds because $\sigma(x_{\mathbf{U}}) \approx \sigma(x)$ for $|x_{\mathbf{U}}| > |x_{\mathbf{V}}|$.

Acknowledgement:

I am deeply grateful to my advisor Elias M. Stein for those stimulating talks and unforgettable lectures.

References

- [1] G. H. Hardy and J. E. Littlewood, *Some Properties of Fractional Integrals*, Mathematische Zeitschrift **27**: no.1, 565-606, 1928.
- [2] S. L. Sobolev, *On a Theorem of Functional Analysis*, Matematicheskii Sbornik **46**: 471-497, 1938.
- [3] E. M. Stein and G. Weiss, *Fractional Integrals on n -Dimensional Euclidean Space*, Journal of Mathematics and Mechanics **7**: 503-514, 1958.
- [4] E. M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality and Oscillatory Integrals*, Princeton University Press, 1993.
- [5] B. Muckenhoupt and R. L. Wheeden, *Weighted Norm Inequality for Fractional Integrals*, Transactions of the American Mathematical Society **192**: 261-274, 1974.
- [6] E. T. Sawyer and R. L. Wheeden, *Weighted Inequalities for Fractional Integrals on Euclidean and Homogeneous Spaces*, American Journal of Mathematics **114**: no. 4, 813-874, 1992.
- [7] E. T. Sawyer and Z. Wang, *The Product Stein-Weiss Theorem*, Studia Mathematica **256**: 259-309, 2021.
- [8] C. Fefferman and B. Muckenhoupt, *Two Nonequivalent Conditions for Weight Functions*, Proceedings of the American Mathematical Society **45**: 99-104, 1974.
- [9] R. R. Coifman and C. Fefferman, *Weighted Norm Inequalities for Maximal Functions and Singular Integrals*, Studia Mathematica **51**: 241-250, 1974.
- [10] C. Perez, *Two Weighted Norm Inequalities for Riesz Potentials and Uniform L^p -Weighted Sobolev Inequalities*, Indiana University Mathematics Journal **39**: no.1, 31-44, 1990.
- [11] A. Cordoba and R. Fefferman, *A geometric Proof of the Strong Maximal Theorem*, Annals of Mathematics **102**: no.1, 95-100, 1975.
- [12] R. Fefferman, *Harmonic Analysis on Product Spaces*, Annals of Mathematics **126**: no.1, 109-130, 1987.
- [13] R. Fefferman and E. M. Stein, *Singular Integrals on Product Spaces*, Advances in Mathematics **45**: no.2, 117-143, 1982.
- [14] D. Müller, F. Ricci, E. M. Stein, *Marcinkiewicz Multipliers and Multi-parameter structures on Heisenberg (-type) group, I*, Inventiones Mathematicae **119**: no.2, 199-233, 1995.
- [15] J. L. Journé, *Calderón-Zygmund Operators on Product Spaces*, Revista Matemática Iberoamericana **1**: no.3, 55-91, 1985.
- [16] J. Pipher, *Journé's Covering Lemma and Its Extension to Higher Dimensions*, Duke Mathematics Journal **53**: no.3, 683-690, 1986.

Westlake University, Hangzhou, 310010, China
email: wangzipeng@westlake.edu.cn