



Dynamic Guideline: WHO-Based SARS - Diagnosis and/or Treatment and/or Management

I. WHO Latest Guideline Summary (Summary Date: 29 April 2026)

Scope and critical clarification

This report addresses **severe acute respiratory syndrome caused by SARS-CoV-1**, not SARS-CoV-2/COVID-19. WHO describes SARS as a **viral respiratory disease caused by a SARS-associated coronavirus** [WHO SARS topic page](#).

The most recent directly relevant and verifiable WHO SARS guideline identified is:

WHO guidelines for the global surveillance of severe acute respiratory syndrome (SARS): updated recommendations, October 2004

WHO states that this document provides **revised global surveillance and reporting guidance** for SARS as an ongoing strategy for **rapid case detection** [WHO publication page](#). The WHO IRIS PDF indicates that the October 2004 guidance **replaces previous WHO SARS surveillance guidance** [WHO IRIS PDF](#).

A separate WHO publication titled “**Meaningful youth engagement in a WHO guideline development process: case study of WHO abortion care guideline (2022)**” was identified, but it is **not a SARS guideline** and is not linked to SARS or infectious disease guideline development in the retrieved evidence [WHO publication page](#), [WHO event page](#).

1.1 Guideline Overview

Item	Details
Guideline Title	WHO guidelines for the global surveillance of severe acute respiratory syndrome (SARS): updated recommendations, October 2004
Publication Date	15 October 2004 / October 2004
Latest SARS-specific WHO guideline status as of	29 April 2026
Target Population	National health authorities, surveillance systems, public-health authorities, laboratories, health-care workers, infection-prevention teams, and outbreak-response authorities
Disease/ Condition	Severe acute respiratory syndrome caused by SARS-CoV-1
Scope	Global SARS surveillance, case detection, reporting, laboratory-supported surveillance, preparedness, and public-health response
Objectives	Rapidly detect SARS cases; guide national and international reporting; prevent national and international spread; support sustainable surveillance without overburdening health-system capacity
Authoritative Source	WHO publication page and WHO IRIS record WHO publication page , WHO IRIS item

1.1.1 Exactly Five WHO SARS-Related Guideline/Document Records Screened

The following table lists **exactly five WHO SARS-related guideline/document records or categories** identified in the supplied evidence. Only the first is fully verifiable from the provided WHO/IRIS sources; the remaining four were named in the supplied search summaries but lacked complete verifiable WHO URLs, titles, and recommendation text.

#	WHO SARS-related item	Date	Verification status	Scope / relevance	Source
1	WHO guidelines for the global surveillance of severe acute respiratory syndrome (SARS): updated recommendations, October 2004	15 October 2004	Fully verifiable	Global surveillance, reporting, rapid detection, public-health response	WHO publication page , WHO IRIS PDF
2	WHO SARS Laboratory Safety guidance/document category	2003	Named but not fully verifiable from supplied sources	Laboratory biosafety and SARS testing context; full recommendations not available in supplied evidence	WHO IRIS PDF
3	WHO SARS Clinical Management guidance/document category	2003	Named but not fully verifiable from supplied sources	Clinical care and treatment context; full WHO text not available in supplied evidence	Not fully verifiable in supplied sources

#	WHO SARS-related item	Date	Verification status	Scope / relevance	Source
4	WHO SARS Infection Control guidance/document category	2003	Named but not fully verifiable as a separate document from supplied sources	Infection prevention and control; risk-based infection-control measures are verifiable within the 2004 surveillance PDF	WHO IRIS PDF
5	WHO SARS Epidemiology guidance/document category	2003	Named but not fully verifiable from supplied sources	Epidemiologic description and surveillance context; separate WHO document not fully verifiable in supplied evidence	WHO SARS topic page

1.2 Key Recommendations

Because the 2004 WHO SARS guidance predates modern WHO GRADE presentation, the following ratings are **retrospective GRADE judgments** based on the evidence described in the retrieved sources. For outbreak-control recommendations, evidence is largely observational but biologically plausible, consistent, and supported by the observed interruption of SARS transmission.

Recommendation	GRADE Evidence Quality	Strength
Maintain ongoing global SARS surveillance capable of	Moderate	Strong

Recommendation	GRADE Evidence Quality	Strength
rapid detection and reporting of suspected re-emergence.		
Use clinical and epidemiological case definitions , supplemented by laboratory testing where appropriate, for SARS surveillance.	Low	Strong
Report SARS signals rapidly through national public-health channels and, when criteria are met, to WHO/international systems, including case counts, deaths, geographic distribution, and local transmission areas.	Low	Strong
Implement immediate isolation of suspected/probable SARS cases and apply standard, droplet, and contact precautions in health-care settings.	Moderate	Strong
Use contact tracing and quarantine/monitoring of close contacts when SARS transmission is suspected or confirmed.	Moderate	Strong
Health-care workers should apply risk-based infection-control measures during care of patients with suspected or confirmed SARS to prevent or interrupt transmission.	Moderate	Strong
Provide supportive clinical care as the foundation of SARS treatment; manage hypoxemia, pneumonia, and acute respiratory distress syndrome according to standard critical-care principles.	Low	Strong
Do not use ribavirin, corticosteroids, lopinavir/ritonavir, interferons, IVIG, or convalescent plasma as routine SARS therapy outside research or exceptional outbreak-response protocols, because efficacy is unproven.	Very Low	Conditional
During inter-epidemic periods, avoid indiscriminate SARS laboratory testing when pre-test probability is extremely low,	Low	Conditional

Recommendation	GRADE Evidence Quality	Strength
because false positives may trigger unnecessary public-health actions.		
Establish preparedness systems: command-and-control structures, reporting pathways, laboratory referral plans, infection-control capacity, and communication systems before re-emergence occurs.	Low	Strong

1.3 Guideline Development Methodology

Evidence Review Process

The October 2004 WHO SARS guidance was developed after the 2003 SARS outbreak and superseded earlier WHO SARS surveillance and response guidance [WHO IRIS PDF](#). WHO describes the document as revised global surveillance and reporting guidance intended to rapidly detect SARS cases [WHO publication page](#).

The retrieved evidence indicates that SARS case definitions were based on **clinical and epidemiological data** and supplemented by laboratory testing [WHO case definitions page](#).

Consensus Method

The supplied sources do not provide full details of the WHO consensus process used for the 2004 SARS surveillance guideline. However, the document was issued as WHO global guidance and replaced prior WHO SARS surveillance guidance [WHO IRIS PDF](#).

Conflict of Interest Management

The supplied sources do not provide a detailed conflict-of-interest declaration process for the 2004 SARS guideline. This is an important limitation because current WHO guideline-development standards generally include formal management of competing interests, but the retrieved SARS-specific documents do not show a complete modern GRADE/COI framework.

Meaningful Youth Engagement

No retrieved evidence indicates that youth were engaged in development of the WHO SARS surveillance guideline. WHO's later evidence brief on meaningful youth engagement concerns the **WHO abortion care guideline (2022)**, not SARS [WHO publication page](#). Related UN guidance supports meaningful youth engagement in epidemics and pandemics, but this does not establish that such engagement occurred in the SARS guideline-development process [UNFPA MAYE guide](#).

II. Evidence Summary After WHO Latest Guideline Release (WHO guidelines for the global surveillance of severe acute respiratory syndrome (SARS): updated recommendations) (Evidence Cutoff Date: 29 April 2026)

2.1 New Evidence Overview

Search Strategy

The evidence synthesis focused on literature and authoritative sources published or indexed after the WHO October 2004 SARS surveillance guidance, with priority given to:

- WHO and WHO IRIS sources;
- CDC and other public-health sources;
- peer-reviewed reviews and systematic reviews;
- SARS-CoV-1 treatment, diagnosis, surveillance, infection prevention, and preparedness literature;
- clinical-trial registry evidence where available.

SARS-CoV-2/COVID-19 evidence was excluded unless clearly used only to distinguish it from original SARS-CoV-1 evidence.

Databases and Sources Searched

Sources represented in the supplied evidence include:

- WHO publication pages and WHO IRIS;
- WHO SARS topic page;
- CDC archived and infection-control materials;
- NCBI Bookshelf;
- PubMed Central;
- peer-reviewed systematic reviews;
- [ClinicalTrials.gov](https://clinicaltrials.gov) and registry analyses, where retrieved.

Inclusion Criteria

Included evidence met one or more of the following criteria:

1. Directly addressed **SARS-CoV-1** diagnosis, surveillance, treatment, management, infection prevention, or preparedness.
2. Summarized WHO SARS outbreak response or reporting mechanisms.
3. Provided systematic-review evidence on SARS treatment.
4. Described SARS-specific clinical or infection-control evidence.
5. Evaluated SARS-CoV-1 therapeutic reports or trial evidence.

Exclusion Criteria

Excluded evidence included:

- SARS-CoV-2/COVID-19 treatment guidelines, except where clearly identified as non-applicable context;
 - COVID-19 clinical trials not involving SARS-CoV-1;
 - non-WHO youth-engagement evidence not linked to SARS guideline development, except as contextual ethical/public-health material.
-

2.2 Key New Studies

Study / Source	Design	Key Findings	GRADE Quality
Stockman et al., SARS treatment systematic review	Systematic review of SARS treatment studies; searched MEDLINE, EMBASE, BIOSIS, and CENTRAL through February 2005	Included 54 SARS treatment studies, 15 in vitro studies, and 3 ARDS studies. Could not determine whether ribavirin, corticosteroids, lopinavir/ritonavir, interferon, IVIG, or convalescent plasma were effective.	Very Low
Annual Reviews SARS review	Narrative expert review	SARS-CoV was identified as the cause of SARS; WHO announced the definitive cause on 16 April 2003. Supportive care remained the only therapy with evidence-based support; public-health measures included vigilance, traveler screening, isolation, and quarantine.	Low
Clinical Medicine SARS review	Narrative clinical review	SARS resembled other respiratory infections; many patients deteriorated toward the end of the second week. Nosocomial transmission was a major feature, making infection control central to management.	Low
WHO/NCBI Bookshelf summary of global SARS	Public-health response summary	WHO reporting included daily country summaries with new cases, new deaths, cumulative probable cases, geographic distribution, and areas of	Low

Study / Source	Design	Key Findings	GRADE Quality
reporting		local transmission.	
SARS preparedness and laboratory-testing article	Preparedness review	In the absence of global SARS transmission, SARS testing should be limited to higher-risk clinical scenarios to avoid false positives and unnecessary public-health disruption.	Low
PMC SARS review on control measures and current status	Narrative review	WHO announced the outbreak over in July 2003; after laboratory-associated infections, no SARS infections were reported worldwide since early 2004. Prompt detection, isolation, standard/ droplet precautions, and contact tracing reduced transmission.	Low
Potential specific therapies review summarizing SARS-CoV-1 reports	Review of antiviral/ immunologic therapies	SARS-CoV-1 evidence for convalescent plasma, lopinavir/ ritonavir plus ribavirin, interferon plus corticosteroids, and steroids was non-randomized, retrospective, or uncontrolled.	Very Low
Clinical-trial registry scan	Registry search summary	No verifiable SARS-CoV-1-specific registered trial record was identified for ribavirin, corticosteroids, interferons, lopinavir/ritonavir, IVIG, convalescent plasma, monoclonal antibodies, vaccines, or diagnostics.	Very Low / Not applicable

Sources: [SARS treatment systematic review](#), [Annual Reviews](#), [Clinical Medicine review via PMC](#), [NCBI Bookshelf](#), [PMC preparedness article](#), [PMC SARS review](#), [Potential specific therapies in COVID-19](#).

2.2.1 Diagnosis and Case Detection

SARS clinical diagnosis is challenging because illness can resemble other acute respiratory infections [Clinical Medicine review via PMC](#). WHO surveillance case definitions were based on clinical and epidemiological data and supplemented by laboratory tests [WHO case definitions page](#).

Updated diagnostic interpretation as of 29 April 2026:

- In the absence of known SARS-CoV-1 circulation, diagnosis should require a high index of suspicion plus epidemiologic risk.
- Laboratory testing should be coordinated through public-health authorities.
- Testing low-risk patients indiscriminately may cause false positives and unnecessary outbreak responses [PMC preparedness article](#).

GRADE Table: Diagnosis

Evidence statement	Quality of Evidence	Strength of Recommendation
Use clinical-epidemiologic case definitions as the first step in SARS detection.	Low	Strong
Supplement case definitions with laboratory testing when epidemiologic risk or compatible severe respiratory disease exists.	Low	Strong
Avoid broad low-pretest-probability SARS testing during inter-epidemic periods.	Low	Conditional

2.2.2 Treatment and Clinical Management

No SARS-specific therapy has been proven effective. A systematic review found that available studies could not determine whether ribavirin, corticosteroids, lopinavir/ritonavir, interferon,

IVIg, or convalescent plasma were effective [SARS treatment systematic review](#).

Supportive care remains the only evidence-supported treatment approach identified in the retrieved SARS-specific evidence [Annual Reviews](#).

GRADE Table: Treatment

Intervention / Management Strategy	Evidence Summary	Quality of Evidence	Strength of Recommendation
Supportive care, oxygen therapy, and critical-care management for respiratory failure/ ARDS	No proven antiviral therapy; standard supportive management remains foundational.	Low	Strong
Ribavirin	Evidence inconclusive; potential toxicity; no proven benefit.	Very Low	Conditional against routine use
Corticosteroids	Evidence inconclusive; adverse effects reported; may be considered only for standard non-SARS indications or research protocols.	Very Low	Conditional against routine SARS-specific use
Lopinavir/ritonavir plus ribavirin	Historically controlled reports suggested possible benefit, but no definitive evidence.	Very Low	Conditional against routine use
Interferons	Small uncontrolled reports; no reliable efficacy conclusion.	Very Low	Conditional against routine use
IVIg	No reliable SARS-CoV-1 efficacy evidence identified.	Very Low	Conditional against routine

Intervention / Management Strategy	Evidence Summary	Quality of Evidence	Strength of Recommendation
			use
Convalescent plasma	Non-randomized reports suggested possible association with improved outcomes when given early, but evidence is highly uncertain.	Very Low	Conditional; use only in research or exceptional outbreak protocols

2.2.3 Infection Prevention and Control

Nosocomial transmission was a major feature of SARS [Clinical Medicine review via PMC](#). Infection prevention therefore remains central to all SARS management.

Evidence supports the following measures:

- rapid case detection;
- patient isolation;
- standard precautions;
- droplet precautions;
- contact precautions;
- contact tracing;
- quarantine or monitoring of close contacts;
- risk-based health-care worker protection.

A SARS review states that prompt case detection, isolation, standard and droplet precautions, and contact tracing reduced transmission [PMC SARS review](#). CDC infection-control references include SARS literature on emergency-room outbreaks, health-care worker protection, hospital surface contamination, airborne detection, and evidence of airborne transmission [CDC infection-control references](#).

GRADE Table: Infection Prevention and Control

Recommendation	Evidence Basis	Quality of Evidence	Strength
Immediately isolate suspected or probable SARS cases.	Outbreak observational evidence and consistent transmission-control experience.	Moderate	Strong
Use standard, droplet, and contact precautions for suspected or confirmed SARS.	Nosocomial transmission evidence and infection-control plausibility.	Moderate	Strong
Use airborne precautions for aerosol-generating procedures or high-risk clinical settings.	Evidence of airborne detection/transmission risk and precautionary principle.	Low	Strong
Perform contact tracing and quarantine/monitor close contacts.	SARS outbreak containment experience.	Moderate	Strong
Protect health-care workers with risk-based infection-control measures.	Health-care-associated transmission was prominent in SARS.	Moderate	Strong

2.2.4 Surveillance, Reporting, and Preparedness

WHO's SARS reporting during the outbreak relied on daily country summaries including:

- new cases;
- new deaths;
- cumulative probable cases;
- geographic distribution;
- areas of local transmission.

These reports were submitted by national public-health authorities to WHO Geneva, with copies to WHO country and regional offices [NCBI Bookshelf](#).

Preparedness measures identified in related guidance include command-and-control systems, surveillance, case reporting, contact surveillance, fever clinics, laboratory pathways, and communication mechanisms [CDC archive PDF](#), [PMC preparedness article](#).

GRADE Table: Surveillance and Preparedness

Recommendation	Evidence Basis	Quality of Evidence	Strength
Maintain capacity for rapid SARS case detection and public-health notification.	WHO surveillance strategy and outbreak-response experience.	Moderate	Strong
Use phased preparedness plans that can escalate from inter-epidemic vigilance to active outbreak response.	WHO surveillance framework and public-health preparedness principles.	Low	Strong
Maintain laboratory referral networks and biosafety capacity.	Need for rapid confirmation and avoidance of laboratory-associated infection.	Low	Strong
Maintain public communication systems to reduce misinformation and panic.	Preparedness literature and outbreak-response experience.	Low	Strong

2.3 Evidence Gaps and Future Research Needs

1. No proven SARS-specific antiviral therapy

- Existing treatment evidence is observational, uncontrolled, or historically controlled.
- Future SARS-like outbreaks require rapid adaptive randomized trials.

2. No robust SARS-CoV-1 clinical-trial registry evidence

- No verifiable SARS-CoV-1-specific registered trials were identified for ribavirin, corticosteroids, interferons, lopinavir/ritonavir, IVIG, convalescent plasma, monoclonal antibodies, vaccines, or diagnostics.

3. Limited modern diagnostic validation for inter-epidemic SARS

- Testing algorithms need validation for low-prevalence settings to minimize false positives.

4. Incomplete WHO documentation transparency by modern standards

- The 2004 guideline predates current GRADE and WHO evidence-to-decision methods.
- Conflict-of-interest and stakeholder-engagement processes are not fully visible in retrieved sources.

5. Youth and community engagement not documented in SARS guideline development

- WHO's meaningful youth-engagement evidence brief concerns abortion care, not SARS [WHO publication page](#).
- Future outbreak guidelines should include affected communities, health-care workers, youth representatives where relevant, and marginalized populations.

6. Preparedness for laboratory-associated and zoonotic re-emergence

- Reviews note that after the 2003 outbreak, a few laboratory-associated infections occurred, but no SARS infections were reported worldwide since early 2004 [PMC SARS review](#).

2.4 Updated Recommendations Based on New Evidence

Original Recommendation	New Evidence Impact	Updated Recommendation
Maintain global SARS surveillance for rapid detection.	No post-2004 evidence overturns this; SARS has not circulated globally since early 2004, but re-emergence remains theoretically possible.	Continue targeted, sustainable SARS surveillance integrated with severe acute respiratory infection and unusual pneumonia reporting.

Original Recommendation	New Evidence Impact	Updated Recommendation
Use case definitions based on clinical and epidemiologic features, supplemented by laboratory testing.	Later evidence reinforces that low-prevalence testing can cause false positives and unnecessary response burden.	Use clinical-epidemiologic screening first; reserve SARS-CoV-1 testing for compatible severe illness with epidemiologic risk, clusters, or public-health concern.
Report cases rapidly to WHO and public-health authorities.	WHO outbreak reporting experience supports rapid daily reporting during active transmission.	If SARS is suspected or confirmed, notify public-health authorities immediately and escalate to international reporting according to WHO/IHR mechanisms.
Implement isolation and infection-control measures.	Later reviews confirm nosocomial transmission was central to SARS spread.	Immediately isolate suspected/probable cases and use standard, droplet, contact, and risk-based airborne precautions, especially during aerosol-generating procedures.
Use contact tracing and quarantine.	SARS outbreak containment experience continues to support this approach.	Trace, risk-stratify, and monitor/quarantine close contacts according to exposure risk and public-health authority guidance.
Provide clinical management.	Systematic review evidence shows no proven specific SARS therapy.	Provide supportive care; do not use ribavirin, steroids, lopinavir/ritonavir, interferons, IVIG, or convalescent plasma routinely outside research or exceptional outbreak protocols.
Maintain preparedness.	Preparedness literature emphasizes command systems, surveillance,	Maintain SARS re-emergence preparedness as part of broader high-consequence respiratory

Original Recommendation	New Evidence Impact	Updated Recommendation
	reporting, lab pathways, and communication.	pathogen planning.
Stakeholder engagement not documented.	WHO's later youth-engagement guidance shows value of inclusive participation but is not SARS-specific.	Future SARS or SARS-like guidelines should use transparent WHO guideline methods, GRADE, conflict-of-interest management, and meaningful engagement of affected groups.

Final Clinical and Public-Health Position as of 29 April 2026

1. **Latest verifiable WHO SARS guideline:** WHO's October 2004 global SARS surveillance updated recommendations remain the latest directly relevant WHO SARS guideline identified [WHO publication page](#).
2. **Diagnosis:** Use clinical and epidemiological assessment first; add laboratory testing when epidemiologic risk or compatible severe respiratory illness justifies testing.
3. **Treatment:** No SARS-specific drug therapy has proven efficacy. Supportive care remains the standard.
4. **Infection control:** Isolation, standard/droplet/contact precautions, health-care worker protection, contact tracing, and quarantine remain the core SARS-control measures.
5. **Surveillance:** Maintain sustainable surveillance and preparedness for unusual pneumonia clusters or suspected re-emergence.
6. **Research priority:** Any future SARS-CoV-1 outbreak should trigger rapid, ethically approved, adaptive randomized trials and transparent WHO guideline updating using GRADE methods.