

Gram-Schmidt orthogonalization

- Gram-Schmidt method
- Examples
- Orthogonal matrices
- QR decomposition
- Least squares and Gram-Schmidt method

Suppose you have linearly independent vectors v_1, v_2, \dots, v_n . How to find an orthonormal basis for $\text{span}\{v_1, v_2, \dots, v_n\}$?

Step 1. Pick any vector, say v_1 , and make it of length 1:

$$e_1 = v_1/|v_1|.$$

Step 2. Pick any of the remaining vectors, say v_2 , make it orthogonal to v_1 :

$$v'_2 = v_2 - P_{e_1}(v_2),$$

where P_{v_1} is the projection operator on vector e_1 . Hence

$$v'_2 = v_2 - \langle v_2, e_1 \rangle e_1 = v_2 - \langle v_2, v_1 \rangle v_1 / \langle v_1, v_1 \rangle$$

Step 3. make v'_2 of length 1:

$$e_2 = v'_2/|v'_2|$$

Inductive step:

$$v'_k = v_k - \sum_{i < k} \langle v_k, v_i \rangle v_i / \langle v_i, v_i \rangle$$

$$e_k = v'_k/|v'_k|.$$

Question: Is it possible to do the same when you have $v_1, v_2, \dots, v_n, \dots$?

Example 1. Legendre polynomials.

Consider polynomials of degree $n-1$ on $[-1, 1]$ with the L_2 inner product. Then polynomials with the leading coefficient 1

$$1, x, x^2 - 1/3, \dots$$

is an orthogonal basis. Note, that Legendre polynomials are not an orthonormal basis.

Question: Is it possible to make Legendre polynomials orthonormal?

Example 2. Chebyshev Polynomials. Consider the vector space of polynomials on the interval $[-1, 1]$ with the inner product

$$\langle g, h \rangle = \int_{-1}^1 g(x)h(x)/(1-x^2)^{1/2} dx$$

We want to construct an orthogonal basis $T_k, k = 0, 1, 2, 3, \dots$ for such polynomials, such that for every k T_k is a polynomial of degree at most k , $\langle T_0, T_0 \rangle = \pi$ $\langle T_k T_k \rangle = \pi/2, k > 0$:

$$T_0 = 1, T_1 = x, T_2 = 2x^2 - 1, T_{n+1} = 2xT_n - T_{n-1}.$$

Definition A square matrix is called orthogonal if its columns are orthonormal.

Lemma If Q is orthogonal, then $Q^{-1} = Q^t$

Question: Recall the matrix of the Fourier transform, is it orthogonal?

Example 1: Rotation matrix in \mathbb{R}^2

Question: Is a permutation matrix orthogonal?

Question: Is a projection matrix orthogonal?

Lemma Multiplication by an orthogonal matrix preserves inner product.

Question: What does it tell us about angles and lengths? Why?

Lemma The Gram-Schmidt method implies that

$$(v_1 \ v_2 \ \dots \ v_n) = (e_1 \ e_2 \ \dots \ e_n) \begin{pmatrix} \langle v_1, e_1 \rangle & \langle v_2, e_1 \rangle & \dots & \langle v_n, e_1 \rangle \\ 0 & \langle v_2, e_2 \rangle & \dots & \langle v_n, e_2 \rangle \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \langle v_n, e_n \rangle \end{pmatrix}$$

for linearly independent vectors v_1, v_2, \dots, v_n .

Corollary $A = QR$, if columns of A are linearly independent.

Question: Suppose $A = QR$, how does it simplify the least squares method?