



Importance of regional PM_{2.5} transport and precipitation washout in heavy air pollution in the Twain-Hu Basin over Central China: Observational analysis and WRF-Chem simulation

Weiyang Hu^a, Tianliang Zhao^{a,*}, Yongqing Bai^{b,*}, Shaofei Kong^c, Jie Xiong^b, Xiaoyun Sun^a, Qingjian Yang^{a,d}, Yao Gu^a, Huicheng Lu^a

^a Climate and Weather Disasters Collaborative Innovation Center, Key Laboratory for Aerosol-Cloud-Precipitation of China Meteorological Administration, PREMIC, Nanjing University of Information Science & Technology, Nanjing 210044, China

^b Institute of Heavy Rain, China Meteorological Administration, Wuhan 430205, China


^c Department of Atmospheric Sciences, School of Environmental Studies, China University of Geosciences (Wuhan), Wuhan 430074, China

^d Henan Meteorological Observatory, Zhengzhou 450003, China

HIGHLIGHTS

- A quasi 2-day regional PM_{2.5} transport from northern to central China
- Cold air invasion and terrain block facilitated air pollution over the THB.
- Regional transport to the THB contributed 70.5% to heavy PM_{2.5} pollution.
- Precipitation washout largely dispelling the THB's heavy PM_{2.5} pollution

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 19 August 2020

Received in revised form 9 November 2020

Accepted 9 November 2020

Available online 14 November 2020

Editor: Pingqing Fu

Keywords:

PM_{2.5} pollution

Twain-Hu Basin

Regional transport

Precipitation washout

WRF-Chem

ABSTRACT

With observational analysis and WRF-Chem simulation on a heavy air pollution event in January 2019 over the Twain-Hu Basin (THB) in Central China, this study characterized the regional transport of PM_{2.5} emitted from the North China Plain (NCP) to the THB region in Central China and quantitatively assessed the influence of the regional PM_{2.5} transport and precipitation washout on PM_{2.5} change in the wintertime heavy air pollution over the THB. It was found that the THB's heavy air pollution event was exacerbated by the strong northeasterly winds driving a quasi 2-day time lag of regional PM_{2.5} transport from the NCP to the THB. The multi-scale atmospheric circulations of cold air invasion influenced by East Asian winter monsoon and the terrain block of THB altered the structures of regional PM_{2.5} transport in deteriorating air quality to the THB. It was assessed for the THB region that the enhancing contribution of regional PM_{2.5} transport to the high air pollution level reached up to 70.5% in the heavy air pollution, and the precipitation washout could contribute the 55.3% PM_{2.5} removal to dissipating the PM_{2.5} pollution over the THB with frequent precipitation and wet environment, distinguishing from the dominance of wind-cleaning air pollution in the other regions in China.

© 2020 Elsevier B.V. All rights reserved.

* Corresponding authors.

E-mail addresses: tlzhao@nuist.edu.cn (T. Zhao), 2007byq@163.com (Y. Bai).

1. Introduction

China is confronted with air pollution of fine particulate matter $PM_{2.5}$ with aerodynamic diameters equal to or less than $2.5 \mu m$ in the ambient atmosphere over the recent decades (Guo et al., 2014; T. Wang et al., 2019; Zhao et al., 2017). Apart from posing a threat to human health (Chen et al., 2017; Liu et al., 2016; Pope and Dockery, 2006), severe $PM_{2.5}$ episodes can lead to lower atmospheric visibility (Tao et al., 2012), and long term effects of high $PM_{2.5}$ levels can impact climate changes (Kan et al., 2012; J. Li et al., 2017), which has become an urgent issue in environmental sciences (Yang et al., 2013; Ye et al., 2016).

Air pollution events occur in the stagnantly meteorological conditions with air temperature inversion, weak winds, and stable boundary layers in addition to high anthropogenic emissions of air pollutants (Y. Wang et al., 2019; Xu et al., 2016; H. Zhang et al., 2015; Zhang et al., 2018; Zhong et al., 2019). Furthermore, regional transport of air contaminants is also an important factor that worsens regional air pollution (Z. Li et al., 2017; Miao et al., 2017; L. Chen et al., 2019; Jiang et al., 2015). Governed by strong winds, air pollutants can be transported easily from the upstream source regions to the downwind receptor areas, which can result to the excessive levels of air pollutants including $PM_{2.5}$ at receptors in the downwind region (Wang et al., 2017; Wu et al., 2017; Miao et al., 2019). Regional transport of gaseous precursors of $PM_{2.5}$ could contribute up to 70% secondary particles for the severe haze pollution in Beijing (Hua et al., 2016). The regional transport of polluted air masses over the Central and Eastern China (CEC) was the main factor leading to the increases of wintertime $PM_{2.5}$ concentrations in Wuhan, which could contribute 53.4%–65.0% in the heavy air pollution events (Lu et al., 2017; Lu et al., 2019; Yu et al., 2020). The contribution of regional $PM_{2.5}$ transport was ranged from 29% to 60% to $PM_{2.5}$ pollution in East and South China (Q. Chen et al., 2019; Ji et al., 2018; Kang et al., 2019).

Determined by air pollutant emissions and meteorological conditions, air pollution events frequently occur in four regions over China including the North China Plain (NCP) (Han et al., 2018; Sun et al., 2016; Zhang et al., 2019), the Yangtze River Delta (YRD) (Huang et al., 2020) in East China, the Sichuan Basin (SCB) (Du et al., 2020; Ning et al., 2018) in Southwest China and the Pearl River Delta (PRD) (Huang et al., 2018) in South China. The Twain-Hu Basin (THB) with the lower lands (mainly less than 200 m a. s. l.) of two provinces Hubei and Hunan in Central China is a geographical junction linking NCP in the north, YRD in the east, PRD in the south and SCB in the west (Fig. 1). In recent years, the THB has been confronting severe air pollution (Bai et al., 2018; X.Y. Zhang et al., 2015) and high AOD values (>0.8) in winter based on 19 years of MODIS data (Shen et al., 2020a, 2020b). Importantly, the THB is situated at a crossroads of regional transport of air

pollutants over central and eastern China with the prevailing winds of East Asian monsoons. Previous studies of air pollutants' regional transport were mostly concentrated on the regions of NCP, YRD, SCB and PRD (Chang et al., 2018a, 2018b; Huang et al., 2020; Kang et al., 2019; X. Li et al., 2019; R. Li et al., 2019; Ma et al., 2017; Zhang et al., 2019). However, the structures of regional $PM_{2.5}$ transport with the contribution to the heavy air pollution event in the THB region have been poorly understood.

Wet scavenging is an important process of aerosol removals from the atmosphere, which are sorted as in-cloud and below-cloud scavenging processes (Seinfeld and Pandis, 2006). The in-cloud scavenging process happens when aerosols serve as cloud condensation nuclei (CCN) and ice nuclei (IN), or directly captured by cloud drops, while the below-cloud process is an aerosol washout by precipitating raindrops or snow particles (Chate, 2005). These two wet scavenging processes are regarded as crucial for changing aerosol concentrations in the atmosphere (Luan et al., 2019). Therefore, what extend of aerosol wet scavenging by precipitation in the $PM_{2.5}$ removal over the THB with frequent precipitation and wet environment needs a quantitative assessment.

In this study, we utilized the observed data of air pollutants and meteorology to investigate a heavy air pollution event in January 2019 over the THB associated with regional $PM_{2.5}$ transport from the source regions in Northern China. Furthermore, by employing the air quality model WRF-Chem, we simulated the structures of regional $PM_{2.5}$ transport to the THB and assessed the contribution of regional $PM_{2.5}$ transport and aerosol removal by precipitation washout to heavy air pollution event occurred in the THB. The objective of this study was to understand the importance of regional $PM_{2.5}$ transport and precipitation washout in air pollution over Central China with an implication for regional change of air quality in a downwind region with frequent precipitation and wet environment.

2. Data and methods

2.1. Data

Here, we used the hourly surface $PM_{2.5}$ concentrations over central-eastern China from the China Air Quality Online Monitoring and Analysis Platform (<http://www.mee.gov.cn>), the data of u- and v-wind components at 10 m from the hourly ERA5-Land data - the reanalysis dataset from ECMWF (<https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-land?tab=form>), and the near-surface meteorological data over the THB with a temporal resolution of 1 h from China Meteorological Administration (<http://data.cma.cn/>), including




Fig. 1. (a) The geographical positions of two provinces, Hubei and Hunan (blue areas), in Central China with the red frame covering the WRF-Chem simulation domain and the NCP (grey areas), PRD, YRD and SCB region; (b) the terrain height (m in a.s.l.) in the THB (white dashed line) with eight urban sites Xiangyang (XY), Jingmen (JM), Jingzhou (JZ), Changde (CD), Suizhou (SZ), Xiaogan (XG), Wuhan (WH) and Xianning (XN) and surrounding regions.

air temperature, relative humidity, sea level pressure, wind speed, wind direction and precipitation. Time we mentioned in this study was the local time in China (UTC + 8:00 h).

2.2. Model configuration

The Weather Research and Forecasting model with Chemistry (WRF-Chem) version 3.4.1 online coupling model (Grell et al., 2005) was employed to simulate a PM_{2.5} pollution event over the THB. The simulation domain (Fig. 1) was centred at 30.5° N and 114.06° E with 9 km × 9 km horizontal resolution. The vertical 38 hybrid layers were set from the surface to 50 hPa, the lowest 21 levels in the planetary boundary layer (PBL) to better reproduce the air pollution event. The NCEP Final Global Forecast System Operational Analysis (FNL) 6-hourly dataset with a horizontal resolution of 1° × 1° was selected to be the meteorologically initial driving fields and boundary conditions.

Nudging-based four-dimensional data assimilation (FDDA) was used to effectively reduce model errors with the nudging coefficient of 3.0×10^{-4} for wind, air temperature and water vapour mix ratio, and we selected spectral nudging matching 10 and 5 wavenumbers in zonal and meridional directions respectively.

The parameterization used in the model configuration for the WRF-Chem simulation experiment were summarized in Table 1.

The Multi-resolution Emission Inventory for China (MEIC) with a high spatial resolution (0.25° × 0.25°) (<http://www.meicmodel.org/>) from 2016 was applied for the anthropogenic air pollutant emissions.

A heavy air pollution event with surface PM_{2.5} concentrations over 150 µg m⁻³ was observed from January 4–9, 2019 in the THB region (Fig. S1). In order to simulate the heavy air pollution event, a period of WRF-Chem modelling was chosen from 00:00 a.m. of January 2 to 07:00 a.m. of January 10, 2019, with the spin-up time of the first 36 h.

2.3. Numerical experiments

In addition to the control experiment (CE), we designed two sensitivity experiments TE (transport experiment) with closing all anthropogenic emissions of air pollutants over the THB, and WE (washout experiment) with closing wet scavenging from precipitation to PM_{2.5} concentrations to evaluate the contribution of regional PM_{2.5} transport and precipitation wet removal to PM_{2.5} changes in the heavy air pollution event occurred over the THB. The detail information of the simulation experiments performed in this study is listed in Table 2.

2.4. Modelling validation

The CE modelling results of meteorology and PM_{2.5} concentrations were validated with the observational data in the THB (Figs. 2, S2 and Table 3). The statistical metrics, including correlation coefficient (R), mean bias (MB), the root mean square error (RMSE), the normalized mean bias (NMB), the mean fractional bias (MFB), and the mean fractional error (MFE), were used in the modelling validation. The CE

Table 1

Parameterization schemes used in the WRF-Chem model.

Options	WRF-Chem
Microphysics	Lin scheme
Longwave radiation	RRTMG scheme
Shortwave radiation	Goddard shortwave scheme
Surface layer	MM5 similarity surface layer
Land-surface	the unified Noah land surface model
Urban canopy model	Single-layer UCM scheme
Boundary layer	YSU boundary layer scheme
Cumulus	Grell 3-D ensemble scheme
Photolysis	Fast-J
Chemistry	CBMZ
Aerosol	MOSAIC (4 bins)
Spectral nudging	On

Table 2

The description of three simulation experiments in this study.

Experiments	Description
CE (control experiment)	the base case
TE (transport experiment)	CE with closing all anthropogenic emissions over the THB
WE (washout experiment)	CE with closing PM _{2.5} below-cloud washout from precipitation

simulated meteorology generally agreed well with the observations (Figs. 2, S2). Meanwhile, the simulated PM_{2.5} values were reasonably consistent with the observations (Table 3), meeting the good modelling performance proposed by Boylan and Russell (2006). Overall, the simulations of meteorology and PM_{2.5} predicted by WRF-Chem over the THB presented the good agreements with the observations. Therefore, the simulation data could be used to investigate the regional transport and precipitation washout of PM_{2.5} in the heavy air pollution over the THB.

3. Results and discussion

3.1. Observational analysis

3.1.1. Regional PM_{2.5} transport captured over CEC

To investigate the regional PM_{2.5} transport observed over central-eastern China (CEC) to the heavy air pollution happened in the THB, the observed PM_{2.5} concentrations over CEC with the near-surface wind vectors from the reanalysis meteorology of ERA5-Land were averaged from 03:00 p.m. of January 3 to 7:00 a.m. of January 10, 2019 (Fig. 3). During this period, the strong northeasterly winds prevailed over CEC (mainly in the NCP), driving the regional transport of PM_{2.5} pollutants sourced from the upwind northern CEC areas to the downwind THB region (Fig. 3). Strong winds are in favour of the removal of air pollutants for the local areas and air pollutant transport to the downwind areas (Wei et al., 2015; Yang et al., 2015).

To further investigate the regional PM_{2.5} transport from the upstream CEC region to the THB, we explored the hourly variations of PM_{2.5} concentrations over January 1–10, 2019 at four observational sites Handan, Zhengzhou, Nanyang, Xiangyang (Fig. 4). Driven by the strong northeasterly winds (Fig. 3), the surface PM_{2.5} peaks advanced southwards at 10 p.m. of January 3, from site 1 (Handan), at 12 p.m. of January 4 to site 2 (Zhengzhou), at 12 p.m. of January 5 to site 3 (Nanyang) and at 00 a.m. on January 6 to site 4 (Xiangyang), indicating a large impact of regional PM_{2.5} transport on enhancing PM_{2.5} concentrations to 392 µg m⁻³ for heavy air pollution in the THB with a quasi-2d time lag (Fig. 4).

3.1.2. Connection of PM_{2.5} changes with precipitation over the THB

In order to figure out the connection of changes of PM_{2.5} concentrations and precipitation washout during the heavy air pollution over the THB, we compared the hourly variations in regional averages of observed PM_{2.5} concentrations and precipitation over the THB (Fig. 5). As shown in Fig. 5, there were two periods of apparent precipitation with dropping PM_{2.5} concentrations down, and the changes of PM_{2.5} concentrations were estimated to have negative correlations with the R values of −0.57 and −0.44 respectively, passing the significant level of 0.05, reflecting the importance of precipitation in removal PM_{2.5} concentrations for dispelling the heavy PM_{2.5} pollution. The contribution of precipitation removal to heavy PM_{2.5} pollution in the THB was quantitatively assessed in Section 3.2.3, based on the modelling experiments.

3.2. WRF-Chem simulation

The control experiment and two sensitivity experiments with closing all anthropogenic emissions of air pollutants over the THB, and




Fig. 2. Hourly changes of observed (black dot lines) and simulated (red dots) meteorological elements including surface air temperature (T), relative humidity (RH) and sea level pressure (SLP) as well as near-surface wind speed (WS) and precipitation averaged over the THB from 03:00 p.m. of January 3 to 7:00 a.m. of January 10, 2019.

with closing $PM_{2.5}$ wet scavenging of precipitation were conducted in this study (Table 2). Based on the three WRF-Chem simulation experiments, we investigated the structures of regional $PM_{2.5}$ transport and also assessed the contribution of regional $PM_{2.5}$ transport and the precipitation wet removal to $PM_{2.5}$ changes to the heavy air pollution observed in the THB in the following sections.

3.2.1. The vertical structures of regional $PM_{2.5}$ transport

To characterize the vertical structures of regional $PM_{2.5}$ transport during the heavy air pollution, Fig. 6 presented the vertical sections (based the back dashed line as shown in Fig. 3) of $PM_{2.5}$ concentrations

and wind vectors along the prevailing northeasterly wind direction at 02:00 a.m. of January 2, 02:00 p.m. of January 4 and 5, 2019 based on the control experiment of WRF-Chem simulation. Accompanying a cold air invasion during East Asian winter monsoonal season (Fig.S1), the regional $PM_{2.5}$ transport from the northern CEC region (mainly the NCP) to the THB was generally constrained below 2.6 km in the lower troposphere, where the northerly winds in the lower layers and the southerly winds in the high level made up the vertical circulations in the regional $PM_{2.5}$ transport (Fig. 6). Meteorologically, an invasion of cold air in the northerly winds uplifts warm air in the southerly winds along the surface southward cold front with cold air subsidence behind the cold front (Ding, 1993), which was clearly reflected in the temporal evolution of vertical circulations in the lower troposphere (Fig. 6).

The near-surface $PM_{2.5}$ peaks in the upstream NCP could be emitted from the air pollutant sources there (Fig. 6a–b), the southward movements of the atmospheric column with high $PM_{2.5}$ concentrations along the regional transport were retarded, by blocks of the southerly winds and the windward basin terrain in the THB, and the distinct downward airflow in the regional $PM_{2.5}$ transport with the terrain block of THB was conducive to accumulate near-surface $PM_{2.5}$. Governed by the multi-scale atmospheric circulations, the external transport of $PM_{2.5}$ pollutants emitted from the source region can be transported easily to the downstream receptor areas which can exacerbate acceptor regions' air quality, resulting in a complicated relation of source and receptor in regional transport of air pollutants.

Table 3

Statistical metrics of the comparisons from hourly observed and simulated surface $PM_{2.5}$ concentrations at eight urban sites in the THB.

Indices	NMB (%)	MFB (%)	MFE (%)	RMSE ($\mu\text{g m}^{-3}$)	R
XY	-27.41	-34.58	36.75	81.55	0.85
JM	-23.70	-32.37	41.16	63.41	0.66
JZ	-9.88	-22.63	46.20	57.35	0.55
CD	-24.92	-33.41	44.69	71.12	0.69
SZ	-10.46	-10.73	18.65	35.23	0.86
XG	-6.40	-13.84	29.12	28.10	0.86
WH	-7.89	-10.95	22.46	30.47	0.85
XN	3.39	-4.04	33.31	31.91	0.74




Fig. 3. Spatial distribution of surface $PM_{2.5}$ concentration averages ($\mu g m^{-3}$) observed at the sites (coloured dots) and 10 m wind vectors of ERA5-Land from 03:00 p.m. of January 3 to 7:00 a.m. of January 10, 2019, over CEC with the black dash line and four observational sites (numbers with circles) along the direction of prevailing northeasterly winds during the heavy air pollution event.

3.2.2. Contribution of regional $PM_{2.5}$ transport to the heavy air pollution in the THB

In addition to the control experiment (CE), we designed the transport experiment (TE) with closing all anthropogenic emissions of air pollutants over the THB in CE (Table 2). As the TE-simulated $PM_{2.5}$




Fig. 4. Hourly changes of $PM_{2.5}$ concentrations observed in three upwind sites, Handan, Zhengzhou and Nanyang (sites 1, 2, 3 in Fig. 3) and the downstream THB's site Xiangyang (site 4 in Fig. 3) during 1–10 January 2019. The grey dash lines indicate the heavy air pollution level of $150 \mu g m^{-3}$ for surface $PM_{2.5}$ concentrations. The two arrows indicated the intervals of lag time along with the regional $PM_{2.5}$ transport from Handan to Zhengzhou, Nanyang and Xiangyang during the heavy air pollution episode.

concentrations were divided by those simulated with CE, the contribution of regional $PM_{2.5}$ transport from the upstream CEC to heavy air pollution in the THB was quantitatively estimated. The spatial distribution of the contribution for regional $PM_{2.5}$ transport presented the significantly decline gradients in the general north-south direction over the THB (Fig. 7). Governing by northeasterly winds, the regional transport of $PM_{2.5}$ pollutants dominated the surface $PM_{2.5}$ changes during the air pollution event with the contribution up to 80–85% in the northern THB, where the western areas were greater than the eastern areas (Fig. 7), demonstrating the effect of the large terrain gap in the western THB (Fig. 1b) on the regional $PM_{2.5}$ transport.

Furthermore, the regional $PM_{2.5}$ transport contribution was averaged over the THB region under conditions of light air pollution ($75 \mu g m^{-3} \leq PM_{2.5} < 150 \mu g m^{-3}$) and heavy air pollution ($PM_{2.5} \geq 150 \mu g m^{-3}$) respectively, as listed in Table 4. The contributions of regional $PM_{2.5}$ transport to THB were estimated with 58.9% and 70.5% during the light and heavy air pollution periods, respectively. The results provide an important information that regional $PM_{2.5}$ transport played an vital role in $PM_{2.5}$ sources over the THB.

3.2.3. Removal contribution of precipitation washout to surface $PM_{2.5}$ concentrations

As designed in the WRF-Chem modelling experiments (Table 2), the $PM_{2.5}$ washout experiment WE was with closing aerosol below-cloud washout from precipitation in the control experiment CE. Using the method of Feng and Wang (2012) to calculate the precipitation removal contributions, what extend of aerosol wet scavenging by precipitation in the removal of $PM_{2.5}$ over the THB could be quantitatively assessed from WE and CE. There were two apparent precipitation processes from 09:00 p.m. of January 3 to 01:00 p.m. of January 4 (the formation stage) and from 01:00 p.m. of January 8 to 09:00 p.m. of January 9 (the dissipating episodes), respectively (Fig. 5). Therefore, we evaluated the washout contributions of precipitation to surface $PM_{2.5}$ concentrations in the two apparent rainfall processes (Table 5). In the formation stage of heavy $PM_{2.5}$ pollution, the removal $PM_{2.5}$ contribution of precipitation washout removal contribution reached to 17.0%, which could be offset by the $PM_{2.5}$ accumulation during the evolution to heavy air pollution. In the dissipating stage of heavy air pollution, precipitation removal of $PM_{2.5}$ concentrations was accounted for 55.3%, exhibiting that the precipitation washout has a large influence on dispelling the heavy $PM_{2.5}$ pollution over the THB with frequent precipitation and wet environment, differing from the role wind-cleaning air pollution in the other areas in China.

The integrated source apportionment method in CMAQ model (CMAQ/ISAM) is a reliable method evaluating the local/regional contributions to the air pollution, while the sensitivity experiments of zero local emissions conducted by WRF-Chem is also a reasonable way to assess local and regional contributions in many simulation studies (L. Chen et al., 2019; Q. Chen et al., 2019; L. Zhang et al., 2015; Gao et al., 2011). In order to support the sensitivity test results of zero out the local emission to $PM_{2.5}$ levels, CMAQ/ISAM could assess the contribution of regional transport and local emissions to $PM_{2.5}$ concentrations in the THB with a comprehensive comparison with WRF-Chem sensitivity tests in further study.

4. Conclusions

In this paper, we studied a heavy air pollution event occurred in January 2019 over the THB in terms of regional $PM_{2.5}$ transport and precipitation washout by using meteorological and environmental observations, high spatial-resolution reanalysis data of meteorology and WRF-Chem simulations to explore the importance of regional $PM_{2.5}$ transport and precipitation washout in heavy air pollution over Central China.

The strong northeasterly winds prevailed over the NCP, driving the regional transport of $PM_{2.5}$ from the upwind northern NCP areas to




Fig. 5. Hourly changes of average $PM_{2.5}$ concentrations (red line) and precipitation (black triangles) observed over the THB with the light blue shades marking the periods of apparent precipitation.




Fig. 6. Vertical cross-sections of simulated $PM_{2.5}$ concentrations (colour contours; $\mu g m^{-3}$), wind vectors and terrain (black shades) along the prevailing wind direction (black dash line in Fig. 3) at (a) 02:00 a.m. of January 4, (b) 02:00 p.m. of January 4, and (c) 02:00 p.m. of January 5, 2019, during the heavy air pollution event. Note that the vertical speeds of wind vectors (arrows) are multiplied by 100 for the illustration of vertical circulations. The red frames denote the atmospheric columns over the THB.

the downwind THB region. A quasi 2-day time lag impact of regional $PM_{2.5}$ transport was found on peaking $PM_{2.5}$ concentration in the heavy pollution over the THB region. The multi-scale atmospheric

circulations of cold air invasion of East Asian winter monsoon and block of THB terrain could determine the structures of regional $PM_{2.5}$ transport in deteriorating air quality to the THB. The regional transport contribution displayed the obviously decline gradients from north to south over the THB with the regional average reaching to 70.5% during heavy air pollution stage. A strong influence of precipitation washout on dispelling the heavy $PM_{2.5}$ pollution over the THB, and the removal contribution of precipitation to $PM_{2.5}$ was accounted for 55.3%, which was different from the other air pollution regions with the dominant wind-cleaning air pollution.

Based on the investigation of a wintertime heavy air pollution in the THB region, Central China, this study revealed the regional $PM_{2.5}$ transport over CEC with large contributions of regional $PM_{2.5}$ transport and precipitation washout of aerosols to the THB's heavy air pollution. Further exploration can be done with the support of climatic analyses of multi-source data of long term observations and more accurate




Fig. 7. Spatial distribution of regional $PM_{2.5}$ transport contribution over the THB averaged from 03:00 p.m. of January 3 to 7:00 a.m. of January 10, 2019, with the THB region encircled by the white dash line.

Table 4

Contribution of regional transport and local emissions to surface $PM_{2.5}$ pollutants under different $PM_{2.5}$ pollution levels over the THB.

Air quality	$PM_{2.5}$ levels ($\mu g m^{-3}$)	Regional transport	Local emissions
Light pollution	$75 \leq PM_{2.5} < 150$	58.9%	41.1%
Heavy pollution	$PM_{2.5} \geq 150$	70.5%	29.5%

Table 5

Removal contributions of precipitation washout to surface $PM_{2.5}$ concentrations.

Periods of heavy air pollution	Removal contributions
The formation stage	17.0 %
The dissipating stage	55.3 %

numerical models with the comprehensive parameterizations of physical and chemical processes in the atmosphere.

CRedit authorship contribution statement

Weiyang Hu: Conceptualization, Methodology, Investigation, Writing - original draft, Writing - review & editing. **Tianlaing Zhao:** Conceptualization, Methodology, Funding acquisition, Writing - review & editing, Supervision. **Yongqing Bai:** Conceptualization, Writing - review & editing, Supervision. **Shaofei Kong:** Conceptualization, Data curation, Resources. **Jie Xiong:** Software, Investigation. **Xiaoyun Sun:** Data processing, Investigation. **Qingjian Yang:** Conceptualization. **Yao Gu:** Investigation. Huicheng Lu: Investigation.

Declaration of competing interest

We declare that we have no financial and personal relationships with other people or organizations that can inappropriately influence our work, there is no professional or other personal interest of any nature or kind in any product, service and/or company that could be construed as influencing the position presented in, or the review of, the manuscript entitled "Importance of Regional PM_{2.5} Transport and Precipitation Washout in Heavy Air Pollution in the Twain-Hu Basin over Central China: Observational analysis and WRF-Chem Simulation".

Acknowledgement

This research was financially funded by grants from National Natural Science Foundation of China (41830965; 42075186; 91744209), the National Key Research and Development Program of China (2016YFC0203304), and the Postgraduate Research & Innovation Project of Jiangsu Province (KYCX20_0951).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2020.143710>.

References

- Bai, Y., Qi, H., Zhao, T., Yang, H., Liu, L., Cui, C., 2018. Analysis of meteorological conditions and diurnal variation characteristics of PM_{2.5} heavy pollution episodes in the winter of 2015 in Hubei province. *Acta Meteorol. Sin.* 76 (5), 803–815 (in Chinese).
- Boylan, J.W., Russell, A.G., 2006. PM and light extinction model performance metrics, goals, and criteria for three-dimensional air quality models. *Atmos. Environ.* 40, 4946–4959.
- Chang, X., Wang, S., Zhao, B., Cai, S., Hao, J., 2018a. Assessment of inter-city transport of particulate matter in the Beijing–Tianjin–Hebei region. *Atmos. Chem. Phys.* 18, 4843–4858.
- Chang, X., Wang, S., Zhao, B., Xing, J., Liu, X., Wei, L., Song, Y., Wu, W., Cai, S., Zheng, H., Ding, D., Zheng, M., 2018b. Contributions of inter-city and regional transport to PM_{2.5} concentrations in the Beijing–Tianjin–Hebei region and its implications on regional joint air pollution control. *Sci. Total Environ.* 660, 1191–1200.
- Chate, D.M., 2005. Study of scavenging of submicron-sized aerosol particles by thunderstorm rain events. *Atmos. Environ.* 39, 6608–6619.
- Chen, H., Lin, Y., Su, Q., Cheng, L., 2017. Spatial variation of multiple air pollutants and their potential contributions to all-cause, respiratory, and cardiovascular mortality across China in 2015–2016. *Atmos. Environ.* 168, 23–35.
- Chen, Q., Sheng, L., Gao, Y., Miao, Y., Hai, S., Gao, S., Gao, Y., 2019a. The effects of the trans-regional transport of PM_{2.5} on a heavy haze event in the Pearl River Delta in January 2015. *Atmos. 10* (5), 237.
- Chen, L., Zhu, J., Liao, H., Gao, Y., Qiu, Y., Zhang, M., Liu, Z., Li, N., Wang, Y., 2019b. Assessing the formation and evolution mechanisms of severe haze pollution in the Beijing–Tianjin–Hebei region using process analysis. *Atmos. Chem. Phys.* 19, 10845–10864.
- Ding, Y., 1993. *Monsoons Over China*. Springer Science & Business Media.
- Du, X.X., Shi, G.M., Zhao, T.L., Yang, F.M., Zheng, X.B., Zhang, Y.J., Tan, Q.W., 2020. Contribution of secondary particles to wintertime PM_{2.5} during 2015–2018 in a major urban area of the Sichuan Basin, Southwest China. *Earth. Space. Sci.* 7 (6), e2020EA001194.
- Feng, X., Wang, S., 2012. Influence of different weather events on concentrations of particulate matter with different sizes in Lanzhou, China. *J. Environ. Sci.* 24 (4), 665–674.
- Gao, Y., Liu, X., Zhao, C., Zhang, M., 2011. Emission controls versus meteorological conditions in determining aerosol concentrations in Beijing during the 2008 Olympic Games. *Atmos. Chem. Phys.* 11 (23), 12437.
- Grell, G.A., Peckham, S.E., Schmitz, R., McKeen, S.A., Frost, G., Skamarock, W.C., Eder, B., 2005. Fully coupled "online" chemistry within the WRF model. *Atmos. Environ.* 39 (37), 6957–6975.
- Guo, S., Hu, M., Zamora, M.L., Peng, J., Shang, D., Zheng, J., Du, Z., Wu, Z., Shao, M., Zeng, L., Molina, M.J., Zhang, R., 2014. Elucidating severe urban haze formation in China. *Proc. Natl. Acad. Sci.* 111 (49), 17373–17378.
- Han, S., Liu, J., Hao, T., Zhang, Y., Li, P., Yang, J., Wang, Q., Cai, Z., Yao, Q., Zhang, M., Wang, X., 2018. Boundary layer structure and scavenging effect during a typical winter haze-fog episode in a core city of BTH region, China. *Atmos. Environ.* 179, 187–200.
- Hua, Y., Wang, S., Wang, J., Jiang, J., Zhang, T., Song, Y., Kang, L., Zhou, W., Cai, R., Wu, D., Fan, S., Wang, T., Tang, X., Wei, Q., Sun, F., Xiao, Z., 2016. Investigating the impact of regional transport on PM_{2.5} formation using vertical observation during APEC 2014 Summit in Beijing. *Atmos. Chem. Phys.* 16, 15451–15460.
- Huang, Y., Deng, T., Li, Z., Wang, N., Yin, C., Wang, S., Fan, S., 2018. Numerical simulations for the sources apportionment and control strategies of PM_{2.5} over Pearl River Delta, China, part I: inventory and PM_{2.5} sources apportionment. *Sci. Total Environ.* 634, 1631–1644.
- Huang, X., Ding, A., Wang, Z., Ding, K., Gao, J., Chai, F., Fu, C., 2020. Amplified transboundary transport of haze by aerosol–boundary layer interaction in China. *Nat. Geosci.* 13, 428–434.
- Ji, Y., Qin, X., Wang, B., Xu, J., Shen, J., Chen, J., Huang, K., Deng, C., Yan, R., Xu, K., Zhang, T., 2018. Counteractive effects of regional transport and emission control on the formation of fine particles: a case study during the Hangzhou G20 summit. *Atmos. Chem. Phys.* 18, 13581–13600.
- Jiang, C., Wang, H., Zhao, T., Li, T., Che, H., 2015. Modeling study of PM_{2.5} pollutant transport across cities in China's Jing–Jin–Ji region during a severe haze episode in December 2013. *Atmos. Chem. Phys.* 15, 5803–5814.
- Kan, H., Chen, R., Tong, S., 2012. Ambient air pollution, climate change, and population health in China. *Environ. Int.* 42, 10–19.
- Kang, H., Zhu, B., Gao, J., He, Y., Wang, H., Su, J., Pan, C., Zhu, T., Yu, B., 2019. Potential impacts of cold frontal passage on air quality over the Yangtze River Delta, China. *Atmos. Chem. Phys.* 19, 3673–3685.
- Li, J., Du, H., Wang, Z., Sun, Y., Yang, W., Li, J., Tang, X., Fu, P., 2017a. Rapid formation of a severe regional winter haze episode over a mega-city cluster on the North China Plain. *Environ. Pollut.* 223, 605–615.
- Li, Z., Guo, J., Ding, A., Liao, H., Liu, J., Sun, Y., Wang, T., Xue, H., Zhang, H., Zhu, B., 2017b. Aerosol and boundary-layer interactions and impact on air quality. *Natl. Sci. Rev.* 4, 810–833.
- Li, X., Hu, X.-M., Ma, Y., Wang, Y., Li, L., Zhao, Z., 2019a. Impact of planetary boundary layer structure on the formation and evolution of air-pollution episodes in Shenyang, Northeast China. *Atmos. Environ.* 214, 116850.
- Li, R., Mei, X., Wei, L., Han, X., Zhang, M., Jing, Y., 2019b. Study on the contribution of transport to PM_{2.5} in typical regions of China using the regional air quality model RAMS-CMAQ. *Atmos. Environ.* 214, 116856.
- Liu, J., Han, Y., Tang, X., Zhu, J., Zhu, T., 2016. Estimating adult mortality attributable to PM_{2.5} exposure in China with assimilated PM_{2.5} concentrations based on a ground monitoring network. *Sci. Total Environ.* 568, 1253–1262.
- Lu, M., Tang, X., Wang, Z., Gbaguidi, A., Liang, S., Hu, K., Wu, L., Wu, H., Huang, Z., Shen, L., 2017. Source tagging modeling study of heavy haze episodes under complex regional transport processes over Wuhan megacity, Central China. *Environ. Pollut.* 231, 612–621.
- Lu, M., Tang, X., Wang, Z., Wu, L., Chen, X., Liang, S., Zhou, H., Wu, H., Hu, K., Shen, L., Yu, J., Zhu, J., 2019. Investigating the transport mechanism of PM_{2.5} pollution during January 2014 in Wuhan, Central China. *Adv. Atmos. Sci.* 36, 1217–1234.
- Luan, T., Guo, X., Zhang, T., Guo, L., 2019. Below-cloud aerosol scavenging by different-intensity rains in Beijing city. *J. Meteor. Res.* 33 (1), 126–137.
- Ma, Q., Wu, Y., Zhang, D., Wang, X., Xia, Y., Liu, X., Tian, P., Han, Z., Xia, X., Wang, Y., Zhang, R., 2017. Roles of regional transport and heterogeneous reactions in the PM_{2.5} increase during winter haze episodes in Beijing. *Sci. Total Environ.* 599, 246–253.
- Miao, Y., Guo, J., Liu, S., Liu, H., Zhang, G., Yan, Y., He, J., 2017. Relay transport of aerosols to Beijing–Tianjin–Hebei region by multi-scale atmospheric circulations. *Atmos. Environ.* 165, 35–45.
- Miao, Y., Liu, S., Sheng, L., Huang, S., Li, J., 2019. Influence of boundary layer structure and low-level jet on PM_{2.5} pollution in Beijing: a case study. *Int. J. Environ. Res. Public Health* 16 (4), 616.
- Ning, G., Wang, S., Yim, S.H.L., Li, J., Hu, Y., Shang, Z., Wang, J., Wang, J., 2018. Impact of low-pressure systems on winter heavy air pollution in the Northwest Sichuan Basin, China. *Atmos. Chem. Phys.* 18, 13601–13615.
- Pope, C.A., Dockery, D.W., 2006. Health effects of fine particulate air pollution: lines that connect. *J. Air Waste Manage. Assoc.* 56, 709–742.
- Seinfeld, J.H., Pandis, S.N., 2006. *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*. second ed. John Wiley & Sons, Inc, New Jersey, USA.
- Shen, L., Wang, H., Zhao, T., Liu, J., Bai, Y., Kong, S., Shu, Z., 2020a. Characterizing regional aerosol pollution in Central China based on 19 years of MODIS data: spatiotemporal variation and aerosol type discrimination. *Environ. Pollut.* 114556.
- Shen, L., Zhao, T., Wang, H., Liu, J., Bai, Y., Kong, S., Zheng, H., Zhu, Y., Shu, Z., 2020b. Importance of meteorology in air pollution events during the city lockdown for COVID-19 in Hubei Province, Central China. *Sci. Total Environ.* 754, 142227.
- Sun, Y., Chen, C., Zhang, Y., Xu, W., Zhou, L., Cheng, X., Zheng, H., Ji, D., Li, X., Tang, X., Fu, P., Wang, Z., 2016. Rapid formation and evolution of an extreme haze episode in northern China during winter 2015. *Sci. Rep.* 6 (1), 1–9.
- Tao, M., Chen, L., Su, L., Tao, J., 2012. Satellite observation of regional haze pollution over the North China Plain. *J. Geophys. Res. Atmos.* 117 (D12).
- Wang, Y., Bao, S., Wang, S., Hu, Y., Shi, X., Wang, J., Zhao, B., Jiang, J., Zheng, M., Wu, M., Russell, A.G., Wang, Y., Hao, J., 2017. Local and regional contributions to fine particulate matter in Beijing during heavy haze episodes. *Sci. Total Environ.* 580, 283–296.

- Wang, Y., Duan, X., Wang, L., 2019a. Spatial-temporal evolution of PM_{2.5} concentration and its socioeconomic influence factors in Chinese cities in 2014–2017. *Int. J. Environ. Res. Public Health* 16 (6), 985.
- Wang, T., Gao, T., Zhang, H., Ge, M., Lei, H., Zhang, P., Zhang, P., Lu, C., Liu, C., Zhang, H., Zhang, Q., Liao, H., Kan, H., Feng, Z., Zhang, Y., Qie, X., Cai, X., Li, M., Liu, L., Tong, S., 2019b. Review of Chinese atmospheric science research over the past 70 years: atmospheric physics and atmospheric environment. *Sci. China Earth Sci.* 62, 1903–1945.
- Wei, P., Ren, Z., Wang, W., Su, F., Gao, Q., Cheng, S., Zhang, Y., 2015. Analysis of meteorological conditions and formation mechanisms of lasting heavy air pollution in eastern China in October 2014. *Res. Environ. Sci.* 28 (5), 676–683.
- Wu, J., Li, G., Cao, J., Bei, N., Wang, Y., Feng, T., Huang, R., Liu, S., Zhang, Q., Tie, X., 2017. Contributions of trans-boundary transport to summertime air quality in Beijing, China. *Atmos. Chem. Phys.* 17, 2035–2051.
- Xu, J., Chang, L., Qu, Y., Yan, F., Wang, F., Fu, Q., 2016. The meteorological modulation on PM_{2.5} inter-annual oscillation during 2013 to 2015 in Shanghai, China. *Sci. Total Environ.* 572, 1138–1149.
- Yang, Y., Quan, J., Wang, Y., Zhang, Y., Cao, G., Zhang, Q., Wang, W., Wang, J., Sun, J., Zhang, X., Li, W., 2013. Factors contributing to haze and fog in China. *Sci. Bull.* 58, 1178–1187.
- Yang, Y., Liu, X., Qu, Y., An, J., Jiang, R., Zhang, Y., Sun, Y., Wu, Z., Zhang, F., Xu, W., Ma, Q., 2015. Characteristics and formation mechanism of continuous hazes in China: a case study during the autumn of 2014 in the North China Plain. *Atmos. Chem. Phys.* 15 (7), 10987–11029.
- Ye, X., Song, Y., Cai, X., Zhang, H., 2016. Study on the synoptic flow patterns and boundary layer process of the severe haze events over the North China Plain in January 2013. *Atmos. Environ.* 124, 129–145.
- Yu, C., Zhao, T., Bai, Y., Zhang, L., Kong, S., Yu, X., He, J., Cui, C., Yang, J., You, Y., Ma, G., Wu, M., Chang, J., 2020. Heavy air pollution with a unique “non-stagnant” atmospheric boundary layer in the Yangtze River middle basin aggravated by regional transport of PM_{2.5} over China. *Atmos. Chem. Phys.* 20, 7217–7230.
- Zhang, H., Wang, Y., Hu, J., Ying, Q., Hu, X.M., 2015a. Relationships between meteorological parameters and criteria air pollutants in three megacities in China. *Environ. Res.* 140, 242–254.
- Zhang, L., Wang, T., Lv, M., Zhang, Q., 2015b. On the severe haze in Beijing during January 2013: unraveling the effects of meteorological anomalies with WRF-Chem. *Atmos. Environ.* 104, 11–21.
- Zhang, X.Y., Wang, J.Z., Wang, Y.Q., Liu, H.L., Sun, J.Y., Zhang, Y.M., 2015c. Changes in chemical components of aerosol particles in different haze regions in China from 2006 to 2013 and contribution of meteorological factors. *Atmos. Chem. Phys.* 15, 12935–12952.
- Zhang, H., Yuan, H., Liu, X., Yu, J., Jiao, Y., 2018. Impact of synoptic weather patterns on 24h-average PM_{2.5} concentrations in the North China Plain during 2013–2017. *Sci. Total Environ.* 627, 200–210.
- Zhang, H., Cheng, S., Yao, S., Wang, X., Zhang, J., 2019. Multiple perspectives for modeling regional PM_{2.5} transport across cities in the Beijing–Tianjin–Hebei region during haze episodes. *Atmos. Environ.* 212, 22–35.
- Zhao, B., Wu, W., Wang, S., Xing, J., Chang, X., Liou, K.-N., Jiang, J.H., Gu, Y., Jang, C., Fu, J.S., Zhu, Y., Wang, J., Lin, Y., Hao, J., 2017. A modeling study of the nonlinear response of fine particles to air pollutant emissions in the Beijing–Tianjin–Hebei region. *Atmos. Chem. Phys.* 17, 12031–12050.
- Zhong, J., Zhang, X., Wang, Y., Wang, J., Shen, X., Zhang, H., Wang, T., Xie, Z., Liu, C., Zhang, H., Zhao, T., Sun, J., Fan, S., Gao, Z., Li, Y., Wang, L., 2019. The two-way feedback mechanism between unfavorable meteorological conditions and cumulative aerosol pollution in various haze regions of China. *Atmos. Chem. Phys.* 19, 3287–3306.