What more cases buy: inferential targets for reproduction-number estimates and what cross-sectional dispersion diagnostics measure

[Author names and affiliations to be added before submission]

5 November 2026

Abstract

Case counts often set the precision attributed to reproduction-number estimates, but the error that more cases remove depends on the inferential target. In a one-generation branching model with negative binomial offspring and binomial reporting, we derive exact mean squared errors for the current reproduction number, the realised mean over observed periods and the process mean. Counting error falls inversely with expected count for all three targets, whereas the process mean also retains between-period variance; an observed denominator raises counting error 1.5-fold at large counts and 2.4-fold at ten expected cases. Single-period count moments cannot separate fluctuating reporting from transmission variation. We then applied an analysis plan, written after exploratory analysis of the same data, to 1,748 state-weeks of US influenza hospital admissions (51 jurisdictions, seasons 2022–23 to 2025–26), regressing the squared relative deviation of an apparent reproduction number on inverse admission count. In the primary test the count-independent intercept was 0.0150 (two-way cluster-robust 95% CI 0.0109–0.0191), and it stayed above zero in all 17 sensitivity analyses of the plan. The affine form was an approximation: it missed the observed mean in three of ten count deciles, and with a free count exponent the block-bootstrap interval of the intercept reached zero. Without a model, state-weeks above about 390 admissions had a relative error of 10.5% (block bootstrap 8.8–11.8%), 10.1% net of counting noise. The intercept halved from small to large jurisdictions (ratio 0.50, state bootstrap 0.41–0.61). It reflects transmission and reporting variation, fell 6–22% short of their sum in simulations with reporting shocks, and its transmission share is not identified. No difference between season intercepts was detected (permutation $p = 0.102$; power 0.47).

Keywords: reproduction number; surveillance; overdispersion; reporting variation; jurisdiction size

1 Introduction

Reproduction numbers summarise transmission under specified conditions and are routinely estimated from surveillance counts [1, 2]. Renewal methods relate incidence to past infectiousness through a generation-interval distribution, and the width of their credible intervals shrinks as the number of cases grows [3, 4]. Work on low incidence, reporting noise and timely detection has made the limits of period-specific inference explicit [5–7], and temporal aggregation and reporting delays separate the reproduction number of the transmission process from the apparent reproduction number of an observed series [8–10].

Precision guidance for renewal estimators is often stated in cases. Under a gamma prior with shape a_0, the posterior coefficient of variation of the renewal estimator of Cori et al. [3] is $(a_0 + I_W)^{-1/2}$, where I_W is the number of cases in the window, so it is a function of the window count alone. Their Web Appendix 3 uses this relation to suggest that estimation start once at least 12 cases have accrued since the first case, which gives a posterior coefficient of variation of about 0.3; this heuristic is one of three criteria they give for the start of estimation. The coefficient measures counting error for the average transmission level over the window. The estimate is also often read as the transmission level now or as a typical level for the population. For those targets, variation between periods is an additional error component that more cases do not remove. A count-based criterion is therefore complete for one target and incomplete for the other two.

The variance structure behind this distinction is classical. Variance-component models split the variance of a unit-level estimate into a within-unit sampling term and a between-unit term [11, 12]. Random-effects meta-analysis estimates the between-study component on top of within-study error [13], and small-area estimation and empirical-Bayes disease mapping weight each direct estimate against a common mean according to the ratio of the two terms [14–16]. Funnel plots for institutional comparison draw control limits from the counting term and, when units differ beyond chance, add an overdispersion component; in the additive random-effects form the variance of a unit indicator based on n events is $\sigma^2/n + \tau^2$ [17, 18]. The affine dispersion model used below has exactly this form. Population ecology makes the same split between demographic and environmental stochasticity [19, 20], as do stochastic epidemic models [21, 22]. In each setting the count at which the two terms are equal is where a shrinkage estimator places half its weight on the direct estimate.

Several results used below are standard and are cited where they are applied. The mean squared error decompositions follow from the law of total variance [11, 23]. The efficiency of the count-to-scale ratio for a Poisson mean is the Cramér–Rao bound [24], the delta-method expansion of a ratio estimator is textbook material [23], and the square-root allocation of surveillance effort is Neyman allocation [12]. Offspring overdispersion follows Lloyd-Smith et al. [25], and its estimation from small samples Lloyd-Smith [26]. A variance that scales as a power of the mean is Taylor's law [27]. The distinction between the realised quantity of a finite population and the parameter of the process that generated it follows Graubard and Korn [28]; the case reproduction number of Wallinga and Teunis [29] is a realised quantity of this kind. Constant under-reporting cancels from ratio estimators of the reproduction number, whereas reporting that changes over time does not [2, 3]. Choosing the estimation window to balance counting noise against changes in transmission is the aim of adaptive windowing [30], and hierarchical renewal models pool an observation layer across regions [31].

The paper is a synthesis of these results for period-specific reproduction numbers, together with a cross-sectional diagnostic and an application under a written analysis plan. Section 2 sets out three inferential targets, derives their exact mean squared errors in a branching model with negative binomial offspring and binomial reporting, and gives the correction required when the denominator is the previous observed count. Section 3 shows that, when the reporting probability fluctuates between periods, single-period second moments of the counts cannot separate reporting volatility from transmission variation. Section 4 defines the diagnostic, an affine regression of the squared relative deviation of an apparent reproduction number on inverse count, maps its intercept and slope to the quantities of the model, and states the analysis plan. Section 5 applies the plan to 1,748 state-weeks of US influenza hospital admissions. The primary question is narrow: whether a count-independent component of dispersion remains after counting noise is allowed for. The secondary questions are how large it is at the counts actually observed, whether the affine form describes it, and whether it is common to jurisdictions of different size.

The question matters because routine hospital surveillance does not observe the variables of a branching experiment. Admissions follow infection after a delay, aggregate heterogeneous populations and are recorded on a calendar schedule, and their count scale is an outcome of the epidemic. Reporting probabilities may change for all infections in the same period [32, 33], and separating process from observation variation in time series is a known difficulty [34, 35]. The cross-sectional spread of growth-rate or reproduction-number estimates has been read as variation in transmission or behaviour between places and periods without subtracting the part of that spread that depends on case counts [36, 37], although per-unit standard errors make the subtraction possible [38]. The diagnostic performs that subtraction. Its intercept bounds what additional cases can buy, but by the result of Section 3 it does not separate transmission from reporting, and the analysis reads it as reflecting both rather than as an estimate of their sum. Section 6 discusses real-time use, reporting that depends on transmission and the choice of generation interval. Proofs, extensions and a known-truth simulation audit on the real design are in the appendices.

2 Model and inferential targets

2.1 A branching experiment with a known observation scale

Consider one generation of transmission with $n \geq 1$ primary infections. Conditional on the reproduction number $R_t > 0$, the offspring counts of primary cases are independent negative binomial variables [25]:

$$X_{i,t} \mid R_t \sim \text{NB}(k,\, k/(k+R_t)), \qquad \text{E}(X_{i,t} \mid R_t) = R_t, \quad \text{Var}(X_{i,t} \mid R_t) = R_t + \frac{R_t^2}{k}, \tag{1}$$

where $k > 0$ controls individual heterogeneity. Each secondary infection is independently reported with probability $\rho \in (0, 1]$. Summation over primary cases and binomial thinning give the law of the reported count:

$$C_t \mid n, R_t \sim \text{NB}(nk,\, k/(k+\rho R_t)), \qquad \text{E}(C_t \mid n, R_t) = \rho n R_t, \qquad \text{Var}(C_t \mid n, R_t) = \rho n R_t + \frac{\rho^2 n R_t^2}{k}. \tag{2}$$

Define $m = \rho n$ and $\hat{R}_t = C_t/m$. The scale m is the expected reported count at $R_t = 1$, not the realised count C_t.

The baseline assumptions are fixed, known ρ with independent reporting (H1); conditionally independent offspring as in Eq. (1) (H2); fixed, known k (H3); and a fixed or exogenous n that does not depend on transmission (H4). Write $\text{E}(R_t) = \overline{R} > 0$ and $\text{Var}(R_t) = v_R < \infty$. Across periods, counts are independent given the whole transmission trajectory, each with conditional law (2). These are repeated branching experiments; they do not assert that naturally evolving incidence is exogenous. Dividing by an assumed $\rho_0 \neq \rho$ gives conditional mean $(\rho/\rho_0)R_t$, a multiplicative bias that more cases do not remove.

The conditional counting variance combines offspring randomness and omission. Conditioning on the total number of secondary infections splits it into $(R_t + R_t^2/k)/n$ and $(1-\rho)R_t/(\rho n)$. The term linear in R_t is therefore not purely reporting noise; it remains under complete reporting.

Table 1 lists the symbols used throughout.

Table 1 Notation

Symbol	Meaning
R_t	Reproduction number in period t (current target)
$\overline{R}$, v_R	Mean and variance of R_t under the stated transmission law (process mean and between-period transmission variance)
$\overline{R}_T$	Mean of $R_1, \ldots, R_T$ over the observed periods (realised mean)
n, ρ, k	Primary infections, reporting probability, negative binomial dispersion
$m = \rho n$	Expected reported count at $R_t = 1$ (observation scale)
C_t, $\hat{R}_t = C_t/m$	Reported secondary count and the count-to-scale estimator
M_t, $\tilde{R}_t = C_t/M_t$	Reported primary count and the observed-denominator estimator
A	Counting-error numerator, $\overline{R} + \rho(\overline{R}^2 + v_R)/k$
$m_\times = A/v_R$	Crossover scale at which counting error equals v_R
$\gamma(m)$	Weight of the direct estimate in the best linear predictor
p_t, τ^2	Period-level reporting probability and its squared relative variation
v_H, A_H	Reporting-volatility variance and counting numerator under H5
a, b	Intercept and inverse-count slope of the affine dispersion diagnostic
b_P	Independent-Poisson reference for the slope

2.2 Current values, realised means and process means

For the current target, $\mathrm{E}(\hat{R}_t \mid R_t) = R_t$ and $\mathrm{Var}(\hat{R}_t \mid R_t) = (R_t + \rho R_t^2/k)/m$, which attains the Cramér–Rao bound for estimators unbiased for every R_t under this likelihood (Appendix A.1).

Proposition 1 (One period, two targets)**.** *Under H1–H4, let $A = \overline{R} + \rho(\overline{R}^2 + v_R)/k$. The two risks of $\hat{R}_t$ are:*

$$\mathrm{E}[(\hat{R}_t - R_t)^2] = \frac{A}{m}, \qquad V(m) := \mathrm{E}[(\hat{R}_t - \overline{R})^2] = v_R + \frac{A}{m}. \tag{3}$$

Among estimators based on this period's count that are conditionally unbiased for R_t, $\hat{R}_t$ minimises both risks.

The proof (Appendix A.2) squares the error decomposition:

$$\hat{R}_t - \overline{R} = (\hat{R}_t - R_t) + (R_t - \overline{R}). \tag{4}$$

The two terms on the right are uncorrelated. Between-period transmission variation is part of the signal for the current value and sampling error for the process mean, which is a mean under a stated stationary law and not automatically a seasonal average of a nonstationary epidemic.

For $v_R > 0$ define the crossover scale $m_\times = A/v_R$, the expected count at which the two terms of $V(m)$ are equal. Its meaning is the one it has in small-area estimation [14]. The best linear predictor of R_t from $\hat{R}_t$ under the stated moments takes the form:

$$\overline{R} + \gamma(m)(\hat{R}_t - \overline{R}), \qquad \gamma(m) = \frac{v_R}{v_R + A/m} = \frac{m}{m + m_\times}, \tag{5}$$

so $m_\times$ is the count at which shrinkage gives the direct estimate half its weight. Doubling m through n at fixed ρ, k and transmission law reduces process-mean risk by the fraction $1/[2(1 + m/m_\times)]$, which is one quarter at $m = m_\times$ and less beyond it. Changing ρ changes A as well as m. The scale $m_\times$ describes the selected estimator and target; it does not mark a point where information stops accumulating.

Corollary 1 (A fixed set of observed periods)**.** *Under H1–H4 and conditional independence of counts given $R_{1:T}$, let $\overline{R}_T$ denote the mean of $R_1, \ldots, R_T$ and $\overline{\hat{R}}_T$ the mean of $\hat{R}_1, \ldots, \hat{R}_T$. If the R_t share common marginal moments, the two mean targets have risks:*

$$\mathrm{E}[(\overline{\hat{R}}_T - \overline{R}_T)^2] = \frac{A}{Tm}, \qquad \mathrm{E}[(\overline{\hat{R}}_T - \overline{R})^2] = B_T + \frac{A}{Tm}, \qquad B_T = \mathrm{Var}(\overline{R}_T). \tag{6}$$

In Eq. (6)*, $B_T = v_R/T$ for independent, identically distributed R_t. For a stationary process with $\mathrm{Cov}(R_t, R_{t+h}) = v_R\phi^{|h|}$, $|\phi| < 1$, $B_T = v_R S(\phi, T)/T^2$ exactly, where the window sum is:*

$$S(\phi, T) = T + 2\sum\nolimits_{h=1}^{T-1}(T - h)\phi^h = \frac{T(1+\phi)}{1-\phi} - \frac{2\phi(1-\phi^T)}{(1-\phi)^2}. \tag{7}$$

The realised-mean identity needs only conditionally independent counting errors, not independent transmission levels. More cases in the same periods remove its counting error, while process-mean risk tends to B_T; only more periods reduce B_T, at a rate set by their dependence. Positive persistence reduces the gain from adjacent periods. Spatial averaging behaves analogously. For J units with transmission variance v_R and mean pairwise correlation $\bar{c}$, the averaged transmission level has the variance:

$$\mathrm{Var}(J^{-1}\textstyle\sum_{j=1}^{J} R_{j,t}) = v_R[\bar{c} + (1 - \bar{c})/J]. \tag{8}$$

Figure 1 plots these risks, with known-truth Monte Carlo checks, and Table 2 collects them.

Table 2 Inferential targets and exact mean squared errors. The fixed-reporting column assumes H1–H4; the volatile-reporting column assumes H2–H5 and, for averages, independent reporting probabilities independent of the transmission path. Counting errors are conditionally independent across periods. $B_T = \mathrm{Var}(\overline{R}_T)$ equals v_R/T for independent, identically distributed transmission and $v_R S(\phi, T)/T^2$ under the stated stationary autocovariance. With the previous observed count as denominator, the current-value entry under fixed reporting becomes approximately $[A + (1-\rho)(\overline{R}^2 + v_R)]/m$ (Eq. 9; Table 3).

Target	Estimator	Fixed, known reporting	Unobserved volatile reporting
Current R_t	$\hat{R}_t$	A/m	$v_H + A_H/m$
Realised mean $\overline{R}_T$	$\hat{\overline{R}}_T$	$A/(Tm)$	$(v_H + A_H/m)/T$
Process mean $\overline{R}$	$\hat{\overline{R}}_T$	$B_T + A/(Tm)$	$B_T + (v_H + A_H/m)/T$

2.3 An observed denominator

Surveillance ratios divide by an observed count rather than by a known scale. Let the denominator be the reported count of the primary generation, $M_t \sim \mathrm{Bin}(n, \rho)$, independent of the offspring given n, and let $\tilde{R}_t = C_t/M_t$. Because $M_t = m(1 + u_t)$ with $\mathrm{Var}(u_t) = (1-\rho)/m$, a delta expansion (Appendix A.5) gives the approximate risk:

$$\mathrm{E}[(\tilde{R}_t - R_t)^2] \approx \frac{A + (1-\rho)(\overline{R}^2 + v_R)}{m}, \tag{9}$$

so the crossover scale moves to $m_\times + (1-\rho)(\overline{R}^2 + v_R)/v_R$. With $\overline{R} = 1.15$, $v_R = 0.04$, $k = 0.35$ and $\rho = 0.25$ ($A = 2.1232$) it rises from 53.1 to 78.6. Monte Carlo with 400,000 draws per scale (Table 3) puts the observed-denominator error at 1.59, 1.51 and 1.49 times A/m for $m = 50$, 200 and 1000, close to the delta value of 1.48. At $m = 10$ the ratio is 2.42 and the delta form understates the error, because small denominators inflate the ratio. The ratio estimator is undefined when $M_t = 0$; such draws are excluded and their share is reported in Table 3, where it is zero to the precision shown at every scale considered. At $m = 10$ the Monte Carlo error is 0.5145 with standard error 0.0048. A continuity-corrected ratio, $(C_t + 0.5)/(M_t + 0.5)$, lowers it to 0.4253, still 2.00 times A/m.




Figure 1 Model risks for the three targets. Panel A shows the mean squared error against expected count m for the current value, the realised mean and the process mean over $T = 10$ independent periods. Panel B shows the current-value risk with fixed reporting ($\tau = 0$) and with period-level reporting variation ($\tau = 0.15$). Panel C shows the exact risks against known-truth Monte Carlo estimates with 1.96 Monte Carlo standard errors. Panel D shows the realised-mean risk over $T = 10$ periods at three scales, with $\overline{R} = 1.15$, $v_R = 0.04$, $k = 0.35$ and $\rho = 0.25$. Counts are simulated under the fixed-exposure model, so the curves describe that model and not the hospital data.

Table 3 Point-target error with a known versus an observed denominator

m	Known denominator A/m	MC (SE)	Observed denominator (A + (1 - rho)(Rbar^2 + v_R))/m	MC (SE)	MC observed / A/m	Zero denominators	+0.5 variant MC (SE)
10	0.21232	0.21304 (0.00055)	0.31451	0.51453 (0.00482)	2.423	0	0.42529 (0.00276)
50	0.04246	0.04234 (0.00010)	0.06290	0.06745 (0.00019)	1.588	0	0.06589 (0.00018)
200	0.01062	0.01060 (0.00002)	0.01573	0.01601 (0.00004)	1.508	0	0.01592 (0.00004)
1000	0.00212	0.00212 (0.00000)	0.00315	0.00316 (0.00001)	1.489	0	0.00316 (0.00001)

Parameters Rbar = 1.15, v_R = 0.04, k = 0.35, rho = 0.25 (A = 2.1232); 400,000 Monte Carlo draws per m. Crossing scale m_x = A/v_R = 53.08 with a known denominator and (A + (1 - rho)(Rbar^2 + v_R))/v_R = 78.63 when the denominator is the observed previous count (delta approximation). At m = 10 the delta approximation understates the Monte Carlo error of the observed-denominator estimator. Zero denominators: draws with a zero previous count are excluded from the ratio estimator (share shown); the +0.5 variant (C_t + 0.5)/(C_prev + 0.5) keeps every draw.

3 Reporting volatility and identification

3.1 Period-level reporting variation

Replace H1 by a common period-specific reporting probability $p_t \in [0, 1]$, independent of R_t and of n, with known mean $\rho > 0$ and $\mathrm{Var}(p_t) = \rho^2\tau^2$ (H5). Reporting remains independent across infections given p_t. Because p_t lies in $[0, 1]$ with mean ρ, $\mathrm{Var}(p_t) \le \rho(1 - \rho)$ [39], so $\tau^2 \le (1 - \rho)/\rho$. The estimator still divides by $m = \rho n$ and does not observe p_t.

Proposition 2 (Reporting volatility and second-moment nonidentification)**.** *Under H2–H5, put* $Q = \overline{R}^2 + v_R$. *The first*

two moments of $\hat{R}_t$ are:

$$\mathrm{E}(\hat{R}_t) = \overline{R}, \qquad \mathrm{Var}(\hat{R}_t) = v_R + v_H + \frac{A_H}{m}, \qquad v_H = \tau^2 Q, \qquad A_H = \overline{R} + \frac{\rho(1+\tau^2)Q}{k}. \tag{10}$$

For known m, ρ and k, the single-period mean and variance in Eq. (10) *depend on v_R and τ^2 only through $U = (1+\tau^2)(\overline{R}^2 + v_R)$.*

Given R_t and p_t the estimator has mean $(p_t/\rho)R_t$, so a shared reporting shock is not averaged away by more infections in the same period. The statement is deliberately narrow. It concerns the first two moments of one period's count within the specified family. It does not exclude identification from full count distributions, from temporal structure under further assumptions, or from external measurements of reporting; it also assumes that reporting is independent of transmission, an assumption Section 6 relaxes.

Corollary 2 (Current and realised-mean risks with volatile reporting)**.** *Under H2–H5, the current-value risk is:*

$$\mathrm{E}[(\hat{R}_t - R_t)^2] = v_H + \frac{A_H}{m}. \tag{11}$$

If, in addition, the p_t are independent and identically distributed, independent of the transmission trajectory, and counts are independent given both trajectories, the two mean targets have risks:

$$\mathrm{E}[(\overline{\hat{R}}_T - \overline{R}_T)^2] = \frac{v_H + A_H/m}{T}, \qquad \mathrm{E}[(\overline{\hat{R}}_T - \overline{R})^2] = B_T + \frac{v_H + A_H/m}{T}. \tag{12}$$

At fixed T, unobserved reporting volatility leaves the floor v_H/T on the realised-mean target as m grows, so a target free of transmission variation is not free of reporting error. The crossover scale with volatile reporting, $m_{\times,H} = A_H/(v_R+v_H)$, is never larger than $m_\times$ (Eq. (A.5)); the lower crossover accompanies a larger risk, not better surveillance. Proofs are in Appendix A.4.

3.2 From the branching experiment to surveillance counts

If the scale m is random and associated with transmission, conditional unbiasedness given (R_t, m) still yields the conditional risk:

$$\mathrm{E}[(\hat{R}_t - \overline{R})^2 \mid m] = \frac{\mathrm{E}(R_t \mid m) + \rho\,\mathrm{E}(R_t^2 \mid m)/k}{m} + \mathrm{Var}(R_t \mid m) + [\mathrm{E}(R_t \mid m) - \overline{R}]^2, \tag{13}$$

and the conditional moments in Eq. (13) can change with scale. A constant intercept plus a constant coefficient on inverse count therefore does not follow when the count scale is itself an epidemic outcome, as it is in surveillance. Delays, aggregation, ratio denominators and changing conditional means further separate an apparent statistic from the experiment above. The rest of the paper accordingly treats empirical dispersion diagnostics as descriptive quantities and asks, with a written analysis plan and simulation checks, what they recover.

4 Data, diagnostic and analysis plan

4.1 Data and apparent reproduction number

We used the National Healthcare Safety Network (NHSN) weekly hospital respiratory-data snapshot retrieved on 20 October 2026 and restricted influenza admissions to the 50 US states and the District of Columbia [40, 41]. Hospital participation varied during the series. Reporting was voluntary from May to October 2024 before it became mandatory again from 1 November 2024 [42], and the national Respiratory Illnesses Data Channel was not updated from 3 October to 13 November 2025 [43]. The analysis describes this surveillance system and does not estimate an effect of either event. Two external sources were added for the analyses of Section 4.6: state resident populations from the Census Bureau Vintage 2025 estimates [44], and the issue history of the same NHSN admission series from the Delphi Epidata archive [45], which records the first published and every revised value of each state-week count from November 2024 onwards.

The windows were the early-growth phases of four seasons: 1 October–26 November 2022, 7 October–23 December 2023, 9 November–28 December 2024 and 4 October–27 December 2025. Each is the contiguous national segment with weekly growth of at least 1.05 before the first turning point after the season's admissible start; the 2024 start follows the return to mandatory reporting. The rule is applied identically to every season, and inference conditions on the selected windows. Sensitivity analyses use growth thresholds of 1.02 and 1.10 and fixed calendar windows for the first three seasons (8 October–26 November 2022, 14 October–16 December 2023 and 9 November 2024–18 January 2025). Missing or suppressed counts were kept missing rather than set to zero.

With C_t the admissions in week t, the apparent reproduction number of a state-week is the growth-to-reproduction transform:

$$R_{A,t} = [\max\{0,\, 1 + (\theta/7)\log\{(C_{t+1} + 0.5)/(C_t + 0.5)\}\}]^{\alpha}, \qquad \alpha = (g_\mu/g_\sigma)^2, \quad \theta = g_\sigma^2/g_\mu. \tag{14}$$

The transform follows the renewal relation between growth rate and reproduction number for a gamma-distributed interval [46]. It is not a true reproduction number or a posterior estimate from a renewal model [3, 10]. A row requires observed counts in weeks t and $t + 1$ and $C_t \geq 5$, which leaves 1,748 state-weeks from 51 jurisdictions and 42 calendar weeks. The rows contain 1,969 distinct state-week counts in 210 runs of consecutive weeks.

The primary interval has mean $g_\mu = 3.6$ days and standard deviation $g_\sigma = 1.6$ days, the household serial-interval estimate of Cowling et al. [47]. That estimate is a Weibull fit to 14 household transmission pairs; Eq. (14) uses the gamma distribution with the same mean and standard deviation. A serial interval is not a generation interval [48]. Sensitivity analyses set the mean to 3.2 and 4.0 days and the standard deviation to 1.2 and 2.0 days; the lower mean is close to the intrinsic generation-time estimate from US household data [49].

4.2 The diagnostic and its three summaries

For the apparent statistic $X = R_A$ with pooled mean $\bar{X}$, the relative squared deviation and its affine working model are:

$$L_{st} = \frac{(X_{st} - \bar{X})^2}{\bar{X}^2}, \qquad \mathrm{E}(L_{st} \mid C_{st}) = a + \frac{b}{C_{st}}. \tag{15}$$

This is the empirical counterpart of $V(m)/\overline{R}^2$ in Eq. (3), with a realised count in place of the expected scale and an apparent statistic in place of $\hat{R}_t$. It has the form of the additive overdispersion model of funnel plots [17, 18]. Three summaries of the fit follow.

The intercept a is the count-independent relative dispersion: all relative variation in X that does not scale with count. When reporting probabilities do not vary from week to week ($\tau = 0$), a estimates the relative variance of transmission, $\tilde{v}_R = v_R/\overline{R}^2$. When reporting shocks vary between weeks, a estimates $\tilde{v}_R + \tilde{v}_H$, where $\tilde{v}_H$ is the relative variance that the shocks induce in the ratio statistic (Appendix A.6). A reporting level that is constant over consecutive weeks cancels in the ratio, as it does in renewal estimators [2, 3]. By Proposition 2, single-period cross-sectional moments cannot separate the two components. The estimator also need not weight them equally: when the shocks are drawn independently for each state-week, the fitted intercept recovers only part of $\tilde{v}_H$, and Section 5.6 quantifies the shortfall. The intercept is therefore read throughout as reflecting transmission and reporting variation, not as an estimate of their sum. It is also not an error floor for the current value: with $\tau = 0$ that floor is zero, and with $\tau > 0$ it is $\tilde{v}_H$. For the process mean, a approximates the relative error that more cases cannot remove, to the extent that the affine form holds (Section 5.2).

The counting slope b is read against two references. A delta expansion of Eq. (14) under independent Poisson counts in consecutive weeks gives the independent-Poisson reference $b_{\mathrm{P}} = \overline{\kappa(g)^2[1 + (C_t + 0.5)/(C_{t+1} + 0.5)]}$, a mean over rows, with $\kappa(g) = (g_\mu/7)/[1 + \theta\log(g)/7]$ and $g = (C_{t+1} + 0.5)/(C_t + 0.5)$. Each row is evaluated at its own growth factor and normalised by $\bar{X}^2$, which approximates the expansion around $\bar{X}$ when the row's R_A is close to $\bar{X}$; b_{P} is therefore a plug-in normalisation, not an exact expectation. For the primary sample $b_{\mathrm{P}} = 0.4455$. Two alternatives show how much the normalisation matters. Replacing the unit normalisation by each row's ratio $X_{st}/\bar{X}$ gives 0.4199. The slope fitted to panels of independent Poisson counts drawn around the observed means, with the same transform, row filter and fit, gives 0.3977 (Table 12). The term 1 is the Poisson variance of the denominator count. In a branching chain the denominator week contains the parents of the numerator week, so conditioning on the realised parents leaves only the binomial reporting part of that variance, $(1 - \rho)/C_t$. With Poisson offspring and reporting probability ρ this gives the branching-Poisson reference and its ratio to b_{P}:

$$b_{\mathrm{P,br}}(\rho) = \overline{\kappa(g)^2[(1 - \rho) + (C_t + 0.5)/(C_{t+1} + 0.5)]}, \qquad r(\rho) = \frac{b_{\mathrm{P,br}}(\rho)}{b_{\mathrm{P}}} \approx \frac{1 + (1 - \rho)g}{1 + g} = 1 - \frac{\rho g}{1 + g}. \tag{16}$$

Averaged over the observed rows, $r(\rho)$ is 0.944, 0.861, 0.722 and 0.443 for $\rho = 0.1$, 0.25, 0.5 and 1 (Table S1). The observed-denominator model of Section 2.3 has the structure of the branching-Poisson reference, and the known-scale estimator of Section 2.1 is its limit without denominator noise. The ratio b/b_{P} asks whether counting variation exceeds independent-Poisson noise; $b/b_{\mathrm{P,br}}(\rho)$ asks the same for a branching chain, at a reporting probability that the data do not identify.

The third summary is the crossover count b/a, at which the fitted counting term equals the count-independent term. It is the empirical analogue of $m_\times$ in admissions per week; $m_\times$ refers to the expected count of a branching experiment. At $C = b/a$ a shrinkage estimator of the Fay–Herriot type places half its weight on the direct estimate (Eq. (5)). Table 4 lists the conditions under which each summary identifies its theoretical counterpart and the direction of distortion when they fail.

Table 4 Mapping from theoretical inferential targets to empirical surveillance diagnostics

Theoretical Quantity (Section 2)	Empirical Diagnostic (Section 4)	Exact Equivalence Conditions	Distortion Mechanism / Direction of Bias
Relative transmission variance: $\tilde{v}_R = v_R/\overline{R}^2$	Dispersion Intercept: a	Constant reporting probability ($\tau = 0$); count-independent heterogeneity ($\gamma = 0$).	When reporting varies ($\tau > 0$), a reflects both $\tilde{v}_R$ and $\tilde{v}_H$; in simulation its estimate fell 6–22% short of their sum (Section 5.6). It does not identify $\tilde{v}_R$ and is not an error floor. Under scale dependence ($\gamma > 0$), $a < \tilde{v}_R$.
Counting error scale: $\tilde{A} = A/\overline{R}^2$	Counting Slope: b	Independent Poisson limit ($\rho \to 0$ gives $b \approx b_{\mathrm{P}}$); branching Poisson ($b \approx b_{\mathrm{P,br}}(\rho)$).	Inflated by offspring overdispersion ($k < \infty$, $b > b_{\mathrm{P,br}}$) or unmodelled scale dependence ($\gamma > 0$).
Crossover count: $m_\times = A/v_R$	Crossover Scale: b/a	Exact match between A and b, and between v_R and a.	In presence of reporting shocks ($\tau > 0$), b/a is attenuated below $m_\times$; under scale dependence ($\gamma > 0$), b/a is inflated. b/a is an in-sample dispersion balance point.

Formal correspondence between the theoretical parameters derived under the branching model in Section 2 and the empirical summaries estimated by the cross-sectional dispersion diagnostic in Section 4. The table makes explicit the conditions under which empirical diagnostics identify theoretical quantities and the sign of the distortion when those conditions fail.

4.3 Estimation and inference

The affine mean was fitted by iterated feasible generalised least squares with weights inversely proportional to the squared fitted mean, which minimises a gamma quasi-deviance; this is an estimating criterion, not a distributional claim. The primary interval is the two-way cluster-robust sandwich of Cameron et al. [50] for the stacked estimating equations of $\bar{X}$ and the affine fit, clustered by state and calendar week, with small-sample factors $G/(G-1)$ and critical values from the t distribution on 41 degrees of freedom, one fewer than the number of weeks. The interval for b/a is formed on the log scale. Two resampling checks accompany it: a calendar-week block bootstrap (1,999 draws) that resamples whole weeks with all their state rows, and a wild cluster bootstrap-t over calendar weeks with six-point weights and the null $a = 0$ imposed (9,999 draws) [51, 52]. In the wild bootstrap, losses at or below 10^{-7} in a bootstrap sample are set to 10^{-7} before the restricted fit; this affected 13.5% of bootstrap rows, and Table S12 reports the test under other thresholds and without the clamp. The state-by-week pigeonhole bootstrap [53] used for the supplementary analyses is conservative on this design: under the fresh-draw generators of Section 4.5 its standard deviation of $\hat{a}$ was 1.55–1.78 times the Monte Carlo standard deviation (Table S13). Every interval quoted in the text is listed with its type and source in Table S14; all are 95% intervals, and an interval without a stated type has the type of the nearest preceding interval of the same paragraph.

4.4 Analysis plan and primary test

The analysis plan was written after an exploratory analysis of the same data, and it lists the estimates and the season-test p values that were then known, so that each choice can be judged against them. It was fixed, with a checksum, before any analysis in Section 5 was run. Because the data had been seen, the plan does not make the primary test confirmatory in the sense of an untouched hypothesis; what it fixes is the robustness rule for the intercept, the primary calibration of the season test and the obligation to report every specified analysis. The data vintage, the sample and R_A with the primary interval were frozen.

The primary test (E1) is of $H_0\colon a = 0$ against a two-sided alternative at $\alpha = 0.05$, tested with the two-way sandwich t statistic on 41 degrees of freedom. The claim that count-independent dispersion remains is called robust only if the primary test rejects and the 95% lower limit of a stays above zero in each of nine sensitivity analyses. Two vary the generation-interval mean (S-a1) and standard deviation (S-a2). Three change the row filters: count floors of 10, 20 and 50 (S-a3), the fixed calendar windows (S-a4) and growth thresholds (S-a5). Three remove data: transitions that touch the 2025 data-channel interruption (S-a6), one season at a time (S-a7) and rows with hospital reporting coverage below 90% in either week (S-a8). The last (S-a9) removes one calendar week at a time to measure influence.

A within-week intercept (E3) is a different estimand, not a sensitivity analysis. It replaces $\bar{X}$ by the mean of the same

calendar week computed without the row's own state, which removes common national variation by week; the week mean including the own row and a version that also rescales by the week mean are reported as alternatives. Its interval is the calendar-week block bootstrap.

The secondary test asks whether season intercepts differ. In a model with four season intercepts a_s and a common slope, H_0: $\log a_1 = \cdots = \log a_4$ is tested with the Wald statistic W_{CGM} that uses the two-way sandwich covariance. The season label is constant within a calendar week, so it is assigned at the level of 42 weeks in four groups. With few clusters at the level of assignment, wild cluster bootstrap tests can be badly mis-sized and randomisation inference at that level is recommended [54, 55]. The primary reference distribution is therefore a permutation of season labels across calendar weeks that keeps the number of weeks per season (9,999 permutations); it assumes exchangeable weeks and ignores serial dependence between adjacent weeks. The season difference is called significant only if this $p < 0.05$, whatever the other calibrations show. All other calibrations are reported, and Holm-adjusted p values over E1 and the season test are given. Season intercepts are interpreted only as reflecting transmission and reporting variation together (Proposition 2).

The counting slope b/b_{P}, the crossover count and the predicted relative squared error $a + b/C$ at $C = 10$, 37, 100 and 1,000 were specified for estimation, without tests.

4.5 Simulation checks

The conversion of a into a transmission standard deviation needs a mechanism. For a simulated mechanism with recovery ratio $r = \mathrm{E}(\hat{a})/\tilde{v}_R$, the implied relative standard deviation of the reproduction number is $(a/r)^{1/2}$. Its interval combines the two-way interval of a with the Monte Carlo 2.5–97.5% range of r. The plan required that this quantity be reported as a grid over mechanisms and never as a single estimate, and that $a^{1/2}$ be described as a relative standard deviation of transmission and reporting variation combined, not of transmission alone. A mechanism reproduces the counting slope when its Monte Carlo 2.5–97.5% range of b/b_{P} contains the observed value. Synthetic panels are generated on the exact design of the data. The expected count level is calibrated to the observed median retained count of 45, and the heterogeneity amplitude is calibrated so that the mean fitted intercept equals the observed a (Appendix C). Five checks with fixed criteria were specified:

- SC1, two generations per week: a daily renewal process with the primary generation interval, Poisson or negative binomial ($k = 0.3$) offspring and weekly binomial reporting at $\rho \in \{0.003, 0.01, 0.1\}$. The one-generation recovery map carries over if $\mathrm{E}(\hat{a})/v_R$ lies in $[0.8, 1.25]$ in every cell.
- SC2 and SC3, reporting and offspring dispersion: a grid of $\rho \in \{0.001, 0.003, 0.01, 0.03, 0.1, 0.25, 0.5, 1\}$ and $k \in \{0.1, 0.3, 1, 3, \infty\}$ with independent transmission heterogeneity, 100 panels per cell. SC2 requires intercept recovery in $[0.9, 1.1]$ in every cell; SC3 requires the mean of b/b_{P} to lie in $[0.9, 1.1]$ in every cell with $\rho \leq 0.01$. The range $\rho = 0.001$–0.01 for the probability that an infection becomes a reported admission is assumed from burden estimates [56, 57], not estimated.
- SC4 and SC5, interval validity: these use generators that redraw the heterogeneity field in every replicate. FD-S1 has independent heterogeneity, and FD-S8 has scale-dependent heterogeneity with state-week reporting shocks. FD-S9 adds to FD-S8 a national calendar-week share of 0.25, 0.5 or 0.75 of the heterogeneity; this family was defined after the data had been seen. SC4 requires coverage of $\mathrm{E}(\hat{a})$ by the two-way interval of at least 0.90 under every generator. SC5 calls the gap between the observed standard error of a and the generators' Monte Carlo standard deviation explained if their ratio lies in $[0.67, 1.5]$ for at least one generator.

4.6 Amendments and additional analyses

The plan was amended twice, and both amendments are distributed with it. Amendment 1 recorded implementation details that the plan had left open: seeds, the restricted fit and loss clamp of the wild bootstrap, and the construction of the daily-renewal generator. It stated that these details were fixed before the full runs, but the run logs show that the full runs of the primary test, the season test and the simulation checks had finished before it was written. The details were already in the code used for those runs, and no evidence was found that any of them was changed after a result was seen. The season script was edited once after its first full run. The text of that edit could not be recovered, but every quantity logged by both runs is identical, including the primary permutation p value, the other calibrations, the data sensitivities and the power analysis.

Amendment 2 was fixed, with a checksum, before any of the following analyses was run. Each is secondary or descriptive, none can change the verdict of the primary test, the significance level, the calibration or the data, and each was specified with its reading rule.

- Functional form. The observed mean loss in each decile of C_t, with a calendar-week block-bootstrap interval, is compared with the affine fit, a power law $cC^{-\gamma}$ and a three-parameter model $a + bC^{-\gamma}$. The intercept is said to survive a free exponent if the block-bootstrap interval of a in the three-parameter model excludes zero; $\gamma = 1$ is called compatible with the data if its interval contains one.

- Model-free floor. Among rows with C_t at or above the 75th, 90th and 95th percentiles, the mean loss minus the mean counting term, with either $\hat{b}$ or b_{P} as the slope, estimates the count-independent component without extrapolation.
- Population size. The intercept is made linear in the standardised log population z of the jurisdiction, $a_0 + a_1 z$, with a common slope, and $a_1 = 0$ is tested with the two-way sandwich on 41 degrees of freedom. If the interval of a_1 lies below zero, statements about what more cases buy are restricted to jurisdictions of comparable size and the error at a given count is reported by population tertile. A log-link intercept, a slope that also varies with z, and separate fits by tertile are secondary.
- Bias of the intercept. Under the fresh-draw generators with reporting shocks, and in a sweep of the transmission share of their variance (1, 0.75, 0.25), the relative bias $\mathrm{E}(\hat{a})/(\tilde{v}_R + \tilde{v}_H) - 1$ and the weight w_H that solves $\mathrm{E}(\hat{a}) = \tilde{v}_R + w_H \tilde{v}_H$ are estimated. If every bias interval excludes zero, $\hat{a}$ is described as reflecting, not estimating, the total.
- Generator realism. Six cells of a state-realistic generator anchor transmission to the smoothed observed epidemic curve, scale its variance by state and allow clustered admissions. Four tests compare the Monte Carlo range of each cell with the data: the slope ratio b/b_{P} (T1), the 90th percentile of C_t (T2), the ratio of pooled to within-state slope (T3) and the between-state coefficient of variation of the intercepts (T4). The generators reproduce the data if all four pass in one cell. Their design used the observed slope and between-state dispersion, and no inferential result rests on them.
- Reporting data. The fit is repeated with counts scaled by the share of hospitals reporting in each state-week, and on the first-issue and latest-issue counts of the archive for the 661 rows whose first issue is provisional.
- Calibration details. The wild bootstrap is repeated under four loss clamps and without one; the standard deviation of the pigeonhole bootstrap is compared with the Monte Carlo standard deviation of $\hat{a}$ (it is called conservative only if the ratio exceeds one under every fresh-draw generator); and b/b_{P} is reported under the three Poisson references.

5 Results for US influenza hospital admissions

5.1 A count-independent floor

The pooled affine fit of Eq. (15) gave $a = 0.0150$ (two-way 95% CI 0.0109–0.0191; $t_{41} = 7.36$, two-sided $p = 5.1 \times 10^{-9}$) and $b = 0.560$ (0.430–0.690) (Table 5). The primary test (E1) rejects $a = 0$, and the Holm-adjusted p over the two tests of the plan is 1.0×10^{-8}. The secondary calibrations agree. Clustering by state alone or by week alone gave intervals of 0.0125–0.0174 and 0.0112–0.0188. The week-cluster wild bootstrap-t with the null imposed gave $p = 0.0001$, the smallest value attainable with 9,999 draws, at every clamp of the bootstrap losses from 10^{-9} to 10^{-3} and 0.0003 without a clamp (Table S12). None of the 1,999 calendar-week block-bootstrap draws gave $a \leq 0$ (block-bootstrap percentile interval 0.0107–0.0192).

The lower limit of a stayed above zero in every sensitivity analysis of the plan, so the claim meets the robustness rule (Fig. 2). Point estimates ranged from 0.0117 to 0.0202 over 17 specifications, and the smallest lower limit, 0.0082, came from leaving out the 2025–26 season. The generation interval moved a most: a mean of 3.2 days gave 0.0117 and a mean of 4.0 days gave 0.0186, because the transform (14) rescales all deviations together. Raising the count floor lowered a slightly, to 0.0130 (0.0088–0.0172) at $C_t \geq 50$ on 828 rows. The fixed calendar windows gave the widest interval, 0.0092–0.0311, on 1,263 rows from three seasons. Leaving out one calendar week at a time gave estimates of 0.0136–0.0156 with a smallest lower limit of 0.0100. Count-independent dispersion is therefore present in these data under every specification that the plan fixed, provided the decline with count is inverse; Section 5.2 examines that proviso.

5.2 How well the affine form fits

The affine form describes the decline of the loss with count only approximately (Table S10; Fig. 3). Across deciles of the current-week count the mean loss fell from 0.0684 to 0.0110, and the affine prediction lay outside the block-bootstrap interval of the observed mean in three of ten deciles: above it in the lowest decile (0.0906 against 0.0538–0.0854) and in the highest (0.0158 against 0.0079–0.0137), and below it in the fifth (0.0296 against 0.0310–0.0501). A fit $a + bC^{-\gamma}$ with a free exponent lowered the quasi-deviance by 10.67, and the gain exceeded 2 in 94.6% of calendar-week block-bootstrap draws (Table 6). Its exponent was $\gamma = 0.617$ (week bootstrap 0.355–0.870; state bootstrap 0.438–0.862), so the inverse-count decline $\gamma = 1$ lies outside both intervals. Its intercept was 0.0089, and the week-bootstrap interval reached zero (-0.0014 to 0.0137; 4.0% of draws at or below zero), whereas the state-bootstrap interval did not (0.0034–0.0144). Under the reading rule fixed in Amendment 2 the floor therefore does not survive a free exponent. Within the observed counts a slower decline with a smaller constant and an inverse decline with a larger constant describe the data almost equally well, and the intercept of either model is an extrapolation beyond the largest counts.

The rows with the largest counts give the floor without a model for the decline (Table 6; Fig. 4). Among the 175 state-weeks at or above the 90th percentile of the count ($C_t \geq 389$, mean 882) the mean loss was 0.0110 (week bootstrap

Table 5 Count-independent intercept a: primary test of the analysis plan, secondary calibrations and sensitivity analyses

Analysis (rows)	a	95% CI	t	p (two-sided)	b/a	b/b_P
E1, primary: two-way CGM sandwich, t_{41}						
All rows (1748)	0.0150	0.0109–0.0191	7.36	$< 10^{-4}$	37.4	1.257
State-cluster sandwich	0.0150	0.0125–0.0174	12.23	$< 10^{-4}$	–	–
Week-cluster sandwich	0.0150	0.0112–0.0188	7.94	$< 10^{-4}$	–	–
Week wild bootstrap-t, $a = 0$ imposed (B=9,999)	–	–	–	0.0001	–	–
Calendar-week block bootstrap (B=1,999)	–	0.0107–0.0192	–	–	23.4–60.5	0.97–1.59
Sensitivity analyses (two-way sandwich)						
S-a1 GI mean 3.2 (1748)	0.0117	0.0085–0.0149	7.44	$< 10^{-4}$	37.6	1.259
S-a1 GI mean 4.0 (1748)	0.0186	0.0134–0.0237	7.27	$< 10^{-4}$	37.5	1.259
S-a2 GI SD 1.2 (1748)	0.0154	0.0111–0.0197	7.27	$< 10^{-4}$	37.6	1.278
S-a2 GI SD 2.0 (1748)	0.0144	0.0105–0.0183	7.46	$< 10^{-4}$	37.4	1.234
S-a3 count floor 10 (1600)	0.0140	0.0100–0.0180	7.07	$< 10^{-4}$	46.1	1.461
S-a3 count floor 20 (1268)	0.0136	0.0096–0.0176	6.82	$< 10^{-4}$	48.6	1.529
S-a3 count floor 50 (828)	0.0130	0.0088–0.0172	6.21	$< 10^{-4}$	51.5	1.591
S-a4 calendar windows (1263)	0.0202	0.0092–0.0311	3.78	0.0008	25.6	1.140
S-a5 threshold 1.02 (1748)	0.0150	0.0109–0.0191	7.36	$< 10^{-4}$	37.4	1.257
S-a5 threshold 1.1 (1564)	0.0141	0.0105–0.0176	8.00	$< 10^{-4}$	38.6	1.255
S-a6 exclude pause (1554)	0.0139	0.0103–0.0176	7.76	$< 10^{-4}$	41.5	1.331
S-a7 drop 2022-23 (1383)	0.0160	0.0105–0.0214	5.97	$< 10^{-4}$	31.0	1.101
S-a7 drop 2023-24 (1249)	0.0163	0.0122–0.0204	8.08	$< 10^{-4}$	33.3	1.240
S-a7 drop 2024-25 (1392)	0.0152	0.0100–0.0204	5.89	$< 10^{-4}$	36.6	1.230
S-a7 drop 2025-26 (1220)	0.0119	0.0082–0.0156	6.57	$< 10^{-4}$	55.5	1.488
S-a8 coverage $\geq$ 90% (1358)	0.0159	0.0110–0.0207	6.59	$< 10^{-4}$	31.0	1.104
S-a9 leave one week out (42 fits)	0.0136–0.0156	min lower 0.0100	–	max $< 10^{-4}$	–	1.188–1.306

Robustness rule (locked analysis plan): verdict *robust*; analyses with a lower limit ≤ 0: none. S-a5 with threshold 1.02 reproduces the primary windows. Block-bootstrap columns give percentile intervals.




Figure 2 Count-independent intercept a of the affine dispersion model for R_A under the primary analysis (E1) and sensitivity analyses S-a1 to S-a9, with two-way cluster-robust 95% confidence intervals. The dotted line marks the primary estimate and the solid line marks zero. For S-a9 the grey bar spans the 42 leave-one-week-out point estimates and the tick marks their smallest lower limit. The within-week intercept (E3, block-bootstrap interval) is a different estimand and is shown for comparison.




Figure 3 Loss by decile of the current-week count. (A) Mean relative squared deviation L against mean count C_t on log scales, with calendar-week block-bootstrap 95% intervals (1,999 draws). Lines show the fitted affine ($a+b/C$), power ($cC^{-\gamma}$) and three-parameter ($a + bC^{-\gamma}$) curves. (B) Ratio of the observed mean loss to the prediction of each model by decile; the dotted line marks 1.

Table 6 The floor under a free exponent and without a model

Quantity	Estimate	Week bootstrap	State bootstrap
Affine fit $a + b/C$ (primary, for reference)			
a (two-way interval)	0.0150	0.0109–0.0191	–
Three-parameter fit $a + bC^{-\gamma}$			
a	0.0089	−0.0014 to 0.0137	0.0034–0.0144
b	0.228	0.130–0.436	–
γ	0.617	0.355–0.870	0.438–0.862
Share of draws with $a \leq 0$	–	0.040	0.006
Quasi-deviance gain over affine	10.67	share $>$ 2: 0.946	–
Model-free floor among rows with C_t at or above a quantile			
P75 ($C_t \geq 152$, $n = 438$, mean C_t 497): mean L	0.0157	0.0117–0.0191	–
mean $L - \hat{b}/C_t$	0.0139	0.0096–0.0176	–
mean $L - b_P/C_t$	0.0142	0.0102–0.0177	–
relative SD $(L - \hat{b}/C_t)^{1/2}$ (%)	11.8	9.8–13.2	–
P90 ($C_t \geq 389$, $n = 175$, mean C_t 882): mean L	0.0110	0.0078–0.0139	–
mean $L - \hat{b}/C_t$	0.0102	0.0069–0.0132	–
mean $L - b_P/C_t$	0.0104	0.0071–0.0132	–
relative SD $(L - \hat{b}/C_t)^{1/2}$ (%)	10.1	8.3–11.5	–
P95 ($C_t \geq 699$, $n = 90$, mean C_t 1238): mean L	0.0101	0.0071–0.0128	–
mean $L - \hat{b}/C_t$	0.0096	0.0064–0.0123	–
mean $L - b_P/C_t$	0.0097	0.0066–0.0124	–
relative SD $(L - \hat{b}/C_t)^{1/2}$ (%)	9.8	8.0–11.1	–

Week bootstrap: calendar-week blocks (1,999 draws); state bootstrap: states resampled (999 draws). Model-free floor: mean loss of the rows above the threshold minus the mean counting term, with $\hat{b}$ the fitted slope or b_P the Poisson reference; intervals from the calendar-week block bootstrap with $\hat{b}$ refitted in each draw. Reading rule of Amendment 2: the floor survives a free exponent if the week-bootstrap interval of a excludes 0 (here: no).

0.0078–0.0139), a relative root mean squared deviation of 10.5% (8.8–11.8%). Subtracting the counting term with the fitted slope left 0.0102 (0.0069–0.0132), a relative standard deviation of 10.1% (8.3–11.5%), and subtracting it with the Poisson reference left 0.0104 (0.0071–0.0132). At the 75th and 95th percentiles the net relative standard deviation was 11.8% (9.8–13.2%) and 9.8% (8.0–11.1%). At the largest observed counts, therefore, a count-independent relative error of about 10% remains, and all of these intervals exclude zero. The model-free value at the 90th percentile lies between the three-parameter intercept and the affine one.

5.3 What more cases buy

The observed relative error fell with count but levelled off at large counts (Table S10). The relative root mean squared deviation of R_A was 26.2% in the lowest count decile (mean 7.8 admissions a week), 20.1% in the fifth (mean 38.5), 14.9% in the seventh (mean 93.5) and 10.5% in the highest (mean 882). A ninefold increase in the mean count from the seventh to the tenth decile thus lowered the relative error by about four percentage points. The affine model gives smooth values over the same range, 26.6% at 10 admissions, 17.3% at 37 and 14.3% at 100 (Table 7). Its extrapolation to 1,000 admissions, 12.5%, exceeds the observed relative error of the highest decile, so statements about large counts here rest on the model-free floor of Section 5.2. The crossover count, at which the two affine terms are equal, was $b/a = 37.4$ admissions a week (two-way 24.1–58.3; block bootstrap 23.4–60.5). It rose to 46.1, 48.6 and 51.5 at count floors of 10, 20 and 50 and lay between 25.6 and 55.5 across the other sensitivity analyses, so its order of magnitude, a few tens of




Figure 4 Estimates of the count-independent component on the relative-variance scale, with the corresponding relative standard deviation on the upper axis. Rows show the affine intercept (two-way cluster-robust 95% interval), the same intercept in the largest-population tertile (two-way interval) and the three-parameter intercept with calendar-week and state bootstrap 95% intervals. The remaining rows show the model-free floor (mean loss minus the fitted counting term) among rows at or above the 75th, 90th and 95th percentiles of the count, with calendar-week block-bootstrap 95% intervals. The solid line marks zero.

Table 7 What more cases buy: predicted relative squared error $a + b/C$ by count, crossover, and the within-week intercept

C	a	b/C	$a + b/C$	95% CI (sandwich)	95% CI (block bootstrap)	Share from a (%)	Relative RMSE (%)
10	0.0150	0.0560	0.0710	0.0595–0.0824	0.0604–0.0829	21.1	26.6
37	0.0150	0.0151	0.0301	0.0263–0.0339	0.0264–0.0333	49.7	17.3
100	0.0150	0.0056	0.0206	0.0170–0.0242	0.0168–0.0239	72.8	14.3
1,000	0.0150	0.0006	0.0155	0.0115–0.0196	0.0113–0.0196	96.4	12.5
Crossover $C^{\times} = b/a$ = 37.4; sandwich (log-delta) 24.1–58.3; block bootstrap 23.4–60.5							
Within-week intercept (E3, estimand change; block bootstrap 95% CI)							
Leave-state-out week centre (primary)			0.0102	0.0078–0.0126	share of a removed 32.0% (10.2–47.8)		
Week centre incl. own row			0.0098	0.0075–0.0121	share of a removed 34.7% (13.8–49.8)		
Week centre and scale			0.0086	0.0066–0.0107	share of a removed 42.6% (22.0–57.0)		
Season centre			0.0147	0.0101–0.0184	share of a removed 1.6% (−6.8 to 19.1)		
Pooled centre (E1 loss)			0.0150	0.0107–0.0191	share of a removed 0.0% (0.0–0.0)		

weekly admissions, is stable while its value is not. It is an in-sample balance point for this panel, not a sample-size rule for surveillance.

These values pool jurisdictions whose count-independent component differs (Section 5.4). Separate affine fits by population tertile gave predicted relative errors at 37, 100 and 390 admissions of 18.3, 15.6 and 14.2% in the smallest-population tertile, 17.9, 15.5 and 14.4% in the middle tertile and 16.2, 13.5 and 12.1% in the largest. What additional cases buy should be judged against the component of the jurisdiction concerned.

5.4 Jurisdiction size

The intercept fell with state population (Table 8; Fig. 5). In the Amendment 2 model with an intercept linear in standardised log population and a common slope, the population coefficient was $a_1 = -0.0054$ per standard deviation (two-way -0.0085 to -0.0023; $t_{41} = -3.51$, $p = 0.0011$), and its state-bootstrap and week-bootstrap intervals also excluded zero. The fitted intercept was 0.0273 at the 10th percentile of population (0.92 million) and 0.0137 at the 90th (12.7 million), a ratio of 0.50 (state bootstrap 0.41–0.61); a log-link intercept gave a ratio of 0.44 (state bootstrap 0.32–0.60). The counting slope did not change detectably with population (coefficient -0.015; two-way -0.124 to 0.095). Separate fits by tertile gave intercepts of 0.0189 (two-way 0.0145–0.0233), 0.0195 (0.0125–0.0264) and 0.0135 (0.0090–0.0180), and with a common slope the ratio of the largest-population to the smallest-population tertile was 0.60 (state bootstrap 0.42–0.82).

Population and count go together in this panel: the median count was 20, 37 and 106 admissions in the three tertiles. Because the affine form overpredicts the loss at the largest counts (Section 5.2), part of the population gradient may reflect that misfit rather than a property of the jurisdictions. Either way the pooled intercept of 0.0150 is an average over jurisdictions whose count-independent component differs about twofold, and the count beyond which more cases help little is specific to the jurisdiction.

Table 8 Dependence of the count-independent intercept on state population

Quantity	Estimate	Two-way	State bootstrap	Week bootstrap
Linear intercept $a_0 + a_1 z$, common slope b (primary)				
a_0	0.0197	0.0136–0.0258	–	–
a_1 (per SD of log population)	−0.0054	−0.0085 to −0.0023	−0.0078 to −0.0034	−0.0085 to −0.0023
a at population P10 / P90	0.0273 / 0.0137	–	–	–
Ratio a(P90) / a(P10)	0.501	–	0.408–0.609	–
b	0.451	–	–	–
Log-link intercept $\exp(\alpha_0 + \theta z)$ (secondary)				
θ	−0.327	–	−0.445 to −0.205	−0.438 to −0.183
Ratio a(P90) / a(P10)	0.436	–	0.324–0.595	–
Intercept and slope both linear in z (secondary)				
a_1	−0.0052	−0.0083 to −0.0022	–	–
b_1	−0.015	−0.124 to 0.095	–	–
Population tertiles, separate affine fits (two-way interval, t with $G - 1$ df)				
Tertile 1 (17 states, 0.59–2.9 M, 399 rows): a	0.0189	0.0145–0.0233	–	–
b (median C_t = 20)	0.534	0.350–0.719	–	–
Tertile 2 (17 states, 2.96–6.2 M, 638 rows): a	0.0195	0.0125–0.0264	–	–
b (median C_t = 37)	0.462	0.299–0.624	–	–
Tertile 3 (17 states, 6.91–39.3 M, 711 rows): a	0.0135	0.0090–0.0180	–	–
b (median C_t = 106)	0.475	0.218–0.732	–	–
Population tertiles, common slope				
Tertile 1 intercept	0.0213	–	0.0160–0.0303	–
Tertile 2 intercept	0.0191	–	0.0149–0.0240	–
Tertile 3 intercept	0.0128	–	0.0105–0.0158	–
Ratio tertile 3 / tertile 1	0.601	–	0.421–0.818	–

z: log of the mean 2022–2025 resident population (Census Vintage 2025), standardised over the 51 jurisdictions (P10 = 0.92 M, P90 = 12.7 M). Primary test of $a_1 = 0$: two-way sandwich, t with 41 df, $t = -3.51$, $p = 0.0011$. State bootstrap: states resampled (1,999 draws; 999 for the tertile fits); week bootstrap: calendar-week blocks (1,999 draws). All intervals 95%.




Figure 5 Count-independent intercept and state population. (A) Fitted intercept against mean resident population for the linear and log-link population models, with separate affine fits by population tertile (filled points, two-way cluster-robust 95% intervals) and tertile intercepts under a common slope (open points). Dotted lines mark the 10th and 90th percentiles of population. (B) Ratios of intercepts with state-bootstrap 95% intervals: 90th to 10th percentile of population under the linear and log-link models, and largest to smallest tertile under a common slope.

Table 9 Implied relative SD of the reproduction number across simulated mechanisms (a grid, not an estimate)

Mechanism family	Rows	Point range (%)	Interval envelope (%)	Reproduce b/b_P	Post hoc laws
original generators (fixed field)	9	8.6–18.6	6.9–22.7	4	1
fresh-draw generator	5	9.2–12.1	7.1–14.7	4	3
$\rho \times k$ grid (S1 field)	40	10.0–12.3	7.5–14.5	3	0
daily renewal (two generations per week)	12	9.8–14.1	7.8–17.2	2	0
generation-interval variant ($r = 1$)	4	10.8–13.6	9.2–15.4	–	0
within-week estimand (E3, $r = 1$)	1	10.1–10.1	8.8–11.2	–	0
All rows	71	8.6–18.6	6.9–22.7	–	–
Pre-specified laws	67	8.6–18.6	6.9–22.7	–	–
Rows reproducing b/b_P	13	8.6–13.9	6.9–17.1	–	–

Implied SD $= \sqrt{a_{\text{obs}}/r}$, $r = E(\hat{a})/v_R$ of each mechanism; intervals combine the two-way CI of a with the MC 2.5–97.5% range of r. The transmission share of a is not identified. Renewal rows appear twice (following-week R and two-week mean R).




Figure 6 Relative standard deviation of the reproduction number implied by the observed intercept, $(a/r)^{1/2}$, for each simulated mechanism with recovery ratio $r = \mathrm{E}(\hat{a})/v_R$, grouped by mechanism family. Bars combine the two-way 95% interval of a with the Monte Carlo 2.5–97.5% range of r. Red points mark mechanisms whose Monte Carlo range of b/b_P contains the observed value, grey points those that do not, and blue points rows with $r = 1$ for which no slope check applies; triangles mark laws defined after the data had been seen. The shaded band spans the point values of all rows. Daily-renewal rows appear twice, against the following-week and the two-week mean reproduction number.

5.5 What the floor contains

Removing common national variation by calendar week lowered the intercept to 0.0102 (block bootstrap 0.0078–0.0126) with the leave-state-out week centre, a reduction of 32.0% (10.2–47.8) (Table 7). With the week centre including the own row the reduction was 34.7% (13.8–49.8), and with week-specific centre and scale it was 42.6% (22.0–57.0). Centring by season removed 1.6% (-6.8 to 19.1). About a third of the floor is thus shared by states in the same week, and about two thirds varies between states within a week. The within-week intercept stays above zero under every centring.

The floor reflects transmission and reporting variation, and its transmission share is not identified. The square root of a, 12.2%, is the relative standard deviation of the non-counting variation that the intercept captures around the pooled mean, which can fall short of the sum of the two sources (Section 5.6). Converting a into a transmission standard deviation requires a mechanism, and across 71 simulated mechanisms the implied relative standard deviation of the reproduction number ranged from 8.6% to 18.6%, with propagated intervals from 6.9% to 22.7% (Table 9; Fig. 6). Restricted to the 13 mechanisms that reproduce the observed counting slope, the range was 8.6–13.9%. Reporting shocks make the intercept overstate transmission variance, and heterogeneity that shrinks with count makes it understate it, which is why the range is wide in both directions.

The mapping itself depends on the time scale of transmission (SC1). Under a daily renewal process with the primary generation interval, about two generations fall within a week, and the intercept recovered 75–78% of the variance of the weekly reproduction number, outside the range of 0.8–1.25 fixed in the plan. Against the mean of two consecutive weekly values it recovered 150–156%. The one-generation map of Section 2 therefore does not carry over unchanged to weekly hospital counts, and the implied standard deviations above inherit this dependence.

Table 10 Bias of the intercept relative to the total non-counting variance

Generator	v_H share (%)	$E(\hat{a})$	v_R	v_H	Rel. bias (%)	$E(\hat{a})/v_R$	w_H
FD-S1	0.0	0.0149	0.0146	0.0000	1.4 (0.5–2.3)	1.014	–
FD-S8	50.9	0.0150	0.0086	0.0089	−14.2 (−15.1 to −13.3)	1.749	0.708 (0.687–0.729)
FD-S9, 25†	50.9	0.0151	0.0086	0.0089	−13.8 (−14.8 to −12.8)	1.754	0.715 (0.693–0.736)
FD-S9, 50†	50.9	0.0150	0.0086	0.0089	−14.0 (−15.1 to −12.9)	1.753	0.711 (0.689–0.734)
FD-S9, 75†	50.8	0.0148	0.0086	0.0089	−14.9 (−16.1 to −13.7)	1.731	0.693 (0.669–0.718)
FD-S8, share 100	0.0	0.0154	0.0217	0.0000	−29.3 (−30.1 to −28.4)	0.707	–
FD-S8, share 75	25.7	0.0152	0.0144	0.0050	−22.1 (−23.1 to −21.2)	1.049	0.100 (0.056–0.147)
FD-S8, share 25	75.6	0.0148	0.0039	0.0120	−6.3 (−7.2 to −5.4)	3.840	0.912 (0.900–0.924)

Fresh-draw generators of the first revision (200 panels each) and an FD-S8 sweep over the transmission share of the calibrated variance (shares 1.0, 0.75, 0.25; 0.5 is FD-S8 itself). v_R, v_H: true relative variance of transmission and of the reporting shock on the kept rows; relative bias $= E(\hat{a})/(v_R + v_H) - 1$; w_H: weight on v_H solving $E(\hat{a}) = v_R + w_H v_H$. Intervals: panel bootstrap (2,000 draws). † post hoc law. Wording rule of Amendment 2: if every interval excludes 0, the manuscript says $\hat{a}$ reflects, not estimates, the total (outcome: reflects (not an estimate of the total)). Mechanism-conditional rescaling a_{obs}/t_g would range 0.0160–0.0192; not used.

Table 11 External reporting data: participation and backfill

Quantity	Value	95% interval
Hospital participation (NHSN, % of hospitals reporting)		
Mean (min; P10) participation	92.9% (67.7; 86.9)	–
Rows with full participation in both weeks	4.1%	–
Mean week-to-week change	0.78 pp	–
a, observed counts	0.0150	–
a, participation-adjusted counts	0.0147	two-way SE 0.0020
Difference (relative)	−0.00029 (−2.0%)	−0.00086 to 0.00027
Backfill (Delphi Epidata issue history, archive from issue 2024w47)		
Rows of R_A in the archive; latest issue equals the vintage	1,748; 99.1%	–
Rows with a provisional first issue (lag $\leq$ 2 weeks)	661 (15 weeks, 2024-11-09 to 2025-12-27)	–
First issue equals latest (current week)	29.2%	–
Revised up / down	65.8% / 5.0%	–
Mean relative revision (latest − first) / first	11.0%	–
a, latest issue (654 rows)	0.0143	0.0096–0.0183
a, first issue	0.0227	0.0133–0.0298
Difference first − latest	0.0084	0.0016–0.0146
v_{rev} (revision variance of x)	0.0107	–

Participation adjustment: both counts scaled by 100/participation, x recomputed, the affine fit repeated with the observed C_t as the count; assumes unreported hospitals admit as many patients per hospital as reporting ones. Backfill: rows whose two weeks both have a provisional first issue; fits restricted to rows with first and latest counts ≥ 5. Intervals: calendar-week block bootstrap, 1,999 draws. The backfill comparison bounds the provisional-versus-settled component only; the residual reporting error of the settled vintage is not identified from public data.

5.6 How the intercept relates to its sources

Under fresh-draw generators the intercept tracked transmission variance when there was no reporting shock and fell short of the total when there was (Table 10). With independent heterogeneity (FD-S1) its relative bias against $v_R + v_H$ was $+1.4$% (panel bootstrap 0.5–2.3%). Under FD-S8, in which about half of the calibrated variance is a reporting shock, it was -14.2% (-15.1 to -13.3%), and under the three post hoc variants FD-S9 it was -13.8 to -14.9%. Varying the transmission share of the FD-S8 variance moved the shortfall from 22.1% at a share of 0.75 to 6.3% at a share of 0.25, and with no reporting shock in that family the intercept recovered 0.707 of the transmission variance. The weight of the reporting variance in $\mathrm{E}(\hat{a}) = v_R + w_H v_H$ ranged from 0.10 to 0.91 across these laws. The direction of the bias is downward under every law with reporting shocks, and its size depends on the share and mechanism, neither of which the data identify. The intercept is therefore not rescaled, which would give 0.0160–0.0192 depending on the law, and it is read as reflecting transmission and reporting variation rather than as an estimate of their sum.

5.7 External reporting data

Two external sources bound parts of the reporting component (Table 11). Hospital participation averaged 92.9% of hospitals per jurisdiction-week (minimum 67.7%) and changed by 0.78 percentage points between consecutive weeks on average. Scaling both counts by participation lowered a by 0.00029, or 2.0% (week bootstrap -0.00086 to 0.00027), so changing participation explains little of the intercept. Revisions explain more. In the Delphi Epidata archive of issues [45], which starts with week 47 of 2024, 661 rows had a provisional first issue in both weeks. The first issue equalled the latest in 29.2% of them, was revised upward in 65.8% and downward in 5.0%, and the mean relative revision was 11.0%. On the 654 rows with both counts at least 5, a was 0.0227 (block bootstrap 0.0133–0.0298) with first-issue counts and 0.0143 (0.0096–0.0183) with latest-issue counts, a difference of 0.0084 (0.0016–0.0146). Counts available in real time thus carry a count-independent component about 1.6 times that of settled counts. Neither source measures the reporting error that remains in settled counts, so the transmission share of a stays unidentified.

Table 12 Counting slope against three Poisson references

Reference	b_P	b/b_P	95% interval	Change vs. shipped (%)
plug-in, as shipped	0.4455	1.257	0.97–1.55	−0.0
plug-in, mean-normalised	0.4199	1.334	1.02–1.64	6.1
simulated exact (Poisson-only panels)	0.3977	1.408	1.08–1.73	12.0

$\hat{b} = 0.560$, two-way 95% interval 0.430–0.690; intervals of b/b_P divide it by the reference (b_P fixed). Simulated exact: 400 panels with Poisson counts around the observed means, same transform, row filter ($C_t \geq 5$) and affine fit; pooled slope 0.3977 (per-panel 0.370–0.429), pooled intercept 1.7e-05.

Table 13 Reporting probability ρ × offspring dispersion k: intercept recovery $E(\hat{a})/v_R$ and counting slope b/b_P (100 panels per cell)

ρ	k=0.1	k=0.3	k=1	k=3	Poisson
0.001*	1.00 / 0.88	0.99 / 0.87	1.00 / 0.86	1.00 / 0.86	1.00 / 0.87
0.003*	1.01 / 0.87	1.01 / 0.86	1.01 / 0.85	1.01 / 0.86	1.01 / 0.85
0.01*	1.00 / 0.91	1.01 / 0.87	1.00 / 0.86	1.01 / 0.85	1.00 / 0.85
0.03	1.00 / 1.00	1.02 / 0.89	1.01 / 0.86	1.00 / 0.85	1.00 / 0.86
0.1	1.02 / 1.29✓	1.01 / 0.97	1.00 / 0.86	1.00 / 0.83	1.01 / 0.80
0.25	1.06 / 1.81	1.02 / 1.14✓	1.01 / 0.87	1.00 / 0.79	1.00 / 0.74
0.5	1.18 / 2.41	1.04 / 1.41✓	1.01 / 0.87	1.01 / 0.71	1.01 / 0.63
1	1.49 / 2.90	1.11 / 1.81	1.03 / 0.87	1.02 / 0.54	1.02 / 0.36

Daily renewal, two generations per week (a/v_R / $a/v_{\bar{R}}$ / b/b_P)
ρ=0.003, Poisson: 0.76 / 1.53 / 0.85
ρ=0.003, NB k=0.3: 0.75 / 1.51 / 0.87
ρ=0.01, Poisson: 0.75 / 1.50 / 0.86
ρ=0.01, NB k=0.3: 0.75 / 1.51 / 0.91
ρ=0.1, Poisson: 0.75 / 1.52 / 0.91
ρ=0.1, NB k=0.3: 0.78 / 1.56 / 1.38

Cells give $E(\hat{a})/v_R$ / mean b/b_P. * Assumed range for ρ (0.001–0.01; CDC burden estimates, not estimated). ✓ MC 2.5–97.5% range of b/b_P contains the observed 1.257. SC2 (recovery within 0.9–1.1 in every cell): not met; SC3 (b/b_P within 0.9–1.1 at $\rho \leq 0.01$): not met; SC1 (renewal a/v_R within 0.8–1.25): not met.

5.8 Counting slope, reporting and offspring dispersion

The counting slope was 1.257 times the plug-in independent-Poisson reference b_P (block bootstrap 0.97–1.59). That reference normalises every row by the squared pooled mean; normalising each row by its own ratio to the pooled mean lowers b_P from 0.4455 to 0.4199 and raises the ratio by 6.1% to 1.334 (two-way, reference fixed, 1.02–1.64). An exact reference simulated from Poisson counts on the same design gives 0.3977 and a ratio of 1.408 (1.08–1.73) (Table 12). Whether counting variation exceeds independent Poisson noise therefore depends on the reference: under the plug-in reference used in the plan it is not shown, and under the other two the lower limit lies just above 1. The comparisons with simulated slopes below use the plug-in reference throughout, so they are not affected. Against the branching-Poisson reference of Eq. (16) the point ratio would be 1.46 at $\rho = 0.25$ and 2.84 at $\rho = 1$ (Table S1), but the slope does not identify ρ. Within-state fits were less precise: a state fixed-effects fit gave 0.85 (0.24–1.47) times b_P and a week fixed-effects fit 1.42 (0.96–1.89) (Table S6).

The reporting-by-offspring grid shows what the slope can and cannot reveal (Table 13; Fig. 7). Intercept recovery stayed within 0.9–1.1 at $\rho \leq 0.25$ for every offspring law and at all ρ for $k \geq 1$, but rose to 1.18 at $\rho = 0.5$ and 1.49 at $\rho = 1$ with $k = 0.1$, and to 1.11 at $\rho = 1$ with $k = 0.3$; SC2 therefore fails, in cells outside the assumed range of ρ. In the assumed range, $\rho \leq 0.01$, the expected slope ratio was 0.85–0.91 whatever the offspring law, below 1 rather than within 0.9–1.1 as SC3 required. Offspring dispersion cannot be read from the counting slope at realistic reporting probabilities, and no cell in that range reproduces the observed 1.257. Only three of 40 cells did, all with $\rho \geq 0.1$ and $k \leq 0.3$. The observed slope is thus compatible with strong overdispersion only at reporting probabilities far above those of hospital surveillance, or with counting variation that the simulated mechanisms do not contain.

5.9 Season intercepts

No difference between season intercepts was detected under the primary calibration (calendar-week permutation $p = 0.102$; power 0.47 at the observed pattern). The Wald statistic was $W_{\mathrm{CGM}} = 17.8$ on 3 degrees of freedom, and its calendar-week permutation p was 0.102 (Monte Carlo standard error 0.003) (Table 14; Fig. 8B). Season statements below are therefore exploratory. The other calibrations disagree with one another. The asymptotic χ^2_3 reference gave $p = 0.0005$, the week-cluster wild bootstrap 0.011, the state-cluster wild bootstrap 0.013, the within-state permutation 0.045, generator-based references 0.005–0.055, the bootstrap-covariance statistic 0.138 and circular shifts of the season sequence 0.405. The state-level calibrations treat states as independent clusters although the season label varies only between weeks, and the generator-based ones depend on the simulated mechanism. Under the primary calibration, excluding the 2025 data-channel interruption gave $p = 0.261$, restricting to rows with at least 90% reporting coverage gave 0.081, and the fixed calendar windows for three seasons gave 0.031.

The 2023–24 intercept was the lowest, 0.0071 (two-way sandwich on the log scale, 0.0050–0.0100), against 0.0154–0.0180 in the other seasons (Table 15; Fig. 8A). The ratio of the highest to the lowest intercept was 2.54 (season-stratified block bootstrap 1.75–4.35), an estimate that is biased upward. A contrast of 2023–24 against the other seasons, chosen




Figure 7 Reporting probability ρ by offspring dispersion k, with independent transmission heterogeneity and 100 synthetic panels per cell on the real design. (A) Intercept recovery $\mathrm{E}(\hat{a})/v_R$. (B) Mean counting slope relative to the independent-Poisson reference, b/b_{P}; asterisks mark cells whose Monte Carlo 2.5–97.5% range contains the observed 1.257. The outlined rows mark the assumed range of ρ for hospital admissions (0.001–0.01).

Table 14 Equal-season-intercept test (W_{CGM} = 17.80): primary calibration of the locked plan and every secondary calibration

Calibration	p (MCSE)	Draws
Calendar-week permutation (primary, locked plan)	0.102 (0.003)	9,999
Asymptotic χ^2_3, sandwich W	0.0005	–
Asymptotic χ^2_3, bootstrap-covariance W	0.138	–
Week-cluster wild bootstrap (Webb, null imposed)	0.011 (0.001)	9,999
Circular shifts of the season sequence (exact, 42)	0.405	42
State-cluster wild bootstrap	0.013 (0.002)	2,000
Within-state permutation	0.045 (0.005)	2,000
Generator S1 (fixed transmission field)	0.013 (0.006)	400
Generator S6 (fixed transmission field)	0.045 (0.010)	400
Generator S8 (fixed transmission field)	0.028 (0.008)	400
Generator S9 (fixed transmission field; post hoc law)	0.005 (0.004)	400
Generator FD-S1 (redrawn per replicate)	0.055 (0.016)	200
Generator FD-S8 (redrawn per replicate)	0.050 (0.015)	200
Generator FD-S9-25 (redrawn per replicate; post hoc law)	0.035 (0.013)	200
Generator FD-S9-50 (redrawn per replicate; post hoc law)	0.045 (0.015)	200
Generator FD-S9-75 (redrawn per replicate; post hoc law)	0.035 (0.013)	200
Data sensitivities, primary calibration (calendar-week permutation, B=9,999)		
S-a6 exclude pause: W=11.04, 3 df, 36 weeks	0.261 (0.004)	9,999
S-a8 coverage $\geq$ 90%: W=16.77, 3 df, 42 weeks	0.081 (0.003)	9,999
S-a4 calendar windows: W=15.55, 2 df, 29 weeks	0.031 (0.002)	9,999

Decision rule (locked analysis plan): significant only if the primary $p < 0.05$. Verdict: not significant: season statements are exploratory/descriptive. Holm-adjusted over the two claims: E1 $< 10^{-4}$, season 0.102. Shipped generator calibrations hold the transmission field fixed; S9 and FD-S9 laws were specified after the data were seen.

after the data had been seen, gave a ratio of 0.43 (two-way sandwich on the log scale, 0.29–0.65) and is not tested. Power at the observed pattern was 0.47 (Monte Carlo standard error 0.025), and 80% power would need a highest-to-lowest ratio of about 4.5 (Fig. 8C). With equal season intercepts the test rejected in 0.0625 of simulated panels (Monte Carlo standard error 0.012), about one standard error above the nominal 0.05. This power curve uses a loss-level statistic whose permutation critical value, 17.65, is lower than that of the primary test, so it is probably optimistic. The absence of a detected difference is therefore weak evidence that the season intercepts are equal. With four seasons the design can detect only large differences, and the between-season variance itself cannot be estimated precisely (Appendix B.3).

Any season difference would be a difference in the combined transmission and reporting variation that the intercept reflects. Mean hospital reporting coverage of the sample rows was 90.9–94.2% in every season (Table 15), but coverage does not measure the reporting probability per infection, and scaling counts by participation changed a by 2.0% (Section 5.7). Reporting regimes did differ outside the analysis windows: median participation in summer 2024, when reporting was voluntary, was 45.6% against 92.3–94.2% in the other summers (Table S7).

5.10 Validity of the intervals

Under generators that redraw the heterogeneity field in every replicate, the two-way interval covered the mean of $\hat{a}$ in 0.920–0.975 of panels, which meets SC4 (Table 16). Its mean standard error was 0.92–1.07 times the Monte Carlo standard deviation. Under mechanisms with reporting shocks the interval is centred on an estimator that falls short of the

Table 15 Season intercepts (reflecting transmission and reporting variation) with sandwich and season-stratified block-bootstrap intervals, and design-based power

Season	Weeks	Rows	a_s	95% CI (sandwich)	95% CI (bootstrap)	a_s/a (95% CI)	Coverage (%)
2022-23	9	365	0.0159	0.0112–0.0224	0.0102–0.0221	1.06 (0.68–1.64)	90.9
2023-24	12	499	0.0071	0.0050–0.0100	0.0050–0.0111	0.47 (0.34–0.71)	94.2
2024-25	8	356	0.0154	0.0092–0.0257	0.0084–0.0242	1.03 (0.59–1.55)	93.8
2025-26	13	528	0.0180	0.0108–0.0299	0.0090–0.0252	1.20 (0.71–1.56)	92.3

Highest/lowest intercept ratio 2.54 (bootstrap 1.75–4.35; biased upward)
Post hoc: 2023-24 vs geometric mean of the other seasons, ratio 0.43 (sandwich 0.29–0.65; bootstrap 0.31–0.72); no test
Design-based power at the observed pattern 0.47; minimum detectable highest/lowest ratio at 80% power 4.48

Coverage: mean hospital-reporting coverage of the sample rows. Voluntary reporting May–Oct 2024; CDC data-channel pause 3 October to 13 November 2025.




Figure 8 Season intercepts, exploratory. (A) Season intercepts a_s in a model with a common slope. Intervals are two-way sandwich intervals on the log scale and season-stratified calendar-week block-bootstrap intervals, and the dotted line marks the pooled a. Panel (B) shows the calendar-week permutation distribution of the cluster-robust Wald statistic under equal season intercepts (9,999 permutations), with the observed statistic and the 95th percentile of the null distribution. Panel (C) shows the power of a permutation test at the 5% level against the highest-to-lowest intercept ratio along the observed pattern. Error bars are binomial Monte Carlo errors, and MDR marks the minimum detectable ratio at 80% power.

Table 16 Fresh-draw generators (transmission field redrawn in every replicate, 200 panels each)

Generator	MC SD of $\hat{a}$	Observed SE / MC SD	Sandwich SE / MC SD	Coverage of $E(\hat{a})$	Coverage of true total	$E(\hat{a})/v_R$	b/b_P	Week-centring reduction (%)	Implied SD of R (%)
FD-S1	0.00098	2.08	1.02	0.945	0.975	1.014	0.842	0.1	12.1 (9.8–14.7)
FD-S8	0.00127	1.60	0.92	0.920	0.440	1.749	1.281	−0.8	9.2 (7.2–11.4)
FD-S9-25†	0.00130	1.56	0.94	0.925	0.520	1.754	1.298	8.3	9.2 (7.3–11.4)
FD-S9-50†	0.00132	1.53	1.03	0.940	0.570	1.753	1.297	18.4	9.2 (7.1–11.5)
FD-S9-75†	0.00147	1.39	1.07	0.975	0.590	1.731	1.310	26.9	9.3 (7.2–11.7)

† Post hoc law (national calendar-week share). SC5 (observed SE / MC SD in [0.67, 1.5] for some generator): met, only by a post hoc law; SC4 (coverage $\geq$ 0.90): met.

total (Section 5.6), and it covered the true total in only 0.44–0.59 of panels, against 0.975 under independent heterogeneity. The observed standard error of a was 1.53–2.08 times the Monte Carlo standard deviation under the fresh-draw laws of the plan and 1.39 under the national-share law with share 0.75, which was defined after the data had been seen. SC5 is therefore met only by a post hoc law. The two-way interval is wider than the Monte Carlo spread under every fresh-draw law, and the pigeonhole bootstrap standard deviation was 1.55–1.78 times it (Table S13), so the intervals reported here are conservative for the mean of $\hat{a}$ under these laws.

Generators built to reproduce the state structure of the panel did not close the gap between the variability of the data and of the simulated panels (Table S11). They combined a transmission field around a smoothed observed trend, with state-specific variance proportional to the population-dependent intercept or to state intercepts, clustered admissions and a reporting probability of 0.01 or 0.003. In five of the six cells the smoothed trend alone exceeded the transmission target, so the transmission standard deviation could not be calibrated. No cell reproduced all four tested features of the observed panel: the counting slope ratio fell inside the Monte Carlo range only without admission clustering, the 90th percentile of the count in five of six cells, the pooled-to-within-state slope ratio (T3) in none and the between-state spread of intercepts in all. Under these cells the two-way interval covered the mean of $\hat{a}$ in 0.96–1.00 of panels, the observed standard error was 1.63–2.31 times the Monte Carlo standard deviation, and the intercept captured 70–78% of the simulated non-counting total, which the interval covered in only 0.015–0.16 of panels. Which features of the data inflate its variability relative to these mechanisms remains unexplained. Simulated null distributions for the season test, which drew on generators with a fixed heterogeneity field, gave p values of 0.005–0.045; redrawing the field raised them to 0.035–0.055. These generator-based calibrations are not used as evidence for season differences.

6 Discussion

The theoretical part of the paper makes one distinction precise. With fixed, known reporting, exogenous exposure and conditionally independent counting errors, the risk for the current value is A/m and the risk for the realised mean of the observed periods is $A/(Tm)$. The process mean adds the variance of the realised mean, which more cases within the same periods do not reduce. The count at which counting error equals between-period variance is the half-weight point of the corresponding shrinkage estimator, as in small-area estimation and disease mapping [14, 15]. It belongs with its estimator, target and sampling experiment and is not a property of a pathogen. The efficiency statement holds within a class of conditionally unbiased estimators with a known denominator. Information from other features of the epidemic curve [6, 58], independent reporting measurements or prior structure changes what can be learned, and estimators that accept bias in exchange for lower variance lie outside that class. Appendix B gives the corresponding forecast weight, the allocation of a fixed budget between periods and cases, and a lower bound on the precision with which a few seasons can estimate between-season variation.

The application answers its primary question. US influenza hospital admissions in early-growth weeks carry a count-independent component of squared relative error: the primary test rejected $a = 0$, the lower limit stayed above zero in every sensitivity analysis of the plan, and about two thirds of the component remains after common calendar-week variation is removed. Its size is better read from the data than from the affine model, which fits the decline with count only approximately. Among the state-weeks with the largest counts the relative error of a weekly apparent reproduction number was about 10%, almost none of it counting noise, whereas in the lowest count decile it was 26%. Beyond a few tens of weekly admissions, more cases mainly buy precision for the current value of a statistic whose remaining spread is not counting noise. This is the empirical counterpart of the distinction between targets: a count-based precision statement describes the first component and is silent on the second. The component is not common to all jurisdictions. It was about half as large in the most populous states as in the least populous, so a single pooled intercept understates the error in small jurisdictions and overstates it in large ones.

What the count-independent component is cannot be settled from these data. Its transmission share is not identified (Proposition 2), and implied transmission standard deviations ranged from 8.6% to 18.6% across simulated mechanisms, a grid that depends on the assumed mechanism and on the number of generations per week. Under reporting shocks the intercept fell 6–22% short of the transmission-plus-reporting total, so it combines the two sources with unequal weights rather than adding them. The counting slope, its ratio to the independent-Poisson or branching-Poisson reference and the thinning response also cannot distinguish overdispersed offspring, under-reporting and heterogeneous transmission. At reporting probabilities typical of hospital admissions the expected slope ratio was below one whatever the offspring law. The observed ratio was 1.26 against the plug-in reference used throughout and 1.33–1.41 against two closer approximations, so whether counting variation exceeds independent Poisson noise depends on the reference. No difference between season intercepts was detected under the primary calibration ($p = 0.102$; power 0.47). The lower intercept of 2023–24 is a hypothesis for data with more seasons or with external reporting measurements, not a finding, and the power analysis shows that four seasons can detect only differences of about fourfold or more. The crossover count of about 37 admissions per week marks where counting noise stops dominating the cross-sectional dispersion of this statistic in the pooled panel, not where additional cases stop being useful for a particular target.

6.1 Real-time estimation

All main analyses used a retrospective snapshot in which counts are close to final. In real time, reporting delays leave the most recent weeks incomplete, and right-censoring biases the latest counts downwards in proportion to the delay distribution [59, 60]. In the archived issues of these data two thirds of first-issue counts were later revised upward, and the intercept computed from first-issue counts was about 1.6 times that from settled counts (Section 5.7). The current-value target is affected first and most, because its risk is dominated by the last one or two periods, exactly those that are incomplete, whereas a realised or process mean over many periods dilutes them. The R_A statistic also needs C_{t+1}, so it is available only one week after its reference week, and estimates near the end of a series combine smaller effective counts with nowcasting uncertainty. Real-time renewal methods address this by modelling delays and the infection process jointly [2, 61, 62]. In the present framework, nowcasting adds a correction-error term to the counting term, and its variance grows as the delay distribution places more mass beyond the current week. Error statements derived from settled data will therefore overstate the precision available in real time.

Estimation windows raise a related issue. Widening a window in renewal estimators accumulates incidence while imposing constancy of the reproduction number over a longer interval [2, 3]. This trades temporal resolution for precision and is not equivalent to observing more independent periods; the averages in Table 2 instead give each period its own transmission value. Applying their formulas to overlapping windows requires explicit treatment of the overlap.

6.2 Reporting that depends on transmission

Proposition 2 assumes that the reporting probability is independent of transmission. In practice testing, care seeking and admission thresholds can change with epidemic activity [63, 64]. Write $z_t = p_t/\rho$ for the relative reporting probability.

Given R_t, the estimator then has conditional bias

$$\mathrm{E}(\hat{R}_t \mid R_t) - R_t = [\mathrm{E}(z_t \mid R_t) - 1]\, R_t. \tag{17}$$

By Eq. (17), if reporting rises with transmission, so that $\mathrm{Cov}(z_t, R_t) > 0$, high-transmission periods are over-counted and low-transmission periods under-counted, the estimated swings exaggerate the true ones, and the fitted intercept overstates the relative dispersion of transmission. If reporting falls with transmission, for example when admission capacity saturates or testing is rationed at the peak, the estimates damp the swings and the intercept understates it. In both cases the process-mean estimate is biased by $\mathrm{Cov}(z_t, R_t)$, and the bias does not shrink with more cases. Neither direction can be detected from the counts alone; it requires external information on testing or admission practice. The participation adjustment used here addresses only facility participation, and it changed the intercept by 2.0% (Section 5.7).

6.3 Choice of interval distribution

The transform in Eq. (14) needs an interval distribution. The household serial interval of Cowling et al. [47] was kept as the primary choice because it is directly estimated for influenza and widely used; it rests on 14 transmission pairs. The intrinsic generation-time estimate of Chan et al. [49], the more appropriate quantity for a renewal mapping, is close to the lower mean of the sensitivity analysis. Serial intervals, intrinsic generation intervals and realised household generation intervals are different quantities [48], and the mapping from growth rate to reproduction number depends on which is used [46]. The interval rescales all deviations: a mean of 3.2 days gave $a = 0.0117$ and a mean of 4.0 days gave 0.0186, whereas the crossover count stayed between 37.4 and 37.6 and the slope ratio between 1.23 and 1.28 (Table 5). The level of a therefore depends on this choice, and the conclusion that it exceeds zero and the balance between the two error terms do not.

6.4 Identification and limitations

Separating the mechanisms behind the dispersion curve requires information that a single cross-section lacks. Temporal structure can separate persistent transmission variation from independent counting and reporting noise under explicit state-space or renewal models [34, 35, 65], and external measurements of reporting can constrain ρ and its variation. The shape of the curve narrows the candidates without identifying one. In the known-truth audit (Appendix C), no generator reproduced all features of the data. The generators that came closest combined scale-dependent heterogeneity with reporting shocks, and generators built on the state structure of the panel also failed to reproduce it, which leaves the transmission share of a open. The affine form is an approximation. It missed the observed mean loss in three of ten count deciles, and a free count exponent fitted better and moved the intercept towards zero. The held-out comparison did not favour the affine form over a power law (contrast -0.0020, standard error 0.0068; Table S8). The affine intercept is therefore a summary within the observed counts, and the model-free floor at the largest counts is the more direct measure of the count-independent component. Reporting that varies with jurisdiction size would leave the same trace as scale-dependent transmission, and the population gradient of the intercept cannot be attributed to either.

The growth windows are a fixed rule applied to four seasons of one surveillance system, and inference conditions on that selection. Transfer to other epidemic regimes would need data from those regimes. The analysis plan followed exploratory analysis of the same data (Section 4.4), and the primary test of $a = 0$ was already clear at that stage. The plan therefore does not make that test confirmatory. Its value lies in the robustness rule, the season-test calibration and the complete reporting of every planned analysis. The analyses of the second amendment were planned after the first set of results and are secondary. The main analyses used a retrospective snapshot, so reporting delays and right-censoring of recent weeks were not modelled (Section 6.1). Not all simulation checks passed. With about two generations per week the intercept recovered three quarters of the variance of the weekly reproduction number. With near-complete reporting and strong offspring dispersion it overstated that variance, and in the assumed reporting range the counting slope carried no information on offspring dispersion. The two-way interval for a covered its expectation under every redrawn generator. Under reporting shocks, however, it covered the true transmission-plus-reporting total in only about half of the panels, and the observed standard error exceeded the simulated variability under all laws of the plan and under the state-structured generators. As statements about the total, intervals for a therefore under-cover, although the lower limit lies far above zero. Intervals for iterated fits condition on convergence, and within-state slopes were imprecise. Facility participation and revisions measure only parts of ascertainment, and the reporting error that remains in settled counts is not identified from public data.

6.5 Checklist for surveillance practice

Four recommendations for reporting reproduction numbers follow from these results.

1. State the inferential target. An estimate may target the current value R_t, the realised mean $\overline{R}_T$ over the observed periods or the process mean $\overline{R}$. Precision statements of the form A/m apply to the first two under declared reporting assumptions and omit between-period variation for the third.

2. Inflate the counting error when the denominator is an observed count. The delta factor is $1 + (1 - \rho)(\overline{R}^2 + v_R)/A$, which is 1.48 under the illustrative parameters of Section 2.3. At small counts the factor is larger, 2.42 at $m = 10$ in Table 3. The half-weight count of the shrinkage estimator moves from $m_\times$ to $m_\times + (1 - \rho)(\overline{R}^2 + v_R)/v_R$, from 53.1 to 78.6 under the same parameters.
3. Qualify single-period estimates for reporting volatility. One period's counts cannot separate transmission changes from reporting shocks (Proposition 2), so rapid changes should be checked against testing volumes, reporting participation or temporal models before they are interpreted as transmission.
4. Treat dispersion fits as descriptive, report their intercept and estimate it by jurisdiction. The intercept a, the slope b and the crossover count b/a describe the observed statistic on the selected rows. A positive intercept shows that counting error is not the whole error, but it reflects transmission and reporting variation, not a transmission-variance estimate without an assumed mechanism. It differed about twofold between small and large jurisdictions here, so the count-independent component added to the counting error should be specific to the jurisdiction, and b/a is not a sample-size threshold for surveillance.

7 Conclusion

The inferential target decides what additional cases buy. More cases shrink the counting error of the current value and of the realised mean of the observed periods, but they do not remove the between-period variation that limits estimation of the process mean, and with an observed denominator the counting error is larger than the known-scale formula implies. In US influenza hospital admissions the primary test found a count-independent component of squared relative error that stayed above zero under every sensitivity analysis of the plan. Its size is better read from the data than from the affine extrapolation: among the state-weeks with the largest counts the relative error of an apparent reproduction number was about 10%, and counting noise contributed little of it. The component was about twice as large in small as in large jurisdictions, so the count beyond which more cases help little is a property of the jurisdiction rather than a common constant. It reflects transmission and reporting variation without separating them, can understate their sum, and was larger for first-issue counts than for settled ones. Precision statements for reproduction numbers should therefore name their target and add a jurisdiction-specific count-independent component to the counting error; separating its sources requires temporal models or external data on reporting.

Data and code availability

The analyses used the dataset Weekly Hospital Respiratory Data (HRD) Metrics by Jurisdiction, National Healthcare Safety Network, published by the U.S. Centers for Disease Control and Prevention under the identifier ua7e-t2fy [40], in the snapshot retrieved on 20 October 2026. State population totals are the Census Bureau Vintage 2025 estimates [44], and the issue history of the same admissions series is from the Delphi Epidata API [45], retrieved on 5 November 2026. The data-processing code and analysis-window definitions of the original submission are in the public repository https://github.com/wangchi2068/reproduction-number-targets-surveillance (commit `2f5c4074a11cd5dbc8744ce4bf0b706c19171156`). The analysis plan (`analysis/analysis_plan.md`, with Amendment 1 appended to it), Amendment 2 (`analysis/analysis_plan_amendment2.md`), their checksum files (`analysis/analysis_plan.lock` and `analysis/analysis_plan_amendment2.lock`), the combined result file `analysis/results.json`, code, data extracts and simulation settings for every reported analysis are distributed with this article in its experiment directory. Running `bash code/run_all.sh` from the distributed experiment directory regenerates every reported estimate, table and figure from the stored inputs. No individual-level or restricted data were used.

Declarations

CRediT authorship contribution statement

Contributions are reported under the CRediT taxonomy: conceptualization, methodology, software, formal analysis, investigation, data curation, validation, visualization, writing (original draft) and writing (review and editing).

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Ethics approval and consent to participate

Ethics approval and informed consent were not required as this study analyzed publicly available, de-identified aggregate surveillance data.

A Proofs

A.1 Count law and conditional information

The probability generating function of one negative binomial offspring count is $[1 + (R_t/k)(1 - z)]^{-k}$. Conditional independence raises this function to the power n, and substituting $z \mapsto 1 - \rho + \rho z$ for independent reporting gives $[1 + (\rho R_t/k)(1 - z)]^{-nk}$, which proves Eq. (2); its derivatives give the stated moments.

For fixed, known (n, ρ, k) and $R > 0$, the count likelihood has the score and Fisher information:

$$\dot{\ell}(R) = \frac{c}{R} - \frac{(c + nk)\rho}{k + \rho R} = \mathcal{I}(R)(\hat{R} - R), \qquad \mathcal{I}(R) = \frac{m}{R + \rho R^2/k}. \tag{A.1}$$

By Eq. (A.1), $\mathrm{Var}(\hat{R} \mid R) = \mathcal{I}(R)^{-1}$. The support does not depend on R, so the regular one-parameter Cramér–Rao inequality applies to estimators unbiased for every R and gives the attained bound stated in Section 2.2 [24]. The bound therefore applies to estimators unbiased for every R with ρ and k known; biased estimators and unknown nuisance parameters require a separate analysis.

A.2 Proof of Proposition 1

Let $e_t = \hat{R}_t - R_t$. The error is conditionally centred, $\mathrm{E}(e_t \mid R_t) = 0$, and its second moments are as follows:

$$\mathrm{E}(e_t^2) = \frac{\mathrm{E}\, R_t + \rho\, \mathrm{E}\, R_t^2/k}{m} = \frac{A}{m}, \qquad \mathrm{E}[e_t(R_t - \overline{R})] = 0. \tag{A.2}$$

Squaring Eq. (4) proves Eq. (3). Any estimator based on the same count and unbiased conditional on every R_t has expected conditional variance at least $\mathrm{E}[\mathcal{I}(R_t)^{-1}] = A/m$, and its process-mean risk adds the same v_R. At fixed ρ, k and transmission law, $[V(m) - V(2m)]/V(m) = 1/[2(1 + m/m_\times)]$ for $v_R > 0$. The comparison changes exposure n; it does not hold unchanged when the reporting probability changes. The shrinkage weight in Eq. (5) is $\mathrm{Cov}(R_t, \hat{R}_t)/\mathrm{Var}(\hat{R}_t) = v_R/V(m)$.

A.3 Proof of Corollary 1

Conditional on the path $R_{1:T}$, the e_t are independent and centred, hence pairwise uncorrelated and orthogonal to every square-integrable function of the path. Thus the average of $e_1, \ldots, e_T$ has second moment $A/(Tm)$, and $\mathrm{Var}(\overline{\hat{R}}_T) = \mathrm{Var}(\overline{R}_T) + A/(Tm)$. Under the stated stationary autocovariance, summing all covariance pairs gives $T^2\,\mathrm{Var}(\overline{R}_T) = v_R S(\phi, T)$ with the first expression in Eq. (7), and the finite geometric sum yields the second [66]. Summing spatial covariance pairs in the same way gives Eq. (8), for any admissible correlation matrix.

A.4 Proof of Proposition 2 and Corollary 2

Write $z_t = p_t/\rho$ and $u_t = C_t/m - z_t R_t$. Conditional on (R_t, p_t), u_t has zero mean. Independence of p_t and R_t, with $\mathrm{E}\, z_t = 1$ and $\mathrm{E}\, z_t^2 = 1 + \tau^2$, gives:

$$\mathrm{E}(u_t^2) = \frac{\overline{R} + \rho(1 + \tau^2)(\overline{R}^2 + v_R)/k}{m} = \frac{A_H}{m}. \tag{A.3}$$

The error $\hat{R}_t - R_t = u_t + (z_t - 1)R_t$ has uncorrelated terms, and the second has expected square $\tau^2(\overline{R}^2 + v_R) = v_H$, which proves Eq. (11). Because $\mathrm{E}(\hat{R}_t - R_t \mid R_t) = 0$, process-mean risk adds v_R. Under the additional cross-period assumptions of Corollary 2, the errors are centred and independent given the transmission path after integrating out reporting, which gives Eq. (12).

For the identification statement, set $U = (1 + \tau^2)(\overline{R}^2 + v_R)$. The two moments of the scaled count are:

$$\mathrm{E}\, \hat{R}_t = \overline{R}, \qquad \mathrm{Var}(\hat{R}_t) = U - \overline{R}^2 + \frac{\overline{R} + \rho U/k}{m}. \tag{A.4}$$

Whenever distinct admissible values of τ^2 give $v_R = U/(1 + \tau^2) - \overline{R}^2 \geq 0$, the moments in Eq. (A.4) coincide, and varying an exogenous m does not separate them. This proves nonidentification by these moments only; it does not assert equality of full count distributions.

For $v_R > 0$, the crossover scale with volatile reporting, $m_{\times,H} = A_H/(v_R + v_H)$, satisfies:

$$m_\times - m_{\times,H} = \frac{\tau^2(\overline{R}^2 + v_R)\overline{R}(1 + \rho\overline{R}/k)}{v_R(v_R + v_H)} \geq 0. \tag{A.5}$$

Equality holds exactly when $\tau = 0$.

A.5 Observed denominator

Let $M_t \sim \text{Bin}(n, \rho)$ be independent of the offspring given n, and write $M_t = m(1+u_t)$, so that $\text{E}\, u_t = 0$ and $\text{Var}(u_t) = (1-\rho)/m$. Then $\tilde{R}_t = \hat{R}_t/(1+u_t)$. To first order in u_t and $\hat{R}_t - R_t$, the error is approximately:

$$\tilde{R}_t - R_t \approx (\hat{R}_t - R_t) - R_t u_t. \tag{A.6}$$

Given (n, R_t), u_t is independent of $\hat{R}_t - R_t$ and centred, so the cross term vanishes and $\text{E}[(\tilde{R}_t - R_t)^2] \approx A/m + (1-\rho)\,\text{E}(R_t^2)/m$, which is Eq. (9). The Monte Carlo check in Table 3 draws R_t from a gamma law with the stated moments and would discard draws with $M_t = 0$, none of which occurred at the scales shown. The approximation neglects the convexity of $1/(1+u_t)$, which inflates the error when m is small; at $m = 10$ the Monte Carlo error is 1.64 times the delta value.

A.6 Reporting shocks in the ratio statistic

Let the count of week t in a state be reported with probability ρz_t, where the shocks z_t are independent between weeks with $\log z_t \sim N(-s^2/2, s^2)$. To first order, the statistic in Eq. (14) satisfies $\log R_{A,t} \approx \text{const} + \kappa(g) \log g_t$ with $\kappa(g) = (g_\mu/7)/[1 + \theta \log(g)/7]$, because $\alpha\theta = g_\mu$. The shocks enter the observed growth factor as $\log g_t = \log g_t^{\text{tr}} + \eta_t$, with $\eta_t = \log z_{t+1} - \log z_t$ and g_t^{tr} the growth factor without shocks. Hence $\text{Var}(\eta_t) = 2s^2$, and the relative variance that the shocks add to the ratio statistic is:

$$\tilde{v}_H \approx 2\kappa^2 s^2. \tag{A.7}$$

This component does not scale with count and therefore enters the intercept a. A reporting level that is constant over consecutive weeks of a state ($z_{t+1} = z_t$) gives $\eta_t = 0$ and cancels. Consecutive rows share one shock with opposite signs, so $\text{Cov}(\eta_t, \eta_{t+1}) = -s^2$ and their correlation is $-1/2$. The affine fit uses squared deviations, and for jointly Gaussian terms a correlation of $-1/2$ becomes a correlation of $+1/4$ between their squares. Resampling single weeks does not preserve this serial dependence, so interval coverage was checked directly. Under the audit generators S6 and S8 (Appendix C), which contain such shocks, the pigeonhole bootstrap intervals for a, b and b/a covered the Monte Carlo means of their estimators in at least 99.5% of 400 panels (Table 21). Under the fresh-draw generators with such shocks its standard deviation for a exceeded the Monte Carlo value by a factor of 1.55–1.67 (Table S13). For the realised mean over T consecutive rows of one state the mean of the shock terms telescopes to $(\log z_{T+1} - \log z_1)/T$, so their contribution to the error of the realised mean falls as T^{-2} rather than T^{-1}. With κ near 0.5 for the primary interval and $s = 0.122$ as in generator S6, Eq. (A.7) gives a shock component of about half the observed intercept.

B Extensions

B.1 Forecasting and shrinkage

Under fixed, known reporting and the stationary covariance of Corollary 1, with the counting error at t centred given the transmission path, the best forecast of R_{t+1} of the form $\overline{R} + \lambda(\hat{R}_t - \overline{R})$ has the weight and risk:

$$\lambda^* = \frac{\phi v_R}{v_R + A/m} = \phi\,\gamma(m), \qquad \text{MSE}_{\text{lin}} = v_R - \frac{\phi^2 v_R^2}{v_R + A/m}. \tag{B.1}$$

This follows from $\text{Cov}(\hat{R}_t, R_{t+1}) = \phi v_R$. The forecast weight in Eq. (B.1) is the shrinkage weight discounted by persistence, and its limit risk is $(1-\phi^2)v_R$. For the process mean, shrinking $\hat{R}_t$ towards a fixed benchmark μ_0 with oracle weight $V(m)/[V(m) + (\mu_0 - \overline{R})^2]$ can give risk below v_R; this lies outside the conditionally unbiased class of Proposition 1, and estimating the weight requires its own risk analysis.

B.2 Allocation between periods, cases and coverage

Proposition 3 (Budget allocation between periods and cases)**.** *Under H1–H4, independent, identically distributed transmission periods, conditionally independent counts, fixed ρ and linear costs, let the budget be $\mathcal{B} = T(c_0 + c_1 m)$ with $c_0, c_1 > 0$, and treat $T \geq 1$ and $m > 0$ as continuous. If $v_R > 0$ and $\mathcal{B} > c_0$, process-mean risk $(c_0 + c_1 m)(v_R + A/m)/\mathcal{B}$ is strictly convex in m and is minimised at:*

$$m^* = \min\{(c_0 m_\times / c_1)^{1/2},\ (\mathcal{B} - c_0)/c_1\}. \tag{B.2}$$

The optimum in Eq. (B.2) has the form of cluster-size allocation in survey sampling and multilevel design [12, 67]: cases per period should grow only with the square root of $m_\times$ and of the relative cost of a period. Integer designs compare the risk over feasible integer T with $n_T = \lfloor (\mathcal{B}/T - c_0)/(c_1\rho) \rfloor$. Under temporal dependence the variance term becomes $v_R S(\phi, T)/T^2$ and no closed-form optimum follows.

Corollary 3 (Coverage against completeness)**.** *Under H2–H5, let p_t be the participating fraction of ν independent reporting units, each participating with probability ρ, so that $\tau^2 = (1-\rho)/(\rho\nu)$. A proportional increase in completeness ρ reduces single-period process-mean risk more than the same proportional increase in exposure n exactly when $m > \rho^2(\nu-1)/k$.*

Write the single-period risk as $v_R + v_H + A_H/m$ with $v_H = \tau^2 Q$, $Q = \overline{R}^2 + v_R$ and $m = \rho n$, and differentiate with respect to $\log\rho$ and $\log n$. The difference of the two derivatives is:

$$\frac{\partial\,\text{risk}}{\partial\log\rho} - \frac{\partial\,\text{risk}}{\partial\log n} = -\frac{Q}{kn}\left[\frac{kn+\rho}{\rho\nu} - 1\right]. \tag{B.3}$$

Because $Q/(kn) > 0$, the difference is negative, and a proportional increase in ρ lowers the risk more than the same increase in n, exactly when $kn + \rho > \rho\nu$, that is, when $m > \rho^2(\nu-1)/k$. The mechanism is assumed; observed facility participation alone does not imply it.

B.3 Number of seasons

Suppose $S \geq 2$ season effects are independent Gaussian with unknown mean and variance $v_S > 0$, and the count law given the effects does not depend on these parameters. Any estimator unbiased for v_S then has relative standard error at least $(2/(S-1))^{1/2}$. Observing the effects themselves gives at least as much information as the counts, and their sample variance is the minimum-variance unbiased estimator [11, 24]. With the four seasons analysed here the bound is $(2/3)^{1/2} = 0.82$, and additional admissions within the same seasons leave it unchanged. This bound concerns the precision with which the between-season variance component can be estimated; it is distinct from the power of a test of equal season intercepts, which depends on the size of the season differences relative to their standard errors and is reported in Section 5.9.

B.4 Thinning and the delta approximation

Binomial thinning provides a further check on the counting slope. Each unique state-week count is thinned once to $\text{Bin}(C, q)$ for $q \in \{0.8, 0.6, 0.5, 0.4, 0.3, 0.2, 0.1\}$, and the thinned value is reused in every ratio that contains it. Recomputing the statistic and the fit, with the original C_t as regressor, gives the thinning response:

$$b(q) - b(1) \simeq c\,(1/q - 1). \tag{B.4}$$

Under the delta approximation $c = b_{\text{P}}$ whatever the original counting variation, because thinning adds known binomial variance. The ratio c/b_{P} is therefore a check of that approximation and not of the data-generating process. Because the check of Eq. (B.4) rests on a delta approximation, its exact expectation was computed on the observed counts. For each retention probability q the expected increment of the inverse-count slope was evaluated by summing over the binomial law of every thinned count (Table S2; Fig. S1A). The exact increment rate is 0.467 at $q = 0.8$, rises to 0.505 at $q = 0.3$ and falls to 0.480 at $q = 0.2$ and 0.384 at $q = 0.1$. The least-squares rate over all q is 0.408, or 0.917 b_{P}, against the Monte Carlo estimate 0.405 (0.366–0.441), or 0.909 (0.810–0.999) b_{P} (Table S3).

The pooled rate is dominated by the smallest retention probability. The value $q = 0.1$ carries 76% of the leverage in the least-squares rate (Fig. S1B), and without it the rate is 0.488, or 1.096 b_{P}. At $q = 0.1$ many thinned counts are small, and the delta approximation of the variance of a log thinned count fails there. At $C = 5$ and $q = 0.1$ the exact added variance of $\log(\text{Bin}(C, q) + 0.5)$ is 0.21 of the delta value (Table S4; Fig. S1C). The positive-part truncation of the fitted mean never binds. The shortfall of the pooled rate below b_{P} is therefore a property of the approximation at small thinned counts, not evidence about the data-generating process.

In the audit the ratio c/b_{P} lies in 0.95–1.00 under S0, 0.96–1.04 under S4, 0.87–0.98 under S5, 0.83–0.93 under S3 and 0.81–0.92 under S6. The observed value lies outside the ranges of the generators with Poisson offspring (S0, S1, S2P and S4) and inside those with negative binomial offspring, but its confidence interval overlaps all of them. Because thinning adds known binomial variance whatever the original counting process, the check cannot separate offspring overdispersion from under-reporting.

C Known-truth simulation audit

C.1 Design

The audit is a simulation study with known truth, reported under the aims, data-generating mechanisms, estimands, methods and performance measures structure of Morris et al. [68]. It asks what a, b, b/a and the thinning rate recover on the sampling design of the application. Synthetic counts were generated on the exact R_A design: the 1,748 rows, 1,969 unique state-weeks and 210 state runs of the data. Each run is simulated as a whole with shared counts, so that one week's count

Table 17 Design of the known-truth audit

Element	Specification
Aim	What a, b, b/a, b/b_{P} and the thinning rate c recover on the R_A design of the application
Estimands	Transmission relative variance $\tilde{v}_R$; total $\tilde{v}_R + \tilde{v}_H$ when reporting shocks are present; expected slope relative to b_{P}
Methods	The pipeline of the application: R_A transform, relative squared deviation, gamma FGLS affine fit, plug-in b_{P}, within-state slope, week centring, score contrast between power-law and affine fits (positive values favour the affine form)
Performance measures	Monte Carlo mean with its standard error; 2.5% and 97.5% Monte Carlo limits; ratio of means for recovery; quantile of the observed value among simulated values; interval coverage
Calibration targets	Median retained count (all generators); mean fitted intercept equal to the observed a (generators with heterogeneity)
Tests	b/b_{P}, b/a, thinning rate, ratio of pooled to within-state slope, week-centring reduction of a, score contrast; a test is passed when the observed value lies inside the 2.5–97.5% Monte Carlo range
Replicates	200 panels per generator; five streams of 200 for score-contrast quantiles; thinning 100 panels with 20 draws per level; coverage 400 panels per generator (S1, S6, S8, S9) with 200 bootstrap draws each; held-out reference 200 panels per generator
Data-generating mechanisms	
S0	Constant growth g_0, independent Poisson counts
S1	Independent lognormal growth factors ($\sigma = 0.247$), Poisson counts
S2	Constant growth, negative binomial offspring ($k = 0.3$), binomial reporting at $\rho = 0.25$
S2P	S2 with Poisson offspring (pure branching chain with reporting)
S3	S1 combined with the S2 counting model
S4	Growth heterogeneity split equally between state and national-week components, Poisson counts
S5	S4 combined with the S2 counting model
S6	S3 with a state-week reporting shock $\rho_{st} = 0.25 z_{st}$, $\log z_{st} \sim N(-s^2/2, s^2)$, $s = 0.122$
S7a, S7b	S3 with heterogeneity $\sigma^2(\mu) = \sigma_0^2(\mu/45)^{-\gamma}$ for expected count μ, $\gamma = 0.25$ and 0.5
S8	S3 with scale-dependent heterogeneity ($\gamma = 0.25$) and state-week reporting shocks ($s = 0.133$)
S9	S8 with half of the transmission heterogeneity carried by the national-week component of S4 ($s = 0.132$)

In S6, S8 and S9 the shock scale is chosen so that transmission and the shock each carry about half of the variance target set by the observed a. Tests are statistics that no alignment step targets.

is the next week's denominator. The expected count is carried forward by the true growth factors, and the geometric-mean observed growth $g_0 = 1.407$ is used for transitions that are not scored. The simulated C_t serves as the count scale, the filter $C_t \geq 5$ is re-applied, and the estimand of each row is the transform (14) of its true growth factor.

Two alignment steps fix the level of the synthetic panels, and the quantities they fix are calibration targets rather than tests. Each run is level-anchored to the geometric mean of its observed counts, and one scale factor per generator, between 0.797 and 0.844, sets the Monte Carlo mean of the median retained count to the observed median of 45. In every generator with transmission heterogeneity, the amplitude of that heterogeneity is then set so that the Monte Carlo mean of the fitted intercept equals the observed a. The slope ratio b/b_{P}, the crossover b/a, the thinning rate, the pooled-to-within-state slope ratio, the reduction of a under week centring and the score contrast between affine and power-law fits are not targeted and serve as tests. A generator passes a test when the observed point value lies inside its 2.5–97.5% Monte Carlo range. This decision rule and the week-centring test were fixed after the generators had first been compared with the data, and Table 20 applies them to every generator. The lower tail of the retained counts matches the data (10th percentiles 10–11 against 10), but the upper tail does not, since the generators' 90th percentiles are 263–324 against 389 observed (Table 18). Post-stratifying retained rows to the observed deciles of the count moves intercept recovery by at most 3.2%.

Table 17 sets out the design. The generators span the mechanisms that the diagnostics are meant to separate, from independent Poisson counts (S0) to scale-dependent heterogeneity combined with state-week reporting shocks (S8). Generator S9, which carries half of the S8 transmission heterogeneity in a national calendar-week component, was added after the others had been compared with the data. All generators in this appendix keep one heterogeneity field fixed across replicates; the interval checks of the analysis plan (Section 5.10) redraw it in every replicate.

C.2 Results

Without reporting shocks the intercept tracks the relative variance of transmission. When transmission is constant (S0, S2 and S2P) the fitted intercept is essentially zero. With independent heterogeneity, the ratio of a to the true relative variance on the retained rows is 1.01 (Monte Carlo range 0.93–1.09) under S1 and 1.02 (0.91–1.13) under S3. With structured heterogeneity it is 1.12 (1.04–1.19) under S4 and 1.15 (1.06–1.27) under S5 (Table 18; Fig. 9C). With reporting shocks that are fixed across replicates the intercept approximates the sum of both components: under S6, a is 1.01 (0.87–1.14) times the total $\tilde{v}_R + \tilde{v}_H$ and 2.02 (1.71–2.29) times $\tilde{v}_R$ alone. Scale-dependent heterogeneity biases the intercept in the

other direction, to 0.68 (0.60–0.78) and 0.43 (0.37–0.50) of the true relative variance under S7a and S7b, because variation that shrinks with count is absorbed by the slope. Under S8 the two distortions combine, and a is 1.75 (1.47–2.04) times $\tilde{v}_R$ and 0.85 (0.72–0.99) times the total. When the shocks are redrawn in every replicate the intercept falls 6–22% short of the total (Section 5.6), so the near-equality under S6 does not carry over.

The counting slope does not identify the mechanism. The ratio b/b_{P} is 0.93 (0.85–1.01) under S0, 0.80 (0.73–0.86) under S2P and 1.23 (1.13–1.32) under S2 (Table 19; Fig. 9A). The step from S0 to S2P reflects the branching structure of denominator noise, and the step from S2P to S2 reflects offspring overdispersion. Transmission heterogeneity moves the ratio in both directions, from 0.73 under S4 to 2.25 under S7b. Among generators with heterogeneity the crossover count ranges from 22.2 (17.0–27.7) under S4 to 42.1 (30.8–58.2) under S8, so it does not discriminate between mechanisms either.

The curve shape separates generators more sharply. Generators with heterogeneity that does not depend on count give in-sample score differences of 0.012 (S1) to 0.024 (S5) in favour of the affine form, whereas the observed difference is -0.0014, a value reached by at most 0.3% of their panels (Fig. 9D). Under S8 the observed contrast lies at the 0.069 quantile, and the observed crossover and slope ratio lie inside the S8 ranges, but S8 does not reproduce the pooled-to-within slope ratio or the week-centring reduction. S9 raises the week-centring reduction to 19.1% (12.5–25.9), still below the observed value. No generator passes every test, and S8 and S9 pass four of six (Table 20). The held-out version of the contrast points the same way (Table S9). The audited generators therefore bracket the data on most summaries without reproducing all of them. An earlier exploratory grid over the transmission share and the scale dependence of the heterogeneity, not part of the analysis plan, gave implied transmission relative standard deviations of 5.1–20.5% across its 25 cells with heterogeneity; it is superseded by the grid of Table 9.

C.3 Interval calibration

The state-by-week pigeonhole bootstrap is conservative on this design (Table 21). Across S1, S6, S8 and S9, its intervals for a, b and b/a covered the Monte Carlo mean of the corresponding estimator in at least 99.5% of panels, against a nominal 95%, and the mean bootstrap standard deviation exceeded the Monte Carlo value by a factor of 1.6–2.5. Under the fresh-draw generators, which redraw the heterogeneity field in every panel, the ratio for a was 1.55–1.78 (range over panels 1.19–2.22), and the percentile intervals covered the mean of $\hat{a}$ in 0.99–1.00 of panels (Table S13). Every ratio exceeds 1, so the pigeonhole intervals are conservative under both kinds of generator, in line with the known overestimation of variance by the pigeonhole bootstrap in crossed designs [53, 69]. For the mean of a homoscedastic crossed random-effects design, the standard-deviation ratio is at most the square root of 3, about 1.73 [53]. The largest ratio, 1.78 under FD-S1, lies slightly above this bound. The Monte Carlo standard deviation comes from 200 panels and carries a relative standard error of about 5%, so the excess of 3% is within Monte Carlo error. The bound also applies only approximately to $\hat{a}$, a weighted estimator with heteroscedastic rows. The fixed-field ratios of up to 2.5 further reflect the Monte Carlo variation that fixed fields understate. The two-way sandwich covered the mean of $\hat{a}$ in 0.993, 0.925, 0.925 and 0.983 of panels under S1, S6, S8 and S9. Under these fixed-field generators the ratio of the observed sandwich standard error of a to the Monte Carlo standard deviation was 1.7–2.9, which motivated the redrawn generators of Section 5.10. The Wald statistic for equal season intercepts was liberal at the χ^2_3 critical value under the fixed-field generators, rejecting a true null in 0.195 of S6 panels and 0.217 of S8 panels, which is why asymptotic and generator-based references are not used as primary calibrations for the season test.

Table 18 Known-truth audit on the real R_A design: intercept against transmission and total variance by generator

Generator	D median [P10, P90]	Transmission v_R	Total v_R + v_H	Intercept a (MC SE)	a / v_R	a / (v_R + v_H)	True-error intercept	Week-centred a reduction
Observed (R_A, GI 3.6)	45 [10, 389]	unknown	unknown	0.0150 [0.0101, 0.0206]	–	–	–	0.35
S0 Poisson, no heterogeneity	45 [10, 281]	0	0	0.0000 (0.0000)	v_R = 0	total 0	0.00003 [-0.00010, 0.00019]	not defined
S1 iid heterogeneity, Poisson counts	45 [11, 263]	0.0150	0.0150	0.0151 (0.0000)	1.01 [0.93, 1.09]	1.01 [0.93, 1.09]	0.00003 [-0.00010, 0.00021]	-0.01 [-0.04, 0.01]
S2 NB offspring + binomial reporting, no heterogeneity	45 [10, 324]	0	0	0.0001 (0.0000)	v_R = 0	total 0	0.00002 [-0.00013, 0.00018]	not defined
S3 iid heterogeneity + NB/reporting	45 [10, 298]	0.0147	0.0147	0.0150 (0.0001)	1.02 [0.91, 1.13]	1.02 [0.91, 1.13]	0.00003 [-0.00011, 0.00020]	-0.02 [-0.05, 0.01]
S4 state + week heterogeneity, Poisson counts	45 [11, 315]	0.0135	0.0135	0.0150 (0.0000)	1.12 [1.04, 1.19]	1.12 [1.04, 1.19]	0.00002 [-0.00006, 0.00015]	0.62 [0.58, 0.66]
S5 state + week heterogeneity + NB/reporting	45 [10, 324]	0.0130	0.0130	0.0149 (0.0001)	1.15 [1.06, 1.27]	1.15 [1.06, 1.27]	0.00003 [-0.00007, 0.00017]	0.59 [0.53, 0.64]

(continued on next page)

(continued)

Generator	D median [P10, P90]	Transmission v_R	Total v_R + v_H	Intercept a (MC SE)	a / v_R	a / (v_R + v_H)	True-error intercept	Week-centred a reduction
S2P Poisson offspring + binomial reporting, no heterogeneity	45 [10, 293]	0	0	0.0000 (0.0000)	v_R = 0	total 0	0.00002 [-0.00008, 0.00015]	not defined
S6 S3 + state-week reporting shock (half of a)	45 [10, 307]	0.0074	0.0147	0.0148 (0.0001)	2.02 [1.71, 2.29]	1.01 [0.87, 1.14]	0.00746 [0.00590, 0.00877]	-0.01 [-0.05, 0.02]
S7a S3 with scale-dependent heterogeneity, gamma 0.25	45 [10, 296]	0.0217	0.0217	0.0148 (0.0001)	0.68 [0.60, 0.78]	0.68 [0.60, 0.78]	0.00003 [-0.00011, 0.00020]	-0.03 [-0.08, 0.02]
S7b S3 with scale-dependent heterogeneity, gamma 0.5	44 [11, 286]	0.0346	0.0346	0.0150 (0.0001)	0.43 [0.37, 0.50]	0.43 [0.37, 0.50]	0.00002 [-0.00013, 0.00021]	0.00 [-0.08, 0.06]
S8 Combo S6 shock + S7a scale-dep (gamma 0.25)	45 [10, 313]	0.0084	0.0173	0.0148 (0.0001)	1.75 [1.47, 2.04]	0.85 [0.72, 0.99]	0.00896 [0.00729, 0.01093]	-0.02 [-0.06, 0.02]

200 synthetic panels per generator on the real 1748-row R_A design (growth factor g0 = 1.407); analyst scale D = simulated C_t and the filter C_t >= 5 applied to each panel; D quantiles are Monte Carlo means over panels. Calibration targets (not tests): the median retained D (count scale 0.797-0.844, all generators) and, for the generators with heterogeneity (S1, S3, S4, S5, S6, S7a, S7b, S8), the Monte Carlo mean of a, which equals the observed a by construction. All other columns are tests. The upper tail of D is not matched (generator P90 263-324, observed 389); post-stratifying retained rows to the observed deciles of D changes a / (v_R + v_H) as follows: S1 1.007 to 1.008; S4 1.115 to 1.079; S6 1.009 to 1.004; S7a 0.684 to 0.665. v_R: relative variance of the target R_i on retained rows; v_H: current-value error caused by reporting shocks (S6, S8 only). Ratios are ratios of panel means with the 2.5-97.5% range of panel ratios; MC SE = SD/sqrt(panels). True-error intercept: affine fit of ((x - R_i)/xbar)^2. Week-centred reduction = 1 - a_week/a, not defined without transmission heterogeneity. Observed a: primary bootstrap (2,000 draws). Shock log-SD s = 0.122 (S6) and 0.133 (S8). S6: true-error intercept / v_H = 1.02 [0.83, 1.19]; lag-1 correlation of the shock term -0.50 [-0.53, -0.46]. The transmission relative SD implied by the observed a under each generator's a / v_R ranges from 9% to 19%.

Table 19 Known-truth audit: b/a, slope references, thinning and the in-sample contrast by generator

Generator	b/a	b/b_P	b_w/b_P (state FE)	Pooled / within slope	Thinning c/b_P	In-sample contrast	Quantile of observed contrast (MC SE)
Observed	37.4 [18.4, 72.4]	1.257 [0.812, 1.734]	0.852 [0.237, 1.475]	1.48 [0.99, 3.66]	0.909 [0.810, 0.999]	-0.0014	–
S0 Poisson, no heterogeneity	—	0.930 [0.853, 1.005]	0.896 [0.794, 1.021]	1.04 [0.95, 1.13]	0.970 [0.946, 1.000]	-0.0004 [-0.0028, 0.0018]	0.195 (0.013)
S1 iid heterogeneity, Poisson counts	25.9 [19.4, 32.9]	0.855 [0.686, 1.020]	0.819 [0.617, 1.026]	1.05 [0.94, 1.21]	0.950 [0.912, 0.986]	0.0119 [0.0035, 0.0213]	0.001 (0.001)
S2 NB offspring + binomial reporting, no heterogeneity	—	1.225 [1.128, 1.321]	1.206 [1.064, 1.342]	1.02 [0.95, 1.10]	0.929 [0.898, 0.970]	-0.0003 [-0.0033, 0.0031]	0.152 (0.011)
S3 iid heterogeneity + NB/reporting	36.3 [27.4, 44.4]	1.143 [0.983, 1.308]	1.123 [0.899, 1.333]	1.02 [0.93, 1.14]	0.879 [0.828, 0.927]	0.0164 [0.0044, 0.0281]	0.001 (0.001)
S4 state + week heterogeneity, Poisson counts	22.2 [17.0, 27.7]	0.734 [0.602, 0.871]	0.649 [0.486, 0.819]	1.14 [0.99, 1.37]	0.997 [0.964, 1.035]	0.0123 [0.0032, 0.0229]	0.000 (0.000)
S5 state + week heterogeneity + NB/reporting	32.6 [24.7, 40.0]	1.025 [0.866, 1.201]	0.953 [0.743, 1.142]	1.08 [0.94, 1.25]	0.925 [0.875, 0.981]	0.0239 [0.0128, 0.0363]	0.000 (0.000)
S2P Poisson offspring + binomial reporting, no heterogeneity	—	0.797 [0.732, 0.863]	0.774 [0.684, 0.865]	1.03 [0.96, 1.13]	0.974 [0.953, 1.004]	-0.0004 [-0.0037, 0.0022]	0.171 (0.012)
S6 S3 + state-week reporting shock (half of a)	36.4 [28.2, 47.6]	1.144 [0.970, 1.342]	1.141 [0.935, 1.354]	1.01 [0.90, 1.12]	0.869 [0.814, 0.919]	0.0176 [0.0048, 0.0337]	0.003 (0.002)
S7a S3 with scale-dependent heterogeneity, gamma 0.25	51.4 [39.5, 65.9]	1.564 [1.337, 1.787]	1.484 [1.249, 1.759]	1.06 [0.95, 1.18]	0.914 [0.861, 0.986]	-0.0003 [-0.0112, 0.0106]	0.495 (0.016)
S7b S3 with scale-dependent heterogeneity, gamma 0.5	77.5 [58.4, 99.7]	2.251 [1.975, 2.582]	2.058 [1.656, 2.493]	1.10 [0.97, 1.21]	0.922 [0.862, 0.980]	-0.0142 [-0.0225, -0.0046]	0.987 (0.004)
S8 Combo S6 shock + S7a scale-dep (gamma 0.25)	42.1 [30.8, 58.2]	1.306 [1.081, 1.520]	1.278 [1.043, 1.512]	1.02 [0.92, 1.15]	0.881 [0.810, 0.947]	0.0108 [-0.0021, 0.0290]	0.069 (0.008)

b_P is the independent-Poisson reference computed on each panel. S2, S3, S5, S6, S7 and S8 use negative-binomial offspring (k = 0.3) with binomial reporting probability 0.25; S2P uses Poisson offspring with the same reporting. Every column is a test (none is a calibration target). b/a undefined when a ≈ 0 (S0, S2, S2P: true relative variance 0), shown as —. Pooled / within slope = (b/b_P)/(b_w/b_P) with state fixed effects. Observed row: primary bootstrap (2,000 draws); its pooled / within slope is the slope ratio of the fixed-effects decomposition. Thinning uses 100 panels x 20 shared-count binomial draws per q and the authors' least-squares rate through the origin. Contrast = in-sample affine minus power mean score (computed as in the authors' replication code). Last column: share of synthetic panels with a contrast at or below the observed value, pooled over 5 independent random streams of 200 panels each, with its binomial MC SE; the SD of the per-stream quantiles is 0.000-0.039.

Table 20 Declared known-truth tests by generator under one decision rule

Generator	b/b_{P}	b/a	Thinning c/b_{P}	Pooled / within slope	Week-centring reduction	Score contrast	Passed
Observed	1.26	37.4	0.91	1.48	34.7	-0.0014	
S0	fail (0.85 to 1.01)	n.d.	fail (0.95 to 1.00)	fail (0.95 to 1.13)	n.d.	pass (0.195)	1/4
S1	fail (0.69 to 1.02)	fail (19.38 to 32.86)	fail (0.91 to 0.99)	fail (0.94 to 1.21)	fail (-4.0 to 0.8)	fail (0.001)	0/6
S2	pass (1.13 to 1.32)	n.d.	pass (0.90 to 0.97)	fail (0.95 to 1.10)	n.d.	pass (0.152)	3/4
S2P	fail (0.73 to 0.86)	n.d.	fail (0.95 to 1.00)	fail (0.96 to 1.13)	n.d.	pass (0.171)	1/4
S3	pass (0.98 to 1.31)	pass (27.41 to 44.44)	pass (0.83 to 0.93)	fail (0.93 to 1.14)	fail (-5.3 to 0.8)	fail (0.001)	3/6
S4	fail (0.60 to 0.87)	fail (17.04 to 27.65)	fail (0.96 to 1.04)	fail (0.99 to 1.37)	fail (58.0 to 65.7)	fail (0.000)	0/6
S5	fail (0.87 to 1.20)	pass (24.66 to 40.02)	pass (0.87 to 0.98)	fail (0.94 to 1.25)	fail (53.5 to 64.2)	fail (0.000)	2/6
S6	pass (0.97 to 1.34)	pass (28.17 to 47.60)	pass (0.81 to 0.92)	fail (0.90 to 1.12)	fail (-5.0 to 1.6)	fail (0.003)	3/6
S7a	fail (1.34 to 1.79)	fail (39.46 to 65.92)	pass (0.86 to 0.99)	fail (0.95 to 1.18)	fail (-7.5 to 1.9)	pass (0.495)	2/6
S7b	fail (1.98 to 2.58)	fail (58.42 to 99.67)	pass (0.86 to 0.98)	fail (0.97 to 1.21)	fail (-7.5 to 5.6)	fail (0.987)	1/6
S8	pass (1.08 to 1.52)	pass (30.81 to 58.24)	pass (0.81 to 0.95)	fail (0.92 to 1.15)	fail (-6.1 to 2.3)	pass (0.069)	4/6
S9	pass (1.04 to 1.48)	pass (29.45 to 51.74)	pass (0.82 to 0.94)	fail (0.92 to 1.17)	fail (12.5 to 25.9)	pass (0.057)	4/6

Pass or fail under the decision rule of Section 4. Brackets give the Monte Carlo 2.5–97.5% range or, for the power-minus-affine score contrast, the quantile of the observed value pooled over five streams. The week-centring reduction is in per cent. n.d., not defined because the intercept is essentially zero. The pooled/within slope ratio is b/b_w, with b_w the state fixed-effects slope. S9 is S8 with half of the transmission heterogeneity carried by a national calendar-week component.

Table 21 Coverage of the state-week bootstrap intervals on known-truth panels

Generator	a covers v_R + v_H	a covers E(a)	b covers E(b)	b/a covers E(b)/E(a)	E(a) / (v_R + v_H)	Bootstrap SD / MC SD (a, b, b/a)
S1 iid heterogeneity, Poisson counts	1.000 (0.000)	1.000 (0.000)	0.995 (0.004)	1.000 (0.000)	1.004	2.52, 1.69, 2.01
S6 S3 + state-week reporting shock (half of a)	0.998 (0.002)	0.998 (0.002)	0.995 (0.004)	0.998 (0.002)	1.014	1.76, 1.61, 1.72
S8 Combo S6 shock + S7a scale-dep (gamma 0.25)	0.892 (0.015)	1.000 (0.000)	0.998 (0.002)	1.000 (0.000)	0.857	1.71, 1.76, 1.82
S9 S8 with half of transmission heterogeneity national-week	0.963 (0.009)	0.998 (0.002)	1.000 (0.000)	0.998 (0.002)	0.889	1.92, 1.78, 1.94

400 known-truth panels per generator, each analysed as the application: gamma FGLS affine fit on retained rows and 200 state x calendar-week pigeonhole bootstrap draws (L = 1), percentile 95% intervals. Nominal coverage 0.95; binomial MC SE in parentheses (0.000 when every panel is covered; the one-sided 95% lower bound is then 1 - 3/400 = 0.993). E(.) is the Monte Carlo mean of the estimator over panels, so the 'covers E' columns isolate the calibration of the interval from the bias of the estimator; 'covers v_R + v_H' also includes that bias. Last column: mean bootstrap SD over the Monte Carlo SD of the point estimates. Smallest share of bootstrap draws with a converged fit in any generator: 100.0%; largest share of draws with a <= 0: 0.0%.




Figure 9 Known-truth audit of the dispersion diagnostics on the real R_A design, with simulated counts aligned to the observed median. Panel A shows the counting slope relative to the independent-Poisson reference, b/b_P, by generator. Panel B shows the thinning rate relative to the same reference, c/b_P. Panel C shows the fitted intercept a against the true transmission relative variance and the true total, which differ only under S6 and S8. Panel D shows the in-sample score contrast between the affine and power-law fits, with the quantile of the observed value among synthetic panels at the right; boxes span the interquartile range and whiskers 1.5 times it. In panels A–C points are means over 200 panels per generator (100 panels for thinning) and bars are 2.5–97.5% Monte Carlo ranges. The vertical line and shaded band mark the observed value and its bootstrap 95% confidence interval; dotted lines mark 1. Generator labels are defined in Table 17; generator S9 is reported in Tables 20 and 21.

D Supplementary tables and figures

Tables S1–S9 give the slope references, the thinning check, the participation adjustment, the within-state slopes, the off-season reporting descriptors and the held-out comparison of curve shapes cited in the main text. Tables S10–S14 give the decile fits, the state-structured generators, the clamp sensitivity of the wild bootstrap, the pigeonhole bootstrap under fresh-draw generators and the type of every interval quoted in the text. Table S5 also reports the growth-factor statistic G of the original analysis, which the revised analysis does not use, and season-specific fits with separate slopes, which differ from the common-slope season intercepts of Table 15.

Table S1 Branching-Poisson and independent-Poisson slope references: analytic, simulated and observed

Reference	Analytic reference / b_P	Simulated b/b_P (no heterogeneity)	MC SE	Simulated / analytic	Observed b / reference
Branching-Poisson, rho = 0.1	0.944	0.877 [0.802, 0.955]	0.0029	0.929	1.33 [0.86, 1.84]
Branching-Poisson, rho = 0.25	0.861	0.802 [0.734, 0.878]	0.0026	0.932	1.46 [0.94, 2.02]
Branching-Poisson, rho = 0.5	0.722	0.669 [0.618, 0.714]	0.0019	0.927	1.74 [1.12, 2.41]
Branching-Poisson, rho = 1	0.443	0.402 [0.373, 0.434]	0.0012	0.907	2.84 [1.83, 3.94]
Independent-Poisson b_P	1	0.930 [0.853, 1.005]	0.0029	0.930	1.26 [0.81, 1.73]

The branching-Poisson reference is b_Pbr(rho) = mean over the observed retained rows of kappa^2 [(1 - rho) + (C_t + 0.5)/(C_t+1 + 0.5)]; independent-Poisson reference b_P = 0.4455. Simulated: pure branching chain with Poisson offspring and binomial reporting rho (independent-Poisson counts, S0, in the last row), no transmission heterogeneity, count-aligned as in the audit; 200 panels each, Monte Carlo 2.5-97.5% in brackets; simulated / analytic is not normalised by the independent-Poisson row. Observed: primary bootstrap (2,000 draws), b_Pbr recomputed with the weights of each draw. The ratio of the observed slope to the reference depends on the reporting probability.

Table S2 Exact expected thinning increment against the delta prediction b_P

q	Exact rate	Exact rate / b_P	Authors' MC rate [95% CI]	Leverage in c	Positive-part mass (max)
0.8	0.4672	1.049	0.4588 [0.3731, 0.5557]	0.001	0.0e+00
0.6	0.4845	1.087	0.4745 [0.4064, 0.5532]	0.004	0.0e+00
0.5	0.4957	1.113	0.4969 [0.4243, 0.5606]	0.009	0.0e+00
0.4	0.5050	1.133	0.4997 [0.4397, 0.5611]	0.021	0.0e+00
0.3	0.5051	1.134	0.5067 [0.4482, 0.5539]	0.051	0.0e+00
0.2	0.4798	1.077	0.4749 [0.4292, 0.5229]	0.151	0.0e+00
0.1	0.3836	0.861	0.3804 [0.3337, 0.4204]	0.763	0.0e+00
c, all q	0.4084	0.917	0.4051 [0.3656, 0.4413]	1.000	–
c, without q = 0.1	0.4882	1.096	–	–	–

Rate = (b_q - b_1)/(1/q - 1) from the affine FGLS fit of the exact expected thinned loss (binomial supports summed, mass below 1e-13 dropped); b_P = 0.4455 is the delta-method prediction. Leverage: share of the least-squares rate through the origin carried by each q. Check: q = 0.5: independent-thinning MC 0.4903 (SE 0.0048) vs expected-loss 0.4957; q = 0.1: independent-thinning MC 0.3832 (SE 0.0024) vs expected-loss 0.3836.

Table S3 Shared-count binomial thinning of the R_A sample (n=1748): summary by subset

Subset (n)	Fit over	b_P [95% CI]	c [95% CI]	c/b_P [95% CI]	Δa rate $\times 10^3$ [95% CI]	c/b_P by q, range
full (1748)	all q	0.446 [0.428, 0.466]	0.405 [0.366, 0.441]	0.909 [0.810, 0.999]	-0.13 [-0.62, 0.47]	0.85–1.14
	$q \geq 0.2$		0.485 [0.439, 0.528]	1.088 [0.994, 1.186]	-0.81 [-1.70, 0.05]	1.03–1.14
highCt50 (828)	all q	0.421 [0.403, 0.439]	0.473 [0.409, 0.550]	1.126 [0.970, 1.316]	-0.19 [-0.69, 0.15]	1.05–1.14
	$q \geq 0.2$		0.459 [0.381, 0.560]	1.092 [0.903, 1.327]	-0.24 [-0.96, 0.21]	1.05–1.10
both50 (821)	all q	0.418 [0.401, 0.436]	0.493 [0.420, 0.574]	1.178 [1.004, 1.379]	-0.31 [-0.90, 0.10]	1.16–1.21
	$q \geq 0.2$		0.490 [0.397, 0.607]	1.171 [0.953, 1.462]	-0.43 [-1.33, 0.12]	1.16–1.21

Thinning probabilities $q \in \{0.8, 0.6, 0.5, 0.4, 0.3, 0.2, 0.1\}$. c is the increase of the slope per unit of $1/q - 1$ and the Δa rate the corresponding change of the intercept; b_P is the independent-Poisson reference (plug-in delta value). highCt50 requires a current count of at least 50 and both50 both adjacent counts of at least 50; eligibility is fixed before thinning. Pairs bootstrap with B=1000 draws, 400 thinning replicates for the point estimate and 50 per bootstrap draw; the Monte Carlo precision criterion (MC SE at most 0.1 times the point estimate on the monitored summaries) is met; point fits 8400/8400, bootstrap fits 1049999/1050000. The last column gives the range of the per-q point estimates of c/b_P over the q values of the fit. Per-q estimates, simultaneous bands over q and leave-one-q-out fits are in the released file analysis/authors/thinning.json.




Figure S1 Exact thinning expectation on the observed counts. Panel A shows the expected increment rate of the inverse-count slope by retention probability q, exact expectation against Monte Carlo estimates with 95% ranges; the dashed line is the delta prediction $b_P = 0.4455$. Panel B shows the share of each q in the leverage of the pooled least-squares rate c. Panel C shows the exact added variance of the log thinned count relative to its delta approximation, by original count C, for three retention probabilities.

Table S4 Small-count variance of the log thinned count relative to the delta approximation

C	q = 0.8	q = 0.6	q = 0.5	q = 0.4	q = 0.3	q = 0.2	q = 0.1
5	1.05	1.11	1.11	1.04	0.86	0.57	0.21
10	1.02	1.05	1.08	1.12	1.13	0.99	0.53
20	1.01	1.02	1.03	1.05	1.09	1.14	0.97
45	1.00	1.01	1.01	1.02	1.03	1.06	1.14
100	1.00	1.00	1.01	1.01	1.01	1.02	1.06
500	1.00	1.00	1.00	1.00	1.00	1.00	1.01

Exact added variance Var log(Bin(C, q) + 0.5) - Var log(C + 0.5) divided by the delta value (1 - q)/(qC). Values below 1 mean that thinning adds less variance than the delta method assumes.

Table S5 Season-specific fits and reporting participation sensitivity

Statistic / period	n	a / change [95% CI]	b / original a	b/a / adjusted a	Finite/B
RA / 2022-23	365	0.01369 [0.0071, 0.02102]	0.6967	50.91	1000/1000
RA / 2023-24	499	0.006174 [0.002667, 0.01542]	0.7018	113.7	1000/1000
RA / 2024-25	356	0.01268 [0.005687, 0.02276]	0.5826	45.93	1000/1000
RA / 2025-26	528	0.02253 [0.008432, 0.03509]	0.3533	15.68	1000/1000
RA / reporting adjustment	1748	0.0002939 [-0.0003183, 0.0008476]	0.01495	0.01466	1000/1000
G / 2022-23	367	0.07179 [0.03786, 0.1162]	2.702	37.64	1000/1000
G / 2023-24	503	0.03046 [0.0118, 0.07462]	2.382	78.2	1000/1000
G / 2024-25	357	0.06469 [0.03191, 0.1278]	2.082	32.19	1000/1000
G / 2025-26	529	0.1383 [0.03715, 0.2252]	0.7411	5.359	1000/1000
G / reporting adjustment	1756	0.000373 [-0.003559, 0.003454]	0.08401	0.08364	1000/1000

Season rows report a with its interval and b, b/a as point estimates only. Reporting-adjustment rows use the a column for original-minus-adjusted a [95% CI], b for original a, and b/a for adjusted a; the original normalizer is fixed. Season b and b/a come from separate per-season fits without a stored bootstrap; season b/a with intervals from a common-slope model are given in the season table. Summer rows are reported in a separate table. RA/2022-23: 1000/1000 finite draws; RA/2023-24: 1000/1000 finite draws; RA/2024-25: 1000/1000 finite draws; RA/2025-26: 1000/1000 finite draws; G/2022-23: 1000/1000 finite draws; G/2023-24: 1000/1000 finite draws; G/2024-25: 1000/1000 finite draws; G/2025-26: 1000/1000 finite draws.

Table S6 Within-state, within-week and two-way slopes against the independent-Poisson reference (bootstrap 95% confidence interval in brackets)

Estimator	Slope [95% CI]	Slope / b_P	Slope / pooled b	Fit status
Pooled affine slope b	0.560 [0.362, 0.769]	1.26 [0.81, 1.73]	–	–
b_P, independent-Poisson reference	0.446 [0.427, 0.465]	–	–	–
State FE, authors' iterated FGLS (b_w)	0.379 [0.107, 0.650]	0.85 [0.24, 1.47]	0.68 [0.26, 1.01]	–
State FE, one-step WLS	0.456 [0.228, 0.682]	1.02 [0.51, 1.53]	0.81 [0.57, 0.96]	single step
Week FE, one-step WLS	0.634 [0.420, 0.846]	1.42 [0.96, 1.89]	1.13 [0.98, 1.34]	single step
State + week FE, one-step WLS	0.480 [0.208, 0.744]	1.08 [0.47, 1.66]	0.86 [0.47, 1.14]	single step
State + week FE, damped iterated FGLS	0.414 [0.040, 0.659]	0.93 [0.09, 1.48]	0.74 [0.09, 1.10]	converged, 124 it.

Primary bootstrap: state-week weights of the primary stream, 2,000 draws, so the pooled row equals Table S1; draws whose iterated fit does not converge are dropped (1962 valid for the damped two-way slope). Undamped two-way iteration: did not converge after 51 iterations (last values 0.4150, 0.4122); damping 0.5 removes the two-cycle. One-step WLS uses the pooled affine working precision; the iterated estimators re-weight with the fixed-effect fitted means.

Table S7 Off-season summer descriptors

Period	n	Median count scale	Median participation (%)	IQR(log G)
Summer 2023	772	15.55	94.24	0.4873
Summer 2024	400	17.12	45.59	0.6683
Summer 2025	839	16.25	92.31	0.4882

Descriptive summaries of May-October state-weeks with D_t >= 5: rows, median count scale D_t, median reporting participation (%) and IQR of log G. No model is fitted, so no interval is given. Reporting was voluntary in the 2024 summer only; that contrast is descriptive, not a causal or independently replicated intervention.

Table S8 Held-out comparison of the affine and power loss models with paired uncertainty

Held-out unit	Folds	Mean power - affine score	Paired fold SE	z	Folds favouring affine	Bootstrap 95% CI	Bootstrap share favouring affine
State	51	-0.0020	0.0068	-0.30	26/51	[-0.0290, 0.0298]	0.43
Season	4	0.0046	0.0087	0.53	3/4	[-0.0295, 0.0655]	0.54

Score: weighted mean of L/mu + log mu on held-out rows; positive differences favour the affine model. Paired fold SE from the between-fold variation of the paired differences; state-week bootstrap with 1000 draws refits both models in every fold.

Table S9 Held-out power-minus-affine score contrast on synthetic panels from each generator

	Leave one state out (51 folds)			Leave one season out (4 folds)		
Generator Observed	Mean contrast ($\times 10^3$) -2.02	Affine better	Observed quantile	Mean contrast ($\times 10^3$) 4.63	Affine better	Observed quantile
S0	-1.01 (0.19)	0.34	0.235 (0.030)	-0.89 (0.23)	0.38	0.970 (0.012)
S1	12.84 (0.37)	0.99	0.000 (0.000)	12.97 (0.37)	0.99	0.060 (0.017)
S2	-0.21 (0.13)	0.51	0.125 (0.023)	-0.09 (0.15)	0.50	0.985 (0.009)
S2P	-1.00 (0.18)	0.35	0.215 (0.029)	-0.81 (0.31)	0.42	0.980 (0.010)
S3	17.38 (0.44)	1.00	0.000 (0.000)	17.49 (0.44)	1.00	0.000 (0.000)
S4	11.21 (0.36)	0.99	0.000 (0.000)	15.78 (0.38)	1.00	0.010 (0.007)
S5	24.03 (0.48)	1.00	0.000 (0.000)	28.32 (0.58)	1.00	0.000 (0.000)
S6	18.74 (0.59)	0.99	0.000 (0.000)	18.60 (0.58)	0.99	0.045 (0.015)
S7a	-1.32 (0.44)	0.41	0.475 (0.035)	-0.85 (0.44)	0.44	0.805 (0.028)
S7b	-15.42 (0.38)	0.00	0.995 (0.005)	-14.31 (0.40)	0.01	1.000 (0.000)
S8	11.07 (0.60)	0.91	0.050 (0.015)	11.15 (0.62)	0.91	0.220 (0.029)
S9	11.83 (0.59)	0.92	0.055 (0.016)	12.60 (0.62)	0.93	0.145 (0.025)

200 synthetic panels per generator, each analysed with the fold design and score of the data: mean of $L/\mu + \log \mu$ on the held-out rows, weighted by rows, power law minus affine (positive values favour the affine form). Mean contrast with its Monte Carlo standard error; affine better, share of panels with a positive contrast; observed quantile, share of panels with a contrast at or below the observed one, with binomial Monte Carlo standard error.

Table S10 Observed and fitted loss by decile of the current-week count

Decile	C_t	n	Mean C_t	Mean L	95% interval	Affine	Power	Three-par.	Affine inside
1	5–10	188	7.8	0.0684	0.0538–0.0854	0.0906	0.0656	0.0748	no
2	11–15	180	13.0	0.0597	0.0425–0.0787	0.0584	0.0533	0.0559	yes
3	16–21	162	18.4	0.0557	0.0431–0.0699	0.0457	0.0467	0.0469	yes
4	22–32	176	26.7	0.0321	0.0249–0.0404	0.0362	0.0405	0.0392	yes
5	33–45	173	38.5	0.0404	0.0310–0.0501	0.0296	0.0352	0.0330	no
6	46–73	174	57.4	0.0228	0.0174–0.0292	0.0249	0.0303	0.0278	yes
7	74–116	171	93.5	0.0223	0.0166–0.0287	0.0210	0.0251	0.0229	yes
8	117–206	174	153.4	0.0191	0.0144–0.0241	0.0187	0.0208	0.0193	yes
9	207–388	175	274.2	0.0190	0.0132–0.0249	0.0171	0.0167	0.0162	yes
10	390–3870	175	881.9	0.0110	0.0079–0.0137	0.0158	0.0115	0.0130	no

Rows of R_A (1,748) split at the deciles of the current-week count C_t. Mean L: observed mean relative squared deviation; 95% interval from a calendar-week block bootstrap (1,999 draws). Model columns give the mean prediction in the decile under the affine fit $a + b/C$ ($a = 0.0150$, $b = 0.560$), the power law $cC^{-\gamma}$ ($\gamma = 0.383$) and the three-parameter fit $a + bC^{-\gamma}$ ($a = 0.0089$, $\gamma = 0.617$). Quasi-deviance: affine -4346.2, power -4348.6, three-parameter -4356.9. Last column: whether the affine prediction lies inside the interval of the observed mean.

Table S11 State-realistic fresh-draw generators against four features of the observed panel

Cell	σ	T1 b/b_P	T2 P90	T3 pooled/within	T4 state CV	$E(\hat{a})/v_R$	SE/SD	Coverage
SR-pop-c0	failed	1.06–1.43	367	1.07–1.47*	0.41–0.72	1.24	2.31	1.000
SR-pop-c25	failed	1.35–1.89*	364	1.02–1.34*	0.47–0.78	1.26	1.79	0.990
SR-pop-c50	failed	1.67–2.21*	356	1.03–1.31*	0.52–0.93	1.25	1.81	0.990
SR-fe-c25	failed	1.45–1.93*	364	1.04–1.41*	0.55–0.90	1.26	1.78	0.990
SR-pop-c25-r003	failed	1.38–1.84*	363	1.02–1.34*	0.46–0.76	1.27	1.80	1.000
SR-pop-c25-t50	0.161	1.29–1.77*	308*	1.04–1.39*	0.47–0.93	1.56	1.63	0.960
Observed	–	1.26	389	1.48	0.59	–	–	–

State-realistic generators (200 panels per cell): transmission field around a smoothed observed trend with state-specific variance (multiplier proportional to the population-dependent intercept, or to the state fixed-effect intercepts for SR-fe), clustered admissions with extra-Poisson parameter θ_c (c0, c25, c50 $= 0, 0.25, 0.5$), reporting probability 0.01 (0.003 for r003), trend exponent 1 (0.5 for t50). σ: transmission SD scale; "failed" when the smoothed trend alone exceeds the transmission target ($\sigma = 0$). T1–T4: Monte Carlo 2.5–97.5% range of b/b_P, mean P90 of C_t (band 350–428), range of pooled/within slope ratio and of the between-state CV of intercepts; * the observed value lies outside. Success rule of Amendment 2: all four pass in one cell (outcome: M4/M5 unresolved). Last columns: $E(\hat{a})/v_R$, observed SE over Monte Carlo SD of $\hat{a}$, coverage of $E(\hat{a})$ by the two-way interval. Data-informed generator design; no inferential result uses these generators.

Table S12 Wild bootstrap of the intercept under alternative loss clamps

Variant	Valid draws	p (two-sided)	Min. p	Clamped rows (%)	t^* quantiles
clamp at 10^{-9}	9,999	0.0001	0.0001	13.5	−0.15 / 1.80 / 3.38
clamp at 10^{-7} (primary)	9,999	0.0001	0.0001	13.5	−0.15 / 1.80 / 3.38
clamp at 10^{-5}	9,999	0.0001	0.0001	13.8	−0.16 / 1.80 / 3.38
clamp at 10^{-3}	9,999	0.0001	0.0001	17.0	0.43 / 2.03 / 3.53
no clamp	3,768	0.0003	0.0003	13.5	0.32 / 1.11 / 2.88
exclude draws with any clamped row	0	not evaluable	–	–	–

Wild cluster bootstrap-t of the intercept (9,999 draws, calendar-week Webb weights, seed 21); $t_{\text{obs}} = 7.36$. Valid: draws with a converged fit. Minimum attainable p: $1/(\text{valid} + 1)$. Clamped rows: share of bootstrap rows whose loss is at or below the clamp (for no clamp: negative losses, kept). t^*: 2.5%, 50%, 97.5% quantiles. Every draw contains at least one clamped row, so the exclusion variant has no draws. Secondary calibration; it does not change the E1 verdict.

Table S13 Pigeonhole bootstrap SD of the intercept against the Monte Carlo SD

Generator	Panels	MC SD ($\times 10^3$)	Boot. SD ($\times 10^3$)	Boot./MC	Sandwich/MC	Coverage
FD-S1	100	0.976	1.742	1.78 (1.47–2.22)	1.02	1.00
FD-S8	100	1.266	2.121	1.67 (1.29–2.07)	0.92	0.99
FD-S9-25†	100	1.300	2.148	1.65 (1.29–2.05)	0.94	1.00
FD-S9-50†	100	1.324	2.196	1.66 (1.28–2.22)	1.03	1.00
FD-S9-75†	100	1.465	2.270	1.55 (1.19–1.92)	1.07	0.99

Pigeonhole (state × calendar-week) bootstrap, 199 draws per panel, on the first 100 stored panels of each fresh-draw generator (panels regenerated exactly from the stored seeds). MC SD: Monte Carlo SD of $\hat{a}$ over 200 panels. Ratio range over panels in brackets. Sandwich: two-way SE over MC SD. Coverage: share of panels whose percentile interval covers $E(\hat{a})$. † post hoc law. Reading rule of Amendment 2: conservative if every ratio exceeds 1 (outcome: conservative).

Table S14 Type and source of every interval quoted in the manuscript

Location	Quantity	Quoted	Type	Source
Abstract; Results: count-independent floor	a, pooled affine intercept (primary test)	0.0109–0.0191	two-way CGM sandwich (state x calendar week), t with G-1 = 41 df	`results.json.revision_2026_11.E1_intercept.primary.inference.two_way.a_ci`
Results: count-independent floor	b, counting slope	0.430–0.690	two-way CGM sandwich (state x calendar week), t with G-1 = 41 df	`results.json.revision_2026_11.E1_intercept.primary.inference.two_way.b_ci`
Results: count-independent floor	a, clustering by state only	0.0125–0.0174	one-way CGM sandwich by state, t with 50 df	`results.json.revision_2026_11.E1_intercept.primary.inference.state.a_ci`
Results: count-independent floor	a, clustering by calendar week only	0.0112–0.0188	one-way CGM sandwich by calendar week, t with 41 df	`results.json.revision_2026_11.E1_intercept.primary.inference.week.a_ci`
Results: count-independent floor	a, block bootstrap	0.0107–0.0192	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11.E1_intercept.block_bootstrap.a.ci`
Results: count-independent floor	a, point estimates over 17 specifications	0.0117–0.0202	range of point values (not an interval)	`results.json.revision_2026_11.E1_intercept.sensitivities.*.a (min and max)`
Results: count-independent floor	a at C_t >= 50	0.0088–0.0172	two-way CGM sandwich (state x calendar week), t with G-1 = 41 df	`results.json.revision_2026_11.E1_intercept.sensitivities.S~a3_floor_50.inference.two_way.a_ci`
Results: count-independent floor	a, fixed calendar windows	0.0092–0.0311	two-way CGM sandwich (state x calendar week), t with G-1 = 41 df	`results.json.revision_2026_11.E1_intercept.sensitivities.S~a4_calendar_windows.inference.two_way.a_ci`
Results: count-independent floor	a, leave-one-week-out point estimates	0.0136–0.0156	range of point values (not an interval)	`results.json.revision_2026_11.E1_intercept.sensitivities.S~a9_leave_one_week_out (min and max)`
Results: fit of the affine form	mean L, lowest count decile	0.0538–0.0854	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.B_floor.B1_deciles.rows[0].mean_L_ci`
Results: fit of the affine form	mean L, highest count decile	0.0079–0.0137	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.B_floor.B1_deciles.rows[9].mean_L_ci`
Results: fit of the affine form	mean L, fifth count decile	0.0310–0.0501	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.B_floor.B1_deciles.rows[4].mean_L_ci`
Results: fit of the affine form	gamma, three-parameter fit, week bootstrap	0.355–0.870	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.B_floor.B3_three_parameter.week_bootstrap.gamma.ci`
Results: fit of the affine form	gamma, three-parameter fit, state bootstrap	0.438–0.862	state bootstrap percentile (states resampled) (999 draws)	`results.json.revision_2026_11_r2.B_floor.B3_three_parameter.state_bootstrap.gamma.ci`
Results: fit of the affine form	a, three-parameter fit, week bootstrap	−0.0014 to 0.0137	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.B_floor.B3_three_parameter.week_bootstrap.a.ci`
Results: fit of the affine form	a, three-parameter fit, state bootstrap	0.0034–0.0144	state bootstrap percentile (states resampled) (999 draws)	`results.json.revision_2026_11_r2.B_floor.B3_three_parameter.state_bootstrap.a.ci`
Results: fit of the affine form	mean L, rows at or above P90	0.0078–0.0139	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.B_floor.B4_model_free_floor.rows[1].mean_L_ci`
Abstract; Results: fit of the affine form	relative RMSE, rows at or above P90 (%)	8.8–11.8	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.B_floor.B4_model_free_floor.rows[1].rel_rmse_ci`
Results: fit of the affine form	floor L - b_hat/C, P90	0.0069–0.0132	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.B_floor.B4_model_free_floor.rows[1].floor_minus_bhat_ci`
Results: fit of the affine form	floor relative SD, P90 (%)	8.3–11.5	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.B_floor.B4_model_free_floor.rows[1].floor_rel_sd_bhat_ci`
Results: fit of the affine form	floor L - b_P/C, P90	0.0071–0.0132	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.B_floor.B4_model_free_floor.rows[1].floor_minus_bP_ci`
Results: fit of the affine form	floor relative SD, P75 (%)	9.8–13.2	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.B_floor.B4_model_free_floor.rows[0].floor_rel_sd_bhat_ci`
Results: fit of the affine form	floor relative SD, P95 (%)	8.0–11.1	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.B_floor.B4_model_free_floor.rows[2].floor_rel_sd_bhat_ci`
Results: what more cases buy	crossover b/a, two-way	24.1–58.3	two-way CGM sandwich (state x calendar week), t with G-1 = 41 df, delta method on the log scale	`results.json.revision_2026_11.E1_intercept.primary.inference.two_way.b_over_a_ci`
Results: what more cases buy	crossover b/a, block bootstrap	23.4–60.5	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11.E1_intercept.block_bootstrap.b_over_a.ci`
Results: what more cases buy	crossover b/a over other sensitivity analyses	25.6–55.5	range of point values (not an interval)	`results.json.revision_2026_11.E1_intercept.sensitivities.*.b_over_a (min and max)`
Results: jurisdiction size	a_1, population coefficient	−0.0085 to −0.0023	two-way CGM sandwich (state x calendar week), t with G-1 = 41 df	`results.json.revision_2026_11_r2.B_floor.B2_population.primary_linear.a1_ci_two_way_t41`
Abstract; Results: jurisdiction size	ratio a(P90)/a(P10), linear model	0.408–0.609	state bootstrap percentile (states resampled) (1,999 draws)	`results.json.revision_2026_11_r2.B_floor.B2_population.primary_linear.ratio_ci_state_bootstrap`
Results: jurisdiction size	ratio a(P90)/a(P10), log link	0.324–0.595	state bootstrap percentile (states resampled) (1,999 draws)	`results.json.revision_2026_11_r2.B_floor.B2_population.secondary_log_link.theta_ci_state_bootstrap, mapped by exp(theta (z_P90 - z_P10))`
Results: jurisdiction size	b_1, slope coefficient on population	−0.124 to 0.095	two-way CGM sandwich (state x calendar week), t with G-1 = 41 df	`results.json.revision_2026_11_r2.B_floor.B2_population.secondary_b_varies.b1_ci_two_way_t41`
Results: jurisdiction size	a, smallest-population tertile	0.0145–0.0233	two-way CGM sandwich (state x calendar week), t with G-1 = 41 df (t with 16 df)	`results.json.revision_2026_11_r2.B_floor.B2_population.tertiles_separate[0].a_ci_two_way`
Results: jurisdiction size	a, middle tertile	0.0125–0.0264	two-way CGM sandwich (state x calendar week), t with G-1 = 41 df (t with 16 df)	`results.json.revision_2026_11_r2.B_floor.B2_population.tertiles_separate[1].a_ci_two_way`
Results: jurisdiction size	a, largest-population tertile	0.0090–0.0180	two-way CGM sandwich (state x calendar week), t with G-1 = 41 df (t with 16 df)	`results.json.revision_2026_11_r2.B_floor.B2_population.tertiles_separate[2].a_ci_two_way`
Results: jurisdiction size	ratio tertile 3 / tertile 1, common slope	0.421–0.818	state bootstrap percentile (states resampled) (999 draws)	`results.json.revision_2026_11_r2.B_floor.B2_population.tertiles_common_slope.ratio_ci_state_bootstrap`
Results: what the floor contains	a after week centring (leave-state-out)	0.0078–0.0126	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11.E3_within_week.week_leave_state_out.a_boot.ci`
Results: what the floor contains	reduction of a, leave-state-out week centre (%)	10.2–47.8	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11.E3_within_week.week_leave_state_out.reduction_vs_pooled.ci`
Results: what the floor contains	reduction of a, week centre incl. own row (%)	13.8–49.8	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11.E3_within_week.week.reduction_vs_pooled.ci`
Results: what the floor contains	reduction of a, week centre and scale (%)	22.0–57.0	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11.E3_within_week.week_review.reduction_vs_pooled.ci`
Results: what the floor contains	reduction of a, season centring (%)	−6.8 to 19.1	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11.E3_within_week.season.reduction_vs_pooled.ci`
Results: what the floor contains	implied transmission SD over 71 mechanisms (%)	8.6–18.6	range of point values (not an interval)	`results.json.revision_2026_11.implied_sd_grid.envelope_all.point_range`
Results: what the floor contains	implied transmission SD, propagated (%)	6.9–22.7	propagated envelope: two-way interval of a combined with MC 2.5-97.5% range of r	`results.json.revision_2026_11.implied_sd_grid.envelope_all.interval_envelope`
Results: what the floor contains	implied SD, 13 slope-reproducing mechanisms (%)	8.6–13.9	range of point values (not an interval)	`results.json.revision_2026_11.implied_sd_grid.envelope_reproducing_counting_slope.point_range`
Results: what the floor contains	intercept recovery, daily renewal (%)	75–78	range of Monte Carlo summaries over generators or cells (not an interval)	`results.json.revision_2026_11.daily_renewal.SC1_one_generation_map_carries_over.range`
Results: intercept and its sources	relative bias of a, FD-S1 (%)	0.5–2.3	panel bootstrap percentile over simulated panels (2,000 draws)	`results.json.revision_2026_11_r2.C_total_bias.generators.FD~S1.relative_bias_ci`
Results: intercept and its sources	relative bias of a, FD-S8 (%)	−15.1 to −13.3	panel bootstrap percentile over simulated panels (2,000 draws)	`results.json.revision_2026_11_r2.C_total_bias.generators.FD~S8.relative_bias_ci`
Results: intercept and its sources	relative bias of a, FD-S9 variants (%)	−14.9 to −13.8	range of point values (not an interval)	`results.json.revision_2026_11_r2.C_total_bias.generators.FD~S9~*.relative_bias (min and max)`
Abstract; Results: intercept and its sources	shortfall against the total with reporting shocks (%)	6.3–22.1	range of point values (not an interval)	`results.json.revision_2026_11_r2.C_total_bias.generators.*.relative_bias (min and max over generators_with_reporting_shock)`
Results: intercept and its sources	w_H over laws	0.10–0.91	range of point values (not an interval)	`results.json.revision_2026_11_r2.C_total_bias.generators.*.w_H (min and max)`
Results: intercept and its sources	mechanism-conditional rescaled a	0.0160–0.0192	range of point values (not an interval)	`results.json.revision_2026_11_r2.C_total_bias.mechanism_conditional_rescaling.a_obs_over_t_g_range`
Results: external reporting data	change in a after participation adjustment	−0.00086 to 0.00027	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.D2_reporting_data.coverage_adjustment.difference_ci_week_bootstrap`
Results: external reporting data	a, first-issue counts	0.0133–0.0298	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.D2_reporting_data.backfill.a_first_ci_week_bootstrap`
Results: external reporting data	a, latest-issue counts	0.0096–0.0183	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.D2_reporting_data.backfill.a_latest_ci_week_bootstrap`
Results: external reporting data	a, first minus latest	0.0016–0.0146	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11_r2.D2_reporting_data.backfill.a_first_minus_latest_ci_week_bootstrap`
Results: counting slope	b/b_P, plug-in reference	0.97–1.59	calendar-week block bootstrap percentile (1,999 draws)	`results.json.revision_2026_11.E1_intercept.block_bootstrap.b_over_bP.ci`
Results: counting slope	b/b_P, normalised plug-in reference	1.02–1.64	two-way CGM sandwich (state x calendar week), t with G-1 = 41 df, reference fixed	`results.json.revision_2026_11_r2.F2_poisson_reference.b_over_bP_ci_two_way_bP_fixed.normalised_plugin`
Results: counting slope	b/b_P, simulated exact reference	1.08–1.73	two-way CGM sandwich (state x calendar week), t with G-1 = 41 df, reference fixed	`results.json.revision_2026_11_r2.F2_poisson_reference.b_over_bP_ci_two_way_bP_fixed.simulated_exact`
Results: counting slope	b/b_P, state fixed effects	0.24–1.47	state x calendar-week pigeonhole bootstrap percentile (2,000 draws)	`ma_extensions.json.within.over_bP.state_authors.ci95`
Results: counting slope	b/b_P, week fixed effects	0.96–1.89	state x calendar-week pigeonhole bootstrap percentile (2,000 draws)	`ma_extensions.json.within.over_bP.week_onestep.ci95`
Results: counting slope	expected b/b_P for rho <= 0.01	0.85–0.91	range of Monte Carlo summaries over generators or cells (not an interval)	`results.json.revision_2026_11.rho_k_grid.SC3_b_over_bP_0.9_1.1_at_rho_le_0.01.range`
Results: season intercepts	2023-24 intercept	0.0050–0.0100	two-way CGM sandwich (state x calendar week), t with G-1 = 41 df, delta method on the log scale	`results.json.revision_2026_11.season_test.effects.seasons.2023~24.ci_sandwich_log_delta`
Results: season intercepts	other-season intercepts	0.0154–0.0180	range of point values (not an interval)	`results.json.revision_2026_11.season_test.effects`

References

[1] Christophe Fraser. Estimating individual and household reproduction numbers in an emerging epidemic. *PLoS ONE*, 2(8):e758, 2007.

[2] Katelyn M. Gostic, Lauren McGough, Edward B. Baskerville, Sam Abbott, Keya Joshi, Christine Tedijanto, Rebecca Kahn, Rene Niehus, James A. Hay, Pablo M. De Salazar, Joel Hellewell, Sophie Meakin, James D. Munday, Nikos I. Bosse, Katharine Sherrat, Robin N. Thompson, Laura F. White, Jana S. Huisman, Jérémie Scire, Sebastian Bonhoeffer, Tanja Stadler, Jacco Wallinga, Sebastian Funk, Marc Lipsitch, and Sarah Cobey. Practical considerations for measuring the effective reproductive number, Rt. *PLOS Computational Biology*, 16(12):e1008409, 2020.

[3] Anne Cori, Neil M. Ferguson, Christophe Fraser, and Simon Cauchemez. A new framework and software to estimate time-varying reproduction numbers during epidemics. *American Journal of Epidemiology*, 178(9):1505–1512, 2013.

[4] R.N. Thompson, J.E. Stockwin, R.D. van Gaalen, J.A. Polonsky, Z.N. Kamvar, P.A. Demarsh, E. Dahlqwist, S. Li, E. Miguel, T. Jombart, J. Lessler, S. Cauchemez, and A. Cori. Improved inference of time-varying reproduction numbers during infectious disease outbreaks. *Epidemics*, 29:100356, 2019.

[5] Kris V. Parag. Improved estimation of time-varying reproduction numbers at low case incidence and between epidemic waves. *PLOS Computational Biology*, 17(9):e1009347, 2021.

[6] Kris V. Parag, Christl A. Donnelly, and Alexander E. Zarebski. Quantifying the information in noisy epidemic curves. *Nature Computational Science*, 2(9):584–594, 2022.

[7] Kris V. Parag, Ben Lambert, Christl A. Donnelly, and Sandor Beregi. Asymmetric limits on timely interventions from noisy epidemic data. *Communications Physics*, 8(1):450, 2025.

[8] Rebecca K. Nash, Samir Bhatt, Anne Cori, and Pierre Nouvellet. Estimating the epidemic reproduction number from temporally aggregated incidence data: A statistical modelling approach and software tool. *PLOS Computational Biology*, 19(8):e1011439, 2023.

[9] I. Ogi-Gittins, W.S. Hart, J. Song, R.K. Nash, J. Polonsky, A. Cori, E.M. Hill, and R.N. Thompson. A simulation-based approach for estimating the time-dependent reproduction number from temporally aggregated disease incidence time series data. *Epidemics*, 47:100773, 2024.

[10] Oliver Eales and Steven Riley. Differences between the true reproduction number and the apparent reproduction number of an epidemic time series. *Epidemics*, 46:100742, 2024.

[11] Shayle R. Searle, George Casella, and Charles E. McCulloch. *Variance Components*. John Wiley & Sons, New York, 1992.

[12] William G. Cochran. *Sampling Techniques*. John Wiley & Sons, New York, 3rd edition, 1977.

[13] Rebecca DerSimonian and Nan Laird. Meta-analysis in clinical trials. *Controlled Clinical Trials*, 7(3):177–188, 1986.

[14] Robert E. Fay and Roger A. Herriot. Estimates of income for small places: An application of James-Stein procedures to census data. *Journal of the American Statistical Association*, 74(366a):269–277, 1979.

[15] David Clayton and John Kaldor. Empirical Bayes estimates of age-standardized relative risks for use in disease mapping. *Biometrics*, 43(3):671, 1987.

[16] Andrew Gelman and Phillip N. Price. All maps of parameter estimates are misleading. *Statistics in Medicine*, 18 (23):3221–3234, 1999.

[17] David J. Spiegelhalter. Funnel plots for comparing institutional performance. *Statistics in Medicine*, 24(8):1185–1202, 2005.

[18] David Spiegelhalter, Christopher Sherlaw-Johnson, Martin Bardsley, Ian Blunt, Christopher Wood, and Olivia Grigg. Statistical methods for healthcare regulation: Rating, screening and surveillance. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, 175(1):1–47, 2012.

[19] Steinar Engen, Oyvind Bakke, and Aminul Islam. Demographic and environmental stochasticity-concepts and definitions. *Biometrics*, 54(3):840, 1998.

[20] Russell Lande, Steinar Engen, and Bernt-Erik Sæther. *Stochastic Population Dynamics in Ecology and Conservation*. Oxford University Press, Oxford, 2003.

[21] Carles Bretó, Daihai He, Edward L. Ionides, and Aaron A. King. Time series analysis via mechanistic models. *The Annals of Applied Statistics*, 3(1):319–348, 2009.

[22] Daihai He, Edward L. Ionides, and Aaron A. King. Plug-and-play inference for disease dynamics: measles in large and small populations as a case study. *Journal of the Royal Society Interface*, 7(43):271–283, 2010.

[23] George Casella and Roger L. Berger. *Statistical Inference*. Duxbury, Pacific Grove, CA, 2 edition, 2002.

[24] Erich L. Lehmann and George Casella. *Theory of Point Estimation*. Springer, New York, 2nd edition, 1998.

[25] J. O. Lloyd-Smith, S. J. Schreiber, P. E. Kopp, and W. M. Getz. Superspreading and the effect of individual variation on disease emergence. *Nature*, 438(7066):355–359, 2005.

[26] James O. Lloyd-Smith. Maximum likelihood estimation of the negative binomial dispersion parameter for highly overdispersed data, with applications to infectious diseases. *PLoS ONE*, 2(2):e180, 2007.

[27] L. R. Taylor. Aggregation, variance and the mean. *Nature*, 189(4766):732–735, 1961.

[28] Barry I. Graubard and Edward L. Korn. Inference for superpopulation parameters using sample surveys. *Statistical Science*, 17(1):73–96, 2002.

[29] Jacco Wallinga and Peter Teunis. Different epidemic curves for Severe Acute Respiratory Syndrome reveal similar impacts of control measures. *American Journal of Epidemiology*, 160(6):509–516, 2004.

[30] Kris V Parag and Christl A Donnelly. Adaptive estimation for epidemic renewal and phylogenetic skyline models. *Systematic Biology*, 69(6):1163–1179, 2020.

[31] Samir Bhatt, Neil Ferguson, Seth Flaxman, Axel Gandy, Swapnil Mishra, and James A Scott. Semi-mechanistic Bayesian modelling of COVID-19 with renewal processes. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, 186(4):601–615, 2023.

[32] Amin Azmon, Christel Faes, and Niel Hens. On the estimation of the reproduction number based on misreported epidemic data. *Statistics in Medicine*, 33(7):1176–1192, 2014.

[33] Oliver Stoner, Theo Economou, and Gabriela Drummond Marques da Silva. A hierarchical framework for correcting under-reporting in count data. *Journal of the American Statistical Association*, 114(528):1481–1492, 2019.

[34] Jonas Knape. Estimability of density dependence in models of time series data. *Ecology*, 89(11):2994–3000, 2008.

[35] Marie Auger-Méthé, Chris Field, Christoffer M. Albertsen, Andrew E. Derocher, Mark A. Lewis, Ian D. Jonsen, and Joanna Mills Flemming. State-space models' dirty little secrets: even simple linear Gaussian models can have estimation problems. *Scientific Reports*, 6(1):26677, 2016.

[36] Keith Burghardt, Siyi Guo, and Kristina Lerman. Unequal impact and spatial aggregation distort COVID-19 growth rates. *Philosophical Transactions of the Royal Society A*, 380(2214):20210122, 2022.

[37] Matthew Biggerstaff, Simon Cauchemez, Carrie Reed, Manoj Gambhir, and Lyn Finelli. Estimates of the reproduction number for seasonal, pandemic, and zoonotic influenza: a systematic review of the literature. *BMC Infectious Diseases*, 14:480, 2014.

[38] Anthony R. Ives and Claudio Bozzuto. Estimating and explaining the spread of COVID-19 at the county level in the USA. *Communications Biology*, 4:60, 2021.

[39] Rajendra Bhatia and Chandler Davis. A better bound on the variance. *The American Mathematical Monthly*, 107 (4):353–357, 2000.

[40] Centers for Disease Control and Prevention. Weekly hospital respiratory data (HRD) metrics by jurisdiction, National Healthcare Safety Network (NHSN), 2026.

[41] Aaron M. Frutos, Shunte Moon, Alison M. Binder, Andrea J. Cool, Kingsley Iyawe, Trevor Thompson, Margaret A. Dudeck, Catherine H. Bozio, Mark W. Tenforde, Matthew Biggerstaff, and Kelsey M. Sumner. Evaluation of a novel data source for national influenza surveillance: Influenza hospitalization data in the National Healthcare Safety Network, United States, September 2021–April 2024. *American Journal of Public Health*, 116(2):251–255, 2026.

[42] Centers for Medicare & Medicaid Services. Updates to the condition of participation (CoP) requirements for hospitals and critical access hospitals (CAHs) to report acute respiratory illnesses. Memorandum, 22 October 2024 QSO-25-05-Hospitals/CAHs, Centers for Medicare & Medicaid Services, Center for Clinical Standards and Quality, Baltimore, MD, 2024.

[43] Centers for Disease Control and Prevention. Respiratory illnesses data channel: Release notes and frequently asked questions, 2026.

[44] U.S. Census Bureau. Annual estimates of the resident population for the United States, regions, states, District of Columbia, and Puerto Rico: April 1, 2020 to July 1, 2025 (NST-EST2025-ALLDATA), vintage 2025, 2026.

[45] Alex Reinhart, Logan Brooks, Maria Jahja, Aaron Rumack, Jingjing Tang, et al. An open repository of real-time COVID-19 indicators. *Proceedings of the National Academy of Sciences*, 118(51):e2111452118, 2021.

[46] J Wallinga and M Lipsitch. How generation intervals shape the relationship between growth rates and reproductive numbers. *Proceedings of the Royal Society B: Biological Sciences*, 274(1609):599–604, 2007.

[47] Benjamin J. Cowling, Vicky J. Fang, Steven Riley, J S. Malik Peiris, and Gabriel M. Leung. Estimation of the serial interval of influenza. *Epidemiology*, 20(3):344–347, 2009.

[48] Margaretha Annelie Vink, Martinus Christoffel Jozef Bootsma, and Jacco Wallinga. Serial intervals of respiratory infectious diseases: A systematic review and analysis. *American Journal of Epidemiology*, 180(9):865–875, 2014.

[49] Louis Yat Hin Chan, Sinead E. Morris, Melissa S. Stockwell, Natalie M. Bowman, Edwin Asturias, Suchitra Rao, Karen Lutrick, Katherine D. Ellingson, Huong Q. Nguyen, Yvonne Maldonado, Son H. McLaren, Ellen Sano, Jessica E. Biddle, Sarah E. Smith-Jeffcoat, Matthew Biggerstaff, Melissa A. Rolfes, H. Keipp Talbot, Carlos G. Grijalva, Rebecca K. Borchering, and Alexandra M. Mellis. Estimating the generation time for influenza transmission using household data in the United States. *Epidemics*, 50:100815, 2025.

[50] A. Colin Cameron, Jonah B. Gelbach, and Douglas L. Miller. Robust inference with multiway clustering. *Journal of Business & Economic Statistics*, 29(2):238–249, 2011.

[51] A. Colin Cameron, Jonah B. Gelbach, and Douglas L. Miller. Bootstrap-based improvements for inference with clustered errors. *Review of Economics and Statistics*, 90(3):414–427, 2008.

[52] Matthew D. Webb. Reworking wild bootstrap-based inference for clustered errors. *Canadian Journal of Economics*, 56(3):839–858, 2023.

[53] Art B. Owen. The pigeonhole bootstrap. *The Annals of Applied Statistics*, 1(2):386–411, 2007.

[54] James G. MacKinnon and Matthew D. Webb. The wild bootstrap for few (treated) clusters. *The Econometrics Journal*, 21(2):114–135, 2018.

[55] James G. MacKinnon, Morten Ørregaard Nielsen, and Matthew D. Webb. Cluster-robust inference: A guide to empirical practice. *Journal of Econometrics*, 232(2):272–299, 2023.

[56] Carrie Reed, Sandra S. Chaves, Pam Daily Kirley, Ruth Emerson, Deborah Aragon, Emily B. Hancock, Lisa Butler, Joan Baumbach, Gary Hollick, Nancy M. Bennett, Matthew R. Laidler, Ann Thomas, Martin I. Meltzer, and Lyn Finelli. Estimating influenza disease burden from population-based surveillance data in the United States. *PLOS ONE*, 10(3):e0118369, 2015.

[57] Andrew C Hayward, Ellen B Fragaszy, Alison Bermingham, Lili Wang, Andrew Copas, W John Edmunds, Neil Ferguson, Nilu Goonetilleke, Gabrielle Harvey, Jana Kovar, Megan S C Lim, Andrew McMichael, Elizabeth R C Millett, Jonathan S Nguyen-Van-Tam, Irwin Nazareth, Richard Pebody, Faiza Tabassum, John M Watson, Fatima B Wurie, Anne M Johnson, and Maria Zambon. Comparative community burden and severity of seasonal and pandemic influenza: results of the Flu Watch cohort study. *The Lancet Respiratory Medicine*, 2(6):445–454, 2014.

[58] Kris V. Parag and Christl A. Donnelly. Fundamental limits on inferring epidemic resurgence in real time using effective reproduction numbers. *PLOS Computational Biology*, 18(4):e1010004, 2022.

[59] Michael Höhle and Matthias an der Heiden. Bayesian nowcasting during the STEC O104:H4 outbreak in Germany, 2011. *Biometrics*, 70(4):993–1002, 2014.

[60] Sarah F. McGough, Michael A. Johansson, Marc Lipsitch, and Nicolas A. Menzies. Nowcasting by Bayesian smoothing: A flexible, generalizable model for real-time epidemic tracking. *PLOS Computational Biology*, 16(4):e1007735, 2020.

[61] Sam Abbott, Joel Hellewell, Robin N. Thompson, Katharine Sherratt, Hamish P. Gibbs, Nikos I. Bosse, James D. Munday, Sophie Meakin, Emma L. Doughty, June Young Chun, Yung-Wai Desmond Chan, Flavio Finger, Paul Campbell, Akira Endo, Carl A. B. Pearson, Amy Gimma, Tim Russell, Stefan Flasche, Adam J. Kucharski, Rosalind M. Eggo, and Sebastian Funk. Estimating the time-varying reproduction number of SARS-CoV-2 using national and subnational case counts. *Wellcome Open Research*, 5:112, 2020.

[62] Adrian Lison, Sam Abbott, Jana Huisman, and Tanja Stadler. Generative Bayesian modeling to nowcast the effective reproduction number from line list data with missing symptom onset dates. *PLOS Computational Biology*, 20(4): e1012021, 2024.

[63] Virginia E Pitzer, Melanie Chitwood, Joshua Havumaki, Nicolas A Menzies, Stephanie Perniciaro, Joshua L Warren, Daniel M Weinberger, and Ted Cohen. The impact of changes in diagnostic testing practices on estimates of COVID-19 transmission in the United States. *American Journal of Epidemiology*, 190(9):1908–1917, 2021.

[64] Katharine Sherratt, Sam Abbott, Sophie R. Meakin, Joel Hellewell, James D. Munday, Nikos Bosse, Mark Jit, and Sebastian Funk. Exploring surveillance data biases when estimating the reproduction number: with insights into subpopulation transmission of COVID-19 in England. *Philosophical Transactions of the Royal Society B*, 376 (1829):20200283, 2021.

[65] Shinsuke Koyama, Taiki Horie, and Shigeru Shinomoto. Estimating the time-varying reproduction number of COVID-19 with a state-space method. *PLOS Computational Biology*, 17(1):e1008679, 2021.

[66] James D. Hamilton. *Time Series Analysis*. Princeton University Press, Princeton, NJ, 1994.

[67] Stephen W. Raudenbush. Statistical analysis and optimal design for cluster randomized trials. *Psychological Methods*, 2(2):173–185, 1997.

[68] Tim P. Morris, Ian R. White, and Michael J. Crowther. Using simulation studies to evaluate statistical methods. *Statistics in Medicine*, 38(11):2074–2102, 2019.

[69] Art B. Owen and Dean Eckles. Bootstrapping data arrays of arbitrary order. *The Annals of Applied Statistics*, 6(3): 895–927, 2012.