
Associations of IBD-family genetic liabilities with FinnGen R12 immune phenotypes



b Effect and multiplicity-adjusted evidence

	OR (95% CI)	BH q
IBD | Ankylosing spondylitis	1.274 (1.179-1.376)	1.1e-08
IBD | Iridocyclitis	1.137 (1.089-1.188)	3.8e-08
IBD | Psoriasis	1.134 (1.071-1.201)	5.3e-05
IBD | Psoriatic arthropathy	1.122 (1.047-1.202)	0.0019
CD | Ankylosing spondylitis	1.164 (1.097-1.236)	2.1e-06
CD | Iridocyclitis	1.078 (1.040-1.118)	0.0001
CD | Psoriasis	1.142 (1.063-1.228)	0.00064
CD | Psoriatic arthropathy	1.122 (1.033-1.217)	0.009
UC | Ankylosing spondylitis	1.111 (0.925-1.333)	0.32
UC | Iridocyclitis	1.052 (0.962-1.150)	0.32
UC | Psoriasis	0.995 (0.914-1.084)	0.98
UC | Psoriatic arthropathy	0.999 (0.905-1.102)	0.98

Points and horizontal bars are odds ratios and 95% confidence intervals. BH correction covers the locked 12-test primary family.
All comparisons showed Cochran-Q heterogeneity and significant MR-PRESSO global tests; statistical association is not causal proof.