



Comment on: “The Effectiveness of Resisted Sled Training (RST) for Sprint Performance: A Systematic Review and Meta-Analysis”

Kai Xu¹ · Yimeng Xu² · Jing Zhou² · Ran Wang¹

Accepted: 29 January 2024

© The Author(s), under exclusive licence to Springer Nature Switzerland AG 2024

Dear Editor,

We read with great interest the article “The Effectiveness of Resisted Sled Training (RST) for Sprint Performance: A Systematic Review and Meta-Analysis” by Alcaraz et al. [1]. The authors made an outstanding contribution to resisted sled training (RST) by systematically reviewing the impact of RST on sprint performance and providing comprehensive answers to practical questions. However, we noticed potential errors that may affect the conclusion of this article.

In a subgroup analysis [1], the authors divided the load of RST into either <20% body mass (BM) or ≥20% BM and reported that these two groups had a standardized mean difference of −0.61 (95% confidence interval [CI] −0.97 to −0.25) and −0.45 (95% CI −0.99 to 0.10), respectively. Therefore, the authors concluded that the effect on performance improvement was moderate when comparing loads <20% BM and loads ≥20% BM, with no significant improvements ($p=0.11$) achieved with loads ≥20% BM.

However, we noticed that Alcaraz et al. [1] appeared to make a mistake when categorizing the data reported by

Kawamori et al. [2]. More specifically, Kawamori et al. [2] found a significant effect for both light and heavy sleds training for a 10-m sprint, with the light sleds group improved from 2.06 ± 0.1 s to 2.00 ± 0.10 s, and the heavy sleds group improved from 2.05 ± 0.07 s to 1.94 ± 0.08 s. Alcaraz et al. [1] categorized the light sleds group data into the ≥20% BM subgroup and the heavy sleds group data into the <20% BM subgroup, which may have affected the subgroup analysis results. In fact, after entering the correct data into Revman 5.4 (see Fig. 1), we found that the <20% BM group and the ≥20% BM group have a standardized mean difference of −0.54 (95% CI −0.87 to −0.21) and −0.71 (95% CI −1.39 to −0.02), respectively. In addition, we found significant improvements for both loads <20% BM ($p=0.001$) and ≥20% BM ($p=0.04$). Although only three studies used loads ≥20% BM and the sample size was only 26, the results mentioned above indicated that RST with loads ≥20% BM might be more effective than RST with loads <20% BM, contradicting Alcaraz et al.’s conclusions.

This letter comments on the article available online at <https://doi.org/10.1007/s40279-018-0947-8>.

A response to this letter is available online at <https://doi.org/10.1007/s40279-024-02000-9>.

✉ Ran Wang
wangran@sus.edu.cn

¹ School of Athletic Performance, Shanghai University of Sport, 399 Changhai Road, Shanghai 200438, China

² School of Physical Education, Shanghai University of Sport, 399 Changhai Road, Shanghai 200438, China




Fig. 1 Standardized mean difference between post-intervention and pre-intervention for sprint time in the acceleration phase based on different loads (corrected subgroup analysis results). Squares represent the standardized mean difference for each trial. Diamonds represent

the pooled standardized mean difference across trials. A acceleration phase, *BM* body mass, *CI* confidence interval, *HL* high load, *IV* independent variable, *LL* low load, *ML* moderate load, *SD* standard deviation, *Std.* standardized

In summary, we suggest that researchers and practitioners interpret the RST loads subgroup analysis results in the Alcaraz et al. [1] review cautiously.

Declarations

Funding No sources of funding were used to assist in the preparation of this letter.

Conflict of interest Kai Xu, Yimeng Xu, Jing Zhou, and Ran Wang have no conflicts of interest that are directly relevant to the content of this letter.

Author contributions KX, YX, JZ, and RW made substantial contributions to the content and writing of the letter. All authors read and approved the final manuscript.

References

1. Alcaraz PE, Carlos-Vivas J, Oponjuru BO, Martínez-Rodríguez A. The effectiveness of resisted sled training (RST) for sprint performance: a systematic review and meta-analysis. *Sports Med.* 2018;48(9):2143–65. <https://doi.org/10.1007/s40279-018-0947-8>.
2. Kawamori N, Newton RU, Hori N, Nosaka K. Effects of weighted sled towing with heavy versus light load on sprint acceleration ability. *J Strength Cond Res.* 2014;28(10):2738–45. <https://doi.org/10.1519/JSC.0b013e3182915ed4>.