

Notes of J.E. Marsden's – *Foundations of Mechanics*

Regoon Wang, ChemE@UNSW
wang.regoon@gmail.com

October 2, 2013

Contents

I	Preliminaries	4
1	Differential Theory	5
1.1	Topology	5
1.2	Finite-Dimensional Banach Sapce	11
1.3	Local Differential Calculus	12
1.4	Manifolds and Mappings	12
1.5	Vector Bundles	12
1.6	The Tangent Bundle	12
1.7	Tensors	12
2	Calculus on Manifolds	13
2.1	Vector Fields as Dynamical Systems	14
2.2	Vector Fields as Differential Operators	14
2.3	Exterior Algebra	14
2.4	Cartan's Calculus of Differential Forms	14
2.5	Orientable Manifolds	14
2.6	Integration on Manifolds	14
2.7	Some Riemannian Geometry	14
II	Analytical Dynamics	15
3	Hamiltonian and Lagrangian Systems	16
3.1	Symplectic Algebra	17
3.2	Symplectic Geometry	17
3.3	Hamiltonian Vector Fields and Poisson Brackets	17
3.4	Integral Invariants, Energy Surfaces, and Stability	17
3.5	Lagrangian Systems	17
3.6	The Legendre Transformation	17
3.7	Mechanics on Riemannian Manifolds	17
3.8	Variational Principles in Mechanics	17
4	Hamiltonian Systems with Symmetry	18
4.1	Lie Groups and Group Actions	18
4.2	The Momentum Mapping	18

4.3	Reduction of Phase Space with Symmetry	18
4.4	Hamiltonian Systems on Lie Groups and the Rigid Body	18
4.5	The Topology of Simple Mechanical Systems	18
4.6	The Topology of the Rigid Body	18
5	Hamilton-Jacobi Theory and Mathematical Physics	19
5.1	Time-Dependent Systems	19
5.2	Canonical Transformations and Hamilton-Jacobi Theory	19
5.3	Lagrangian Submanifolds	19
5.4	Quantization	19
5.5	Introduction to Infinite-Dimensional Hamiltonian Systems	19
5.6	Introduction to Nonlinear Oscillations	19
III	An Outline of Qualitative Dynamics	20
6	Topological Dynamics	21
6.1	Limit and Minimal Sets	21
6.2	Recurrence	21
6.3	Stability	21
7	Differentiable Dynamics	22
7.1	Critical Elements	23
7.2	Stable Manifolds	23
7.3	Generic Properties	23
7.4	Structural Stability	23
7.5	Absolute Stability and Axiom A	23
7.6	Bifurcations of Generic Arcs	23
7.7	A Zoo of Stable Bifurcations	23
7.8	Experimental Dynamics	23
8	Hamiltonian Dynamics	24
8.1	Critical Elements	24
8.2	Orbit Cylinders	24
8.3	Stability of Orbits	24
8.4	Generic Properties	24
8.5	Structural Stability	24
8.6	A Zoo of stable Bifurcations	24
8.7	The General Pathology	24
8.8	Experimental Mechanics	24
IV	Celestial Mechanics	25
9	The Two-Body Problem	26
9.1	Models for Two Bodies	26

9.2	Elliptic Orbits and Kepler Elements	26
9.3	The Delaunay Variables	26
9.4	Lagrange Brackets of Kepler Elements	26
9.5	Whittaker's Method	26
9.6	Poincare Variables	26
9.7	Summary of Models	26
9.8	Topology of the Two-Body Problem	26
10	The Three-Body Problem	27
10.1	Models for Three Bodies	27
10.2	Critical Points in the Restricted Three-Body Problem	27
10.3	Closed Orbits in the Restricted Three-Body Problem	27
10.4	Topology of the Planar n-Body Problem	27

Part I

Preliminaries

Chapter 1

Differential Theory

1.1 Topology

Definition 1.1. A topological space is a set S together with a collect \mathcal{O} of subsets called open sets s.t.

(T1) $\emptyset \in \mathcal{O}$ and $S \in \mathcal{O}$;

(T2) If $U_1, U_2 \in \mathcal{O}$, then $U_1 \cap U_2 \in \mathcal{O}$;

(T3) The union of any collection of open sets is open.

For such a topological space the **closed sets** are the elements of $\gamma = \{A \mid A^c \in \mathcal{O}\}$. An **open neighborhood of a point** u is a topological space S is an open set U s.t. $u \in U$. Similarly, for a subset A of S , U is an **open neighborhood** of A if U is open and $A \subset U$. If A is a subset of a topological space S , the **relative topology** on A is defined by $\mathcal{O}_A = \{U \cap A \mid U \in \mathcal{O}\}$.

Then a **basis** for the topology is a collection \mathcal{B} of opensets s.t. every open set of S is a union of elements of \mathcal{B} . The topology is called **first countable** if for each $u \in S$, there is a countable collection $\{U_n\}$ of neighborhoods of u s.t. for any neighborhood U of u , there is an n so $U_n \subset U$. The topology is called **second countable** if it has a countable basis.

Let $\{u_n\}$ be a sequence of points in S . The sequence is said to **coverge** if there is a point $u \in S$ s.t. for every neighborhood U of u , there is an N s.t. $n \geq N$ implies $u_n \in U$. We say that $\{u_n\}$ converges to u or u is a **limit point** of $\{u_n\}$.

Example 1.1. The standard topology of \mathbb{R} is the unions of open intervals (a, b) . Then \mathbb{R} is second countable (hence first countable) with a basis

$$\left\{ \left(r_n - \frac{1}{m}, r_n + \frac{1}{m} \right) \mid r_n \in \mathbb{Q}, m \in \mathbb{N}^+ \right\}$$

.

Definition 1.2. Let S be a topological space and $A \subset S$. Then the **closure** of A , denoted \overline{A} is the inersection of all colsed sets containing A . The **interior** of A , denoted $\overset{\circ}{A}$ is the union of all open sets contained in A . The **boundary** of A , denoted $\partial A := \overline{A} \cap \overline{A}^c$.

Thus, ∂A is closed, and $\partial A = \partial A^c$. Note that A is open iff $A = \overset{\circ}{A}$ and closed iff $A = \overline{A}$.

Proposition 1.1. *Let S be a topological space and $A \subset S$.*

- (i) $u \in \overline{A}$ iff for every neighborhood U of u , $U \cap A \neq \emptyset$.
- (ii) $u \in \overset{\circ}{A}$ iff there is a neighborhood U of u s.t. $U \subset A$.
- (iii) $u \in \partial A$ iff for every neighborhood U of u , $U \cap A \neq \emptyset$ and $U \cap A^c \neq \emptyset$.

Definition 1.3. A point $u \in S$ is called **isolated** iff $\{u\}$ is open. A subset A of S is called **dense** in S iff $\overline{A} = S$ and is called **nowhere dense** iff $(\overline{A})^c$ is dense in S . Thus, A is nowhere dense iff $\overset{\circ}{\overline{A}} = \emptyset$.

Definition 1.4. A topological space S is called **Hausdorff** iff each two distinct points have disjoint neighborhoods. Similarly, S is called **normal** iff each two disjoint closed sets have disjoint neighborhoods.

Proposition 1.2. (i) *A space S is Hausdorff iff $\Delta_S = \{(u, u) \mid u \in S\}$ is closed in $S \times S$ is the product topology.*

(ii) *A first countable space S is Hausdorff iff all sequences have at most one limit point.*

Proof. If Δ_S is closed and u_1, u_2 are distinct, there is an open rectangle $U \times V$ containing (u_1, u_2) and $U \times V \subset \Delta_S^c$. Then in S , U and V are disjoint because if $\exists p \in U \cap V$, then $(p, p) \in U \times V$ it countered with closed set Δ_S . The converse is similar and we leave it as an exercise. \square

Definition 1.5. Let \mathbb{R}^+ denote the nonnegative real numbers with a point $\{+\infty\}$ adjoined, and topology generated by the open intervals of the form (a, b) . A **metric** on set M is a function $d : M \times M \rightarrow \mathbb{R}^+$ s.t.

- (M1) $d(m_1, m_2) = 0$ iff $m_1 = m_2$;
- (M2) $d(m_1, m_2) = d(m_2, m_1)$;
- (M3) $d(m_1, m_3) \leq d(m_1, m_2) + d(m_2, m_3)$

For $\varepsilon > 0$, the ε **disk** about m is defined by $D_\varepsilon(m) = \{m' \in M \mid d(m', m) < \varepsilon\}$. The collection of subsets of M that are unions of such disks is the metric topology of the metric space (M, d) . Two metrics on a set are called **equivalent** if they induce the same metric topology. $\{u_n\}$ is a **Cauchy sequence** iff for all $\varepsilon > 0$, there is an $N \in \mathbb{N}$ s.t. $n, m \geq N$ implies $d(u_n, u_m) < \varepsilon$. The space M is called **complete** if every Cauchy sequence converges. We define $d(u, A) = \inf\{d(u, v) \mid v \in A\}$ and $d(u, \emptyset) = \infty$.

Proposition 1.3. *Every metric space is normal.*

Proof. Let A and B be closed, disjoint subsets of M , and let

$$U = \{u \in M \mid d(u, A) < d(u, B)\}$$

$$V = \{v \in M \mid d(v, A) > d(v, B)\}$$

It is verified that U and V are open, disjoint and $A \subset U, B \subset V$. \square

Definition 1.6. If $\varphi : S \rightarrow T$ is **continuous** at $u \in S$ if $\forall V \ni \varphi(u), \exists U \ni u \Rightarrow \varphi(U) \subset V$. If $\forall V \subset T, \varphi^{-1}(V) = \{u \in S \mid \varphi(u) \in V\}$ is open in S , φ is **continuous**.

If $\varphi : S \rightarrow T$ is a **bijection**, φ and φ^{-1} are continuous, then φ is a **homeomorphism** and S and T are **homeomorphic**.

Proposition 1.4. φ is continuous iff $\forall A \subset S \Rightarrow \varphi(\overline{A}) \subset \overline{\varphi(A)}$.

Proof. If φ is continuous, then $\varphi^{-1}(\overline{\varphi(A)})$ is closed. But $A \subset \varphi^{-1}(\overline{\varphi(A)})$ and hence $\overline{A} \subset \varphi^{-1}(\overline{\varphi(A)})$ or $\varphi(\overline{A}) \subset \overline{\varphi(A)}$. Conversely, let $B \subset T$ be closed and $A = \varphi^{-1}(B)$. Then $\overline{A} \subset \varphi^{-1}(B) = A$, so A is closed. \square

Proposition 1.5. Let $\varphi : S \rightarrow T$ and S first countable set. Then φ is continuous iff $\forall u_n \rightarrow u \Rightarrow \varphi(u_n) \rightarrow \varphi(u)$.

Proposition 1.6. Let M and N be metric spaces with N complete. Then the collection $C(M, N)$ of all continuous maps $\varphi : M \rightarrow N$ forms a complete metric space with the metric $d^0(\varphi, \phi) = \sup\{d(\varphi(u), \phi(u)) \mid u \in M\}$.

Proof. It is readily verified that d^0 is a metric. Convergence of sequence $f_n \in C(M, N)$ to $f \in C(M, N)$ in the metric d^0 is the same as uniform convergence, that is, for all $\varepsilon > 0$ there is an N s.t. if $n \geq N$, $d(f_n(x), f(x)) \leq \varepsilon$ for all $x \in M$. If f_n is a Cauchy sequence in $C(M, N)$, then since $d(f_n(x), f_m(x)) \leq d^0(f_n, f_m)$, $f_n(x)$ is Cauchy for each point $x \in M$. Thus f_n converges pointwise, define a function $f(x)$. We must show that $f_n \rightarrow f$ uniformly and that f is continuous. First of all, given $\varepsilon > 0$, choose N s.t. $d^0(f_n, f_m) < \varepsilon/2$ if $n, m \geq N$. Then for any $x \in M$, pick $N_x \geq N$ s.t. $d(f_m(x), f(x)) < \varepsilon/2$ if $m \geq N_x$. Thus with $n \geq N$ and $m \geq N_x$, $d(f_n(x), f(x)) \leq d(f_n(x), f_m(x)) + d(f_m(x), f(x)) < \varepsilon/2 + \varepsilon/2 = \varepsilon$. So $f_n \rightarrow f$ uniformly. \square

Definition 1.7. S is called **compact** iff $\forall \cup_\alpha U_\alpha = S$ there is a finite subcovering. A subset $A \subset S$ is called **compact** iff A is compact in the relative topology. A space is called *locally compact* iff each point has a neighborhood whose closure is compact.

Theorem 1.1 (Boizano-Weierstrass). If S is a first countable space and is compact, then every sequence has a convergent subsequence.

Proof. Suppose $\{u_n\}$ contains no convergent subsequences. Then we may assume all points are distinct. Each u_n has a neighborhood \mathcal{O}_n that contains no other u_m . $\{u_n\}$ is closed, so that \mathcal{O}_n together with $\{u_n\}^c$ forms an open covering of S , with no finite subcovering. \square

In a metric space, every compact subset is closed and bounded (Heine Borel theorem).

Proposition 1.7. Let S be a Hausdorff space. Then every compact subset of S is closed. Also, every compact Hausdorff space is normal.

Proof. Let $u \in A^c$ and $v \in A$, where A is compact in S . There are disjoint neighborhoods of u and v and, since A is compact, there are disjoint neighborhood of u and A . Thus A^c is open. The second part is an exercise. \square

Proposition 1.8. *Let S be a Hausdorff space that is locally homeomorphic to a locally compact Hausdorff space. Then S is locally compact.*

Proof. Let $U \subset S$ be homeomorphic to $\varphi(U) \subset T$. There is a neighborhood V of $\varphi(u)$ so $\overline{V} \subset \varphi(U)$ and \overline{V} is compact. Then $\varphi^{-1}(\overline{V})$ is compact, and hence closed in S . $\varphi^{-1}(\overline{V}) \subset \varphi^{-1}(V)$. Thus $\varphi^{-1}(V)$ has compact closure. \square

Definition 1.8. A covering $\{U_\alpha\}$ of S is called a **refinement** of a covering $\{V_i\}$ iff $\forall U_\alpha$ there is a V_i s.t. $U_\alpha \subset V_i$. A covering $\{U_\alpha\}$ of S is called **locally finite** iff each point $u \in S$ has a neighborhood U such that U intersects only a finite number of U_α . A space is called **paracompact** iff every open covering of S has a locally finite refinement of open sets, and S is Hausdorff.

Theorem 1.2. *Second countable, locally compact Hausdorff spaces are paracompact.*

Proof. S is the countable union of open sets U_n s.t. $\overline{U_n}$ is compact and $\overline{U_n} \subset U_{n+1}$. If W_α is a covering of S by open sets, and $K_n = \overline{U_n} - U_{n-1}$ then we can cover K_n by a finite number of open sets each of which is contained in some $W_\alpha \cap U_{n+1}$, and is disjoint from $\overline{U_{n-2}}$. The union of such collections yields the desired refinement of $\{W_\alpha\}$. \square

Theorem 1.3. *Every paracompact space is normal.*

Proof. We first show that if A is closed and $u \in A^c$, there are disjoint neighborhoods of u and A . For each $v \in A$ let U_v, V_v be disjoint neighborhoods of u and v . Let W_α be a locally finite refinement of the covering V_v, A^c and $V = \cup W_\alpha$, the union over those α so $W_\alpha \cap A \neq \emptyset$. A neighborhood U_0 of u meets a finite number of W_α . Let U denote the intersection of U_0 and the corresponding U_v . Then V and U are the required neighborhoods. \square

Theorem 1.4. *If S is a Hausdorff space, the following are equivalent:*

- (i) S is normal;
- (ii) For any two closed nonempty disjoint set A, B there is a continuous function $f : S \rightarrow [0, 1]$ s.t. $f(A) = 0, f(B) = 1$. (Urysohn's Lemma)
- (iii) For any closed set $A \subset S$ and continuous function $f : A \rightarrow [a, b]$, there is a continuous extension $\tilde{f} : S \rightarrow [a, b]$ of f (Tietze extension theorem)

Definition 1.9. The **support** of $f : S \rightarrow \mathbb{R}$ is $\text{supp}(f) = \overline{\{x \in S \mid f(x) \neq 0\}}$. A **partition of unity** on S is a family of continuous mappings $\{\varphi_i : S \rightarrow [0, 1]\}$ s.t.

- (i) $\{\text{supp}(\varphi_i)\}$ is locally finite.
- (ii) $\sum_i \varphi_i(x) = 1$ for all x .

We say that a partition of unity $\{\varphi_i\}$ is **subordinate** to a covering $\{A_\alpha\}$ of S if $\text{supp}(\varphi_i)$ is a refinement of $\{A_\alpha\}$.

Theorem 1.5. *Let S be paracompact and $\{U_i\}$ be any open covering of S . Then there is a partition of unity $\{\varphi_i\}$ subordinate to $\{U_i\}$.*

Definition 1.10. A topological space S is **connected** if \emptyset and S are the only subsets of S that are both open and closed. A subset of S is connected iff it is connected in the relative topology. A **component** A of S is a nonempty connected subset of S s.t. the only connected subset of S containing A is A ; S is called **locally connected** iff each point x has an open neighborhood containing a connected neighborhood of x .

Proposition 1.9. *A space S is not connected iff either of the following holds.*

- (i) *There is a nonempty proper subset of S that is both open and closed.*
- (ii) *S is the disjoint union of two nonempty open sets.*
- (iii) *S is the disjoint union of two nonempty closed sets.*

Proposition 1.10. *Let S be a connected space and $f : S \rightarrow \mathbb{R}$ be continuous. Then f assumes every value between any two values $f(u), f(v)$.*

Proof. Suppose $f(u) < a < f(v)$ and f does not assume the value a . Then $U = \{u_0 \mid f(u_0) < a\}$ is both open and closed. \square

Proposition 1.11. *Let S be a topological space and $B \subset S$ be connected.*

- (i) *if $B \subset A \subset \overline{B}$, then A is connected;*
- (ii) *if B_α are connected and $B_\alpha \cap B \neq \emptyset$, then $B \cup (\bigcup_\alpha B_\alpha)$ is connected.*

Proof. If A is not connected, A is the disjoint union of $U_1 \cap A$ and $U_2 \cap A$ where U_1, U_2 are open in S . Then $U_1 \cap B \neq \emptyset, U_2 \cap B \neq \emptyset$, so B is not connected. \square

Definition 1.11. An **arc** φ in S is a continuous mapping $\varphi : I = [0, 1] \rightarrow S$. If $\varphi(0) = u, \varphi(1) = v$, we say φ joins u and v ; S is called **arcwise connected** iff every two points in S can be joined by an arc in S . A space is called **locally arcwise connected** iff each point has an arcwise connected neighborhood.

Proposition 1.12. *Every arcwise connected space is connected. If a space is connected and locally arcwise connected, it is arcwise connected.*

Proof. If S is arcwise connected and not connected, write $S = U_1 \cup U_2$ where U_1, U_2 are nonempty, disjoint and open. Let $u_1 \in U_1, u_2 \in U_2$ and let φ be an arc joining u_1, u_2 . Now $\varphi(I)$ is connected, and since $\varphi(I) \cap U_i \neq \emptyset$, $\varphi \cap U_1 \cap U_2 \neq \emptyset$. Hence $U_1 \cap U_2 \neq \emptyset$, a contradiction. Let $u \in S$ and U denote all points that can be joined to u by an arc. An easy argument shows U and U^c are open and so $U = S$. \square

Definition 1.12. Let S be a metric space with metric d , and 2^S denote the set of all subsets of S . Define $\tilde{d}(A, B) = \sup\{d(a, B) \mid a \in A\}$. As this is not symmetric, we further define $\bar{d}(A, B) = \sup\{\tilde{d}(A, B), \tilde{d}(B, A)\}$. If $A \neq \emptyset, B = \emptyset, \bar{d}(A, B) = \infty, \bar{d}(\emptyset, \emptyset) = 0$. We call it the **Hausdorff metric**.

Proposition 1.13. *Let S be a metric space and d the Hausdorff metric on 2^S . Then $f : S \rightarrow 2^S$ is continuous at $u_0 \in S$ iff for all $\varepsilon > 0$ there is a $\delta > 0$ s.t. $d(u, u_0) < \delta$ implies:*

(i) for all $a \in f(u)$, there is a $b \in f(u_0)$ s.t. $d(a, b) < \varepsilon$; that is

$$f(u) \subset \cup_{b \in f(u_0)} D_\varepsilon(b)$$

(ii) for all $b \in f(u_0)$, there is an $a \in f(u)$ s.t. $d(b, a) < \varepsilon$.

Definition 1.13. Let S be a set. An **equivalence relation** \sim on S is a binary relation s.t. for all $u, v, w \in S$

(i) $u \sim u$;

(ii) $u \sim v$ iff $v \sim u$;

(iii) $u \sim v, v \sim w \Rightarrow u \sim w$.

The **equivalence class** containing u , denoted $[u]$ is defined by $[u] = \{v \in S \mid u \sim v\}$. The set of equivalence classes is denoted S/\sim , and the mapping $\pi : S \rightarrow S/\sim; u \mapsto [u]$ is called the **canonical projection**.

Definition 1.14. $\{U \subset S/\sim \mid \pi^{-1}(U) \text{ is open in } S\}$ is called the **quotient topology**.

Example 1.2. Consider \mathbb{R}^2 and the relation \sim defined by $(a_1, a_2) \sim (b_1, b_2)$ iff $a_1 - b_1, a_2 - b_2 \in \mathbb{Z}$. Then $T^2 = \mathbb{R}^2/\sim$ is called the **2-torus**. In addition to the quotient topology, it inherits a group structure in the usual way: $[(a_1, a_2)] + [(b_1, b_2)] = [(a_1, a_2) + (b_1, b_2)]$.

Example 1.3. The Klein bottle is obtained by reversing one of the orientations. Notice that K^2 is not "orientable" and does not inherit a group structure from \mathbb{R}^2 .

Definition 1.15. Let Z be a topological space and $c : [0, 1] \rightarrow Z$ a continuous map s.t. $c(0) = c(1) = p \in Z$. We call c a **loop** in Z based at p . The loop c is called **contractible** if there is a continuous map $H : [0, 1] \times [0, 1] \rightarrow Z$ s.t. $H(t, 0) = c(t), H(t, 1) = p, \forall t \in [0, 1]$.

Definition 1.16. A space Z is called **simply connected** iff every loop in Z is contractible.

Definition 1.17. Let X be a topological space and $A \subset X$. Then A is called *residual* iff A is the intersection of a countable family of open dense subsets of X . A space X is called a **Baire space** iff every residual set is dense.

Lemma 1.1. *Let X be a locally Baire space; that is, each point $x \in X$ has a neighborhood U s.t. \overline{U} is a Baire space. Then X is a Baire space.*

Proof. Let $A \subset X$ be residual; $A = \cap_1^\infty O_n$ where $\overline{O_n} = X$. Then $A \cap \overline{U} = \cap_1^\infty (O_n \cap \overline{U})$. Now $O_n \cap \overline{U}$ is dense in \overline{U} for if $u \in \overline{U}, u \in O$ then $O \cap U \neq \emptyset, O \cap U \cap O_n \neq \emptyset$. Hence $\overline{U} \subset \overline{A}, \overline{A} = X$. \square

Theorem 1.6 (Baire category). *Every complete pseudometric space is a Baire space.*

Proof. Let $U \subset X$ be open and $A = \cap_1^\infty O_n$ be residual. We must show $U \cap A \neq \emptyset$. Now as $\overline{O_n} = X, U \cap O_n \neq \emptyset$ and so we can choose a disk of diameter less than one, say V_1 , s.t. $\overline{V_1} \subset U \cap O_1$. Proceed inductively to obtain $\overline{V_n} \subset U \cap O_n \cap V_{n-1}$, where V_n has diameter $< 1/n$. Let $x_n \in \overline{V_n}$. Clearly $\{x_n\}$ is a Cauchy sequence, and by completeness has a convergence subsequence with limit point x . Then $x \in \cap_1^\infty \overline{V_n}, U \cap \cap_1^\infty O_n \neq \emptyset$. \square

Exercises

Exercise 1.1. Let S and T be sets and $f : S \rightarrow T$. Show that f is a bijection iff there is a mapping $g : T \rightarrow S$ s.t. $f \circ g, g \circ f$ are identity mappings.

Exercise 1.2. Let X and Y be topological space with Y Hausdorff. Then show that, for any continuous maps $f, g : X \rightarrow Y$, $\{x \in X \mid f(x) = g(x)\}$ is closed. [Hint: Consider the mapping $x \mapsto (f(x), g(x))$].

Exercise 1.3. Prove that in a Hausdorff space, single points are closed.

Exercise 1.4. Define a topological manifold as a space locally homeomorphic to \mathbb{R}^n . Find a topological manifold that is not Hausdorff and not locally compact. [Hint: Consider $\mathbb{R} \cup \{\pm\infty\}$].

Exercise 1.5. Show that the continuous image of a connected space is connected.

1.2 Finite-Dimensional Banach Space

Definition 1.18. A **norm** on a vector space E is a mapping $\|\bullet\| : E \rightarrow \mathbb{R}$ s.t.

(N1) $\|\bullet\| \geq 0, \forall e \in E$ and $\|e\| = 0$ iff $e = 0$.

(N2) $\|\lambda e\| = |\lambda| \|e\|, \forall e \in E, \lambda \in \mathbb{R}$.

(N3) $\|e_1 + e_2\| \leq \|e_1\| + \|e_2\|, \forall e_1, e_2 \in E$.

A normed space whose induced metric is complete is a **Banach space**.

Definition 1.19. Two norms on a vector space E are **equivalent** iff they induce the same topology on E .

Theorem 1.7. Let E be a finite-dimensional real vector space. Then

(i) there is a norm on E ;

(ii) all norms on E are equivalent;

(iii) all norms on E are complete.

Theorem 1.8. For finite-dimensional real vector spaces, linear and multilinear maps are continuous.

Corollary 1.1. Addition and scalar multiplication in a vector space are continuous maps from $E \times E \rightarrow E, \mathbb{R} \times E \rightarrow E$.

Definition 1.20. Given E, F we let $L(E, F)$ denote the set of all linear maps from E into F together with the natural structure of finite dimensional real vector space. Similarly, $L^k(E, F)$ denote the space of multilinear maps from $E \times \cdots \times E$ into F , $L_s^k(E, F)$, the subspace of symmetric elements of $L^k(E, F)$ [that is, if π is any permutation of $\{1, 2, \dots, k\}$, we have $f(e_1, \dots, e_k) = f(e_{\pi(1)}, \dots, e_{\pi(k)})$] and $L_a^k(E, F)$ the subspace of skew symmetric [That is if π is any permutation of $\{1, 2, \dots, k\}$, we have $f(e_1, \dots, e_k) = (\text{sgn } \pi) f(e_{\pi(1)}, \dots, e_{\pi(k)})$, where $\text{sgn } \pi = \pm 1$ according as π is an even or odd permutation].

Theorem 1.9. There is a natural isomorphism $L(E, L^k(E, F)) \approx L^{k+1}(E, F)$.

Exercises

Exercise 1.6. Let $f \in L(E, F)$ so that f is continuous.

- (a) Show that there is a constant K s.t. $\|f(e)\| \leq K \|e\|$ for all $e \in E$. Define $\|f\|$ as the greatest lower bound of such K .
- (b) Show that this is a norm on $L(E, F)$.
- (c) Prove that $\|f \circ g\| \leq \|f\| \cdot \|g\|$.

Exercise 1.7. Suppose $f \in L(E, F)$ and $\dim(E) = \dim(F)$. Then f is an isomorphism iff it is a monomorphism (one-to-one) and iff it is surjective (onto).

Exercise 1.8. Show that two norms $\|\bullet\|_1, \|\bullet\|_2$ are equivalent iff there is a constant M s.t. $M^{-1} \|e\|_1 \leq \|e\|_2 \leq M \|e\|_1$.

Exercise 1.9. Let E be the set of all C^1 functions $f : [0, 1] \rightarrow \mathbb{R}$ with the norm $\|f\| = \sup_{x \in [0, 1]} |f(x)| + \sup_{x \in [0, 1]} |f'(x)|$. Prove that E is a Banach space.

1.3 Local Differential Calculus

Exercises

1.4 Manifolds and Mappings

Exercises

1.5 Vector Bundles

Exercises

1.6 The Tangent Bundle

Exercises

1.7 Tensors

Exercises

Chapter 2

Calculus on Manifolds

2.1 Vector Fields as Dynamical Systems

Exercises

2.2 Vector Fields as Differential Operators

Exercises

2.3 Exterior Algebra

Exercises

2.4 Cartan's Calculus of Differential Forms

Exercises

2.5 Orientable Manifolds

Exercises

2.6 Integration on Manifolds

Exercises

2.7 Some Riemannian Geometry

Exercises

Part II

Analytical Dynamics

Chapter 3

Hamiltonian and Lagrangian Systems

3.1 Symplectic Algebra

Exercises

3.2 Symplectic Geometry

Exercises

3.3 Hamiltonian Vector Fields and Poisson Brackets

Exercises

3.4 Integral Invariants, Energy Surfaces, and Stability

Exercises

3.5 Lagrangian Systems

Exercises

3.6 The Legendre Transformation

Exercises

3.7 Mechanics on Riemannian Manifolds

Exercises

3.8 Variational Principles₁ in Mechanics

Exercises

Chapter 4

Hamiltonian Systems with Symmetry

4.1 Lie Groups and Group Actions

Exercises

4.2 The Momentum Mapping

Exercises

4.3 Reduction of Phase Space with Symmetry

Exercises

4.4 Hamiltonian Systems on Lie Groups and the Rigid Body

Exercises

4.5 The Topology of Simple Mechanical Systems

Exercises

4.6 The Topology of the Rigid Body

Exercises

Chapter 5

Hamilton-Jacobi Theory and Mathematical Physics

5.1 Time-Dependent Systems

Exercises

5.2 Canonical Transformations and Hamilton-Jacobi Theory

Exercises

5.3 Lagrangian Submanifolds

Exercises

5.4 Quantization

Exercises

5.5 Introduction to Infinite-Dimensional Hamiltonian Systems

Exercises

5.6 Introduction to Nonlinear Oscillations

Part III

An Outline of Qualitative Dynamics

Chapter 6

Topological Dynamics

6.1 Limit and Minimal Sets

Exercises

6.2 Recurrence

Exercises

6.3 Stability

Exercises

Chapter 7

Differentiable Dynamics

7.1 Critical Elements

Exercises

7.2 Stable Manifolds

Exercises

7.3 Generic Properties

Exercises

7.4 Structural Stability

Exercises

7.5 Absolute Stability and Axiom A

Exercises

7.6 Bifurcations of Generic Arcs

Exercises

7.7 A Zoo of Stable Bifurcations

7.8 Experimental Dynamics

Chapter 8

Hamiltonian Dynamics

8.1 Critical Elements

8.2 Orbit Cylinders

Exercises

8.3 Stability of Orbits

Exercises

8.4 Generic Properties

Exercises

8.5 Structural Stability

8.6 A Zoo of stable Bifurcations

8.7 The General Pathology

8.8 Experimental Mechanics

Part IV

Celestial Mechanics

Chapter 9

The Two-Body Problem

9.1 Models for Two Bodies

Exercises

9.2 Elliptic Orbits and Kepler Elements

9.3 The Delaunay Variables

9.4 Lagrange Brackets of Kepler Elements

9.5 Whittaker's Method

9.6 Poincare Variables

Exercises

9.7 Summary of Models

Exercises

9.8 Topology of the Two-Body Problem

Chapter 10

The Three-Body Problem

10.1 Models for Three Bodies

Exercises

10.2 Critical Points in the Restricted Three-Body Problem

Exercises

10.3 Closed Orbits in the Restricted Three-Body Problem

Exercises

10.4 Topology of the Planar n-Body Problem

Appendix

General Theory by Kolmogorov