

Supplementary Material for: “Realization of metallic state in 1T-TaS₂ with persisting long-range coherence of charge density wave”

Xin-Yang Zhu,¹ Shi Wang,¹ Zhen-Yu Jia,¹ Li Zhu,¹ Qi-Yuan Li,¹ Wei-Min Zhao,¹ Cheng-Long Xue,¹
Yong-Jie Xu,¹ Zhen Ma,¹ Jinsheng Wen,^{1,2} Shun-Li Yu,^{1,2} Jian-Xin Li,^{1,2} and Shao-Chun Li^{1,2}

¹*National Laboratory of Solid State Microstructures and Department of Physics, Nanjing University, Nanjing 210093, China*

²*Collaborative Innovation Center of Advanced Microstructures, Nanjing University, Nanjing 210093, China*

(Dated: January 21, 2021)

Here, we show the details of the cluster perturbation theory for the theoretical calculations of the single-particle spectral weight in the main text.




FIG. S1. Tiling of the triangular lattice with 13-site clusters. The nearest-neighbor hoppings within a cluster are represented by solid lines, and intercluster hoppings by dashed lines. Here, each site represents a David star in 1T-TaS₂, and the red sites are the David stars where the K⁺ cations are located in the theoretical calculations.

In the main text, the computational technique for our calculation of the single-particle spectral weight $A_i(\omega)$ on site i is based on the cluster perturbation theory (CPT), which has been successfully applied to many strongly correlated systems [1–7]. As shown in Fig. S1, in the CPT calculations, we divide the original lattice into identical 13-site clusters which constitute a superlattice. Then, the lattice Hamiltonian is written as

$$H = H' + V, \quad (\text{S1})$$

where H' is the cluster Hamiltonian, obtained by severing the hopping terms between different clusters, and V contains the intercluster hoppings. The CPT method is based on a strong-coupling perturbation expansion of the one-body operators V linking the individual clusters[1]. At the lowest order of this expansion, the Green's function \mathbf{G} of the original lattice can be expressed (in matrix

form) as

$$\mathbf{G}(\tilde{\mathbf{k}}, \omega) = \mathbf{G}'(\omega)[1 - \mathbf{V}(\tilde{\mathbf{k}})\mathbf{G}'(\omega)]^{-1}, \quad (\text{S2})$$

where \mathbf{G}' is the cluster Green's function, and $\tilde{\mathbf{k}}$ is the wavevector in the Brillouin zone (BZ) of the superlattice. \mathbf{G}' is independent of $\tilde{\mathbf{k}}$, while \mathbf{V} is frequency independent. The cluster Green's function \mathbf{G}' is calculated by the exact diagonalization method with the Lanczos algorithm, and it has the following expression,

$$G'_{\mu\nu}(\omega) = \langle 0 | c_\mu \frac{1}{\omega - H + E_0 + i\eta} c_\nu^\dagger | 0 \rangle + \langle 0 | c_\nu^\dagger \frac{1}{\omega + H - E_0 + i\eta} c_\mu | 0 \rangle, \quad (\text{S3})$$

where E_0 is the energy of the ground state $|0\rangle$, and η is a small real number introduced in the calculation to shift the poles of the Green's function away from the real axis. Here, μ and ν denote both the site and spin degrees of freedom in a cluster. The spectral function of the local single-particle excitation is given by

$$A_{\mu\mu}(\omega) = -\frac{1}{\pi} \sum_{\tilde{\mathbf{k}}} G_{\mu\mu}(\tilde{\mathbf{k}}, \omega). \quad (\text{S4})$$

-
- [1] D. S  n  chal, D. Perez, and M. Pioro-Ladri  re, *Phys. Rev. Lett.* **84**, 522 (2000).
 - [2] M. G. Zacher, R. Eder, E. Arrigoni, and W. Hanke, *Phys. Rev. Lett.* **85**, 2585 (2000).
 - [3] D. S  n  chal and A.-M. S. Tremblay, *Phys. Rev. Lett.* **92**, 126401 (2004).
 - [4] S.-L. Yu, X. C. Xie, and J.-X. Li, *Phys. Rev. Lett.* **107**, 010401 (2011).
 - [5] J. Kang, S.-L. Yu, T. Xiang, and J.-X. Li, *Phys. Rev. B* **84**, 064520 (2011).
 - [6] W.-H. Leong, S.-L. Yu, T. Xiang, and J.-X. Li, *Phys. Rev. B* **90**, 245102 (2014).
 - [7] S.-L. Yu, W. Wang, Z.-Y. Dong, Z.-J. Yao, and J.-X. Li, *Phys. Rev. B* **98**, 134410 (2018).