

# Iterative Hessian Sketch for Constrained Least-Squares

Lydia Hsu, Shuaiwen Wang

December 9, 2018

- Consider the **constrained least-squares**

$$x_{ls} := \arg \min_{x \in C} \frac{1}{2} \|Ax - y\|_2^2$$

where  $C$ : constraint;  $A \in \mathbb{R}^{n \times p}$ : design matrix;  $y$ : response.

- Consider the **constrained least-squares**

$$x_{ls} := \arg \min_{x \in C} \frac{1}{2} \|Ax - y\|_2^2$$

where  $C$ : constraint;  $A \in \mathbb{R}^{n \times p}$ : design matrix;  $y$ : response.

- Issue: when  $n$  large, computationally **expensive**;

- Consider the **constrained least-squares**

$$x_{ls} := \arg \min_{x \in C} \frac{1}{2} \|Ax - y\|_2^2$$

where  $C$ : constraint;  $A \in \mathbb{R}^{n \times p}$ : design matrix;  $y$ : response.

- Issue: when  $n$  large, computationally **expensive**;
- Idea: “sketch”  $A$  using  $SA$ ;

- Consider the **constrained least-squares**

$$x_{ls} := \arg \min_{x \in C} \frac{1}{2} \|Ax - y\|_2^2$$

where  $C$ : constraint;  $A \in \mathbb{R}^{n \times p}$ : design matrix;  $y$ : response.

- Issue: when  $n$  large, computationally **expensive**;
- Idea: “sketch”  $A$  using  $SA$ ;
- $S \in \mathbb{R}^{m \times n}$ : sketching matrix;  $ES^\top S = I_n$ .

# Common sketching matrices

- sub-Gaussian:  $S_{ij} \sim_{iid} \frac{1}{\sqrt{m}} \mathcal{N}(0, 1);$

# Common sketching matrices

- sub-Gaussian:  $S_{ij} \sim_{iid} \frac{1}{\sqrt{m}} \mathcal{N}(0, 1)$ ;
- random row sampling: randomly select  $m$  rows from  $I_n$  (rescaled by  $\sqrt{\frac{n}{m}}$ );

# Common sketching matrices

- sub-Gaussian:  $S_{ij} \sim_{iid} \frac{1}{\sqrt{m}} \mathcal{N}(0, 1)$ ;
- random row sampling: randomly select  $m$  rows from  $I_n$  (rescaled by  $\sqrt{\frac{n}{m}}$ );
- randomized orthogonal system: randomly select  $m$  rows from an orthogonal matrix  $H$  (rescaled by  $\sqrt{\frac{n}{m}}$ );



- Sketch not only  $A$ , but also  $y$ :

$$x_{cs} := \arg \min_{x \in C} \frac{1}{2} \|SAx - Sy\|_2^2$$

- Sketch not only  $A$ , but also  $y$ :

$$x_{CS} := \arg \min_{x \in C} \frac{1}{2} \|SAx - Sy\|_2^2$$

- Classical sketch is **sub-optimal**: assume a linear ground truth  $y = Ax_* + w$ :

$$\|x_{CS} - x_*\| \sim \|x_{ls} - x_*\| \quad \Rightarrow \quad m = \Theta(n)$$

- Sketch not only  $A$ , but also  $y$ :

$$x_{CS} := \arg \min_{x \in C} \frac{1}{2} \|SAx - Sy\|_2^2$$

- Classical sketch is **sub-optimal**: assume a linear ground truth  $y = Ax_* + w$ :

$$\|x_{CS} - x_*\| \sim \|x_{ls} - x_*\| \Rightarrow m = \Theta(n)$$

- We expect the optimal method satisfies  $m \approx O(p)$  **Think about this part more carefully?**

- Only sketch  $A$ , but not  $y$ :

$$x_{hs} := \arg \min_{x \in C} \frac{1}{2} \|SAx\|_2^2 - \langle A^\top y, x \rangle.$$

- The above serves as the building block for an iterative method

- Only sketch  $A$ , but not  $y$ :

$$x_{hs} := \arg \min_{x \in C} \frac{1}{2} \|SAx\|_2^2 - \langle A^\top y, x \rangle.$$

- Still **sub-optimal** in the same sense;
- The above serves as the building block for an iterative method

# Iterative Hessian Sketch

- Idea: construct a new least-squares problem for which the optimal solution is  $x^{LS} - x^t$ . Then applying HS to the problem will produce a new  $x^{t+1}$  whose distance to  $x^{LS}$  has been reduced by a factor of  $\rho$ .
- $\hat{u} := \operatorname{argmin}_{u \in \{C - x^t\}} \left\{ \frac{1}{2} \|Au\|_2^2 - \langle A^T(y - Ax^t), u \rangle \right\}$ .  
By construction,  $\hat{u} = x^{LS} - x^t$ .
- Advantages: sample size efficiency and computational time saving

# Iterative Hessian Sketch Algorithm

## Algorithm

- 1 Initialize  $x^0 = 0$ .
- 2 For iterations  $t = 0, 1, 2, \dots, N - 1$ , generate independent sketches  $S^{t+1} \in \mathbb{R}^{m \times n}$ , and perform the updates
$$x^{t+1} = \operatorname{argmin}_{x \in C} \left\{ \frac{1}{2m} \|S^{t+1} A(x - x^t)\|_2^2 - \langle A^T(y - Ax^t), x \rangle \right\}.$$
- 3 Return  $\hat{x} = x^N$ .







Sarlos (2006)

Title of the publication

*Journal Name* 12(3), 45 – 678.

# Evaluation for Sketch Performance

## Cost Approximation

In terms of  $f$ -cost, the approximated  $\tilde{x}$  is said to be  $\epsilon$ -optimal if

$$f(x^{LS}) \leq f(\tilde{x}) \leq (1 + \epsilon)^2 f(x^{LS}).$$

## Solution Approximation

In terms of  $f$ -cost, the prediction norm defined as

$$\|\tilde{x} - x^{LS}\|_A := \frac{1}{\sqrt{n}} \|A(\tilde{x} - x^{LS})\|_2.$$

Thank you!