

In [1]:

```
import numpy as np
import pandas as pd
import re
import scipy as sp
import xarray as xr
from scipy import integrate
from matplotlib import pyplot as plt
%matplotlib inline
```

In [2]:

```
#读取文件
df1=pd.read_csv('global.csv')
```

In [3]:

```
df2=pd.read_csv('co2_annmean_mlo.csv')
df2=df2.loc[(df2['year'] >1985)&(df2['year'] <2005)][['year','mean']]
```


1.1 [15 points] Following equation 1-2 (without the buffer effect), build a two-box model to compute the atmospheric CO2 level in ppm (parts per million) from 1987 to 2004

In [4]:

```
def gama(t):
    gama=float(df1.loc[df1['Year']==int(t)]['Total carbon emissions from fossil-fuels (million metric tons of C)']/1000/2.13)
    return gama
def NBEM(f, t, k12, k21):
    N1, N2=f
    dfdt=[-k12*N1+k21*N2+gama(t), k12*N1-k21*N2]
    return dfdt
k12=105/740
k21=102/900
#Initial condition
f0=[740/2.13, 900/2.13]
t=np.linspace(1986, 2004, 19)
f=integrate.odeint(NBEM, f0, t, args=(k12, k21))
N1_NBmodel=f[:, 0]
```

In [5]:

```
plt.plot(df2['year'], df2['mean'], 'ok', label='observations')
plt.plot(t[1:19], N1_NBmodel[1:19], 'k', label='calculation without buffer effect')
plt.title('Non buff effect two box model')
plt.xlabel('Year')
plt.ylabel('CO2 Concentration(ppm)')
plt.xticks([1985, 1990, 1995, 2000])
plt.yticks([340, 360, 380, 400, 420, 430])
plt.legend()
plt.show()
```




1.2 [20 points] Following equation 3-4 (with the buffer effect), build a two-box model to compute the atmospheric CO2 level in ppm from 1987 to 2004.

In [6]:

```
def gama(t):
    gama=float(df1.loc[df1['Year']=="int(t)"]['Total carbon emissions from fossil-fuels (million metric tons of C)']/1000/2.13)
    return gama
def zeta(N1):
    zeta=3.69+1.86/100*N1-1.8*(1e-6)*N1*N1
    return (N20)
def WBEM(f, t, k12, k21, N20):
    N1, N2=f
    dfdt=[-k12*N1+k21*(N20+zeta(N1)*(N2-N20))+gama(t), k12*N1-k21*(N20+zeta(N1)*(N2-N20))]
    return dfdt
N20=821/2.13
k12=105/740
k21=102/900
#Initial condition
f0=[740/2.13, 900/2.13]
t=np.linspace(1986, 2004, 19)
f=integrate.odeint(WBEM, f0, t, args=(k12, k21, N20))
N1_WBmodel=f[:,0]
```

In [7]:


```
plt.plot(df2['year'], df2['mean'], 'ok', label='observations')
plt.plot(t[1:19], N1_WBmodel[1:19], 'k', label='calculation with buffer effect')
plt.title('With buff effect two box model')
plt.xlabel('Year')
plt.ylabel('CO2 Concentration(ppm)')
plt.xticks([1985, 1990, 1995, 2000])
plt.yticks([340, 360, 380, 400, 420, 430])
plt.legend()
plt.show()
```



1.3 [5 points] Based on your results from 1.1 and 1.2, reproduce Figure 2 in Tomizuka (2009) as much as you can.

In [8]:

```
#set figure
fig=plt.figure(figsize=(5,3),dpi=500)
ax=fig.add_subplot(1, 1, 1)
#plot data
plt.plot(df2['year'],df2['mean'],'ok',markersize=3)
plt.plot(t[1:19],N1_NBmodel[1:19],'k')
plt.plot(t[1:19],N1_WBmodel[1:19],'k')
#xticks/yticks
plt.xticks([1985,1990,1995,2000])
plt.yticks([360,380,400,420])
#xlim/ylim
plt.xlim(1983,2005)
plt.ylim(338,430)
#xlabel/ylabel
plt.xlabel('Year',fontsize=9)
plt.ylabel('CO2{2}$ Concentration(ppm)',fontsize=9)
ax.tick_params(labelsize=8)
#set border
ax.spines['right'].set_color('none')
ax.spines['top'].set_color('none')
#set axis format
ax.minorticks_on()
ax.tick_params(axis='y',which='major',direction='in',width=1,length=3.5)
ax.tick_params(axis='y',which='minor',direction='in',width=1,length=2)
ax.tick_params(axis='x',which='major',direction='in',width=1,length=3)
ax.tick_params(axis='x',which='minor',direction='in',width=1,length=3.5,color='none')
ax.spines['bottom'].set_position(('data',340))
ax.spines['left'].set_position(('data',1984))
#add text
ax.text(1985,415,'calculation with buffer effect',fontsize=8)
ax.text(1994,350,'calculation without buffer effect',fontsize=8)
ax.text(1999,385,'observations',fontsize=8)
plt.show()
```



In []:

In []:

In []:

In []: