

Likelihood for Seed Production

Growth data is on individual level, while seed data is on stand level, so I will model individual tree radial growth and stand-level seed production by introducing $\mu_{j,sp,y}$, representing the mean radial growth for stand j , species sp , and year y .

For each individual tree i in stand j , species sp , and year y , the observed radial growth is modeled as, with σ being the variance across individuals:

$$\text{Growth}_{i,j,sp,y} \sim \text{Normal}(\mu_{j,sp,y}, \sigma^2)$$

The seed count for a particular stand j , species sp , and year y follows a negative binomial distribution with mean $\lambda_{j,sp,y}$ and dispersion term ϕ :

$$\text{Seed}_{j,sp,y} \sim \text{NegBinomial}(\lambda_{j,sp,y}, \phi)$$

The modeled seed count ($\lambda_{j,sp,y}$) for stand j , species sp , and year y is:

$$\log(\lambda_{j,sp,y}) = \alpha_0 + \alpha_{sp} + \alpha_j + (\beta_{1,sp} + \beta_{1,j}) \cdot \mu_{j,sp,y} + (\beta_{2,sp} + \beta_{2,j}) \cdot \mu_{j,sp,y-1} + \beta_{3,sp} \cdot \text{Elevation}_j$$

Where:

α_0 is the grand mean across all species, stands and years;

α_{sp} is the species-specific intercept for species sp ;

α_j is the stand-specific intercept for stand j ;

$\beta_{1,sp}$ is the slope for the relationship between current-year growth and seed production for species sp ;

$\beta_{1,j}$ is the slope for the relationship between current-year growth and seed production for stand j ;

$\mu_{j,sp,y}$ is the mean growth for stand j , species sp , and year y ;

$\beta_{2,sp}$ is the slope for the relationship between previous-year growth and seed production as the lag effect for species sp ;

$\beta_{2,j}$ is the slope for the relationship between previous-year growth and seed production as the lag effect for stand j ;

$\beta_{3,sp}$ is the effect of elevation on seed production for species sp .

Priors for Random Slopes

$$\beta_{1,\text{sp}} \sim \text{Normal}(0, \sigma_{\beta_{1,\text{sp}}})$$

$$\beta_{2,\text{sp}} \sim \text{Normal}(0, \sigma_{\beta_{2,\text{sp}}})$$

$$\beta_{1,j} \sim \text{Normal}(0, \sigma_{\beta_{1,j}})$$

$$\beta_{2,j} \sim \text{Normal}(0, \sigma_{\beta_{2,j}})$$

Where:

$\sigma_{\beta_{1,\text{sp}}}$ and $\sigma_{\beta_{2,\text{sp}}}$ represent the standard deviations for the random slopes at the species level;
 $\sigma_{\beta_{1,j}}$ and $\sigma_{\beta_{2,j}}$ represent the standard deviations for the random slopes at the stand level.