TITLE: Evolutionary Explanations for Masting: Linking Traits to Different Hypotheses from Pollination to Seed Survival

AUTHORS: Xiaomao Wang[1], E.M. Wolkovich[1]

[1] Forest & Conservation Sciences, Faculty of Forestry, University of British Columbia, 2424 Main Mall, Vancouver, BC V6T 1Z4, Canada

1 Introduction

In 2025, Japan experienced the highest number of bear attacks in the historical record. Recent research has linked the attacks to food deficit because of disruptions in periodic forest food supply, likely a disruption in seed masting (Xiao *et al.*, 2026). Masting is the phenomenon of synchronized production of seeds or other reproductive structures among individuals within the same population (Kelly, 1994; Kelly & Sork, 2002; Pearse *et al.*, 2016), which has important ecological consequences. Masting has long been studied in the context of seed predation (Janzen, 1971; Kelly, 1994). By producing large seed crops in mast years, plants can overwhelm seed predators, increasing the proportion of seeds that escape predation and potentially enhancing seedling recruitment (Kelly & Sork, 2002). Earlier in the reproductive cycle, higher pollination success can lead to more seed production later in the season. Throughout the whole reproductive cycle, reproduction requires substantial resource investment; the resource availability for reproduction as a result may determine the success of a mast event (Crone & Rapp, 2014). Together, masting is shaped by selective pressures and constraints acting at different stages of the whole reproductive cycle.

Considering different reproductive stages, multiple hypotheses have emerged to explain the mechanism of masting: increased pollination efficiency; satiation of seed predators; constraints associated with resource allocation, and environmental cues or vetoes that trigger or interrupt a mast event (Pearse *et al.*, 2016; Bogdziewicz *et al.*, 2019, 2018). Our understanding of masting has advanced considerably based on these hypotheses with more data collected and looking at the masting at a finer scale. Synchronized flowering can enhance the pollination efficiency when there is pollen limitation in the population (Satake & Iwasa, 2000; Pearse *et al.*, 2015; Bogdziewicz *et al.*, 2017). During the seed maturation stage, masting only makes sense when there is a density-dependent advantage to overwhelm seed predators and increase seed survival. When the synchronized reproduction fails in satiating seed predators, the cost of delayed reproduction or missing a good recruitment year will make masting unfavored (Silvertown, 1980; Qiu *et al.*, 2023).

More recent studies have also explored different resource-based explanations to investigate how species differ in their capacity to respond to resource fluctuations and how this will affect their masting capacity (Journé *et al.*, 2023; Fernández-Martínez *et al.*, 2019). Environmental cues can synchronize reproductive activity among individuals within a population for a potential good year for recruitment (Schauber *et al.*, 2002; Moreira *et al.*, 2015), but they cannot, on their own, explain why species differ in the strength of their masting patterns. These mechanisms are not mutually exclusive, and their relative importance may vary among species according to their life history strategies.

It is then important to connect traits and hypotheses with potential evolutionary drivers when considering what shapes the masting behaviour for certain species. Functional traits are closely related to life history strategies (Adler *et al.*, 2014), which provide a potential way to connect theoretical hypotheses of masting to measurable differences among species. Traits associated with different hypotheses may help evaluate at which stage the selection is more important in shaping masting that overall benefits the species. By examining these traits in relation to masting behaviour, it may be possible to evaluate the relative support for different hypotheses and identify the characteristics associated with variation in reproductive synchrony.

In this study, we investigated the relationships between traits and masting behaviour to evaluate the potential mechanisms underlying masting in North American forests. Specifically, we tested whether traits associated with different hypotheses explain differences in masting behaviours among species and

whether these relationships are similar among closely related species. Here we use North American tree to test this framework since ecologically and evolutionarily they are more closely related.

2 Method

2.1 Data Collection

2.1.1 Trait selection

We selected traits based on three hypotheses proposed to explain the evolution of masting. For the predator-related hypothesis, we selected traits associated with seed attractiveness to predators and the potential for surviving in seed caches, including seed weight, fruit size (cone size for gymnosperms), oil content, dispersal mode, and seed dormancy. For the pollination-related hypothesis, we aimed for traits related to higher pollination efficiency: pollination mode and reproductive type. For resource-related hypotheses, we included traits responsible for higher flexibility in resource use under varying environmental conditions.

2.1.2 Data scraping

For our analyses, we obtained the full species list from the two volumes of *Silvics of North America* (Burns & Honkala, 1990a,b) (hereafter referred as *Silvics*). We extracted masting data primarily from *Silvics*. Pollination mode and seed dispersal data were also obtained mainly from *Silvics*. We obtained seed weight data from the Seed Information Database by the Society for Ecological Restoration (Society for Ecological Restoration *et al.*, 2023), gathered seed dormancy data from *Seeds* by Baskin and Baskin (2014) and accessed the leaf longevity and oil content data through the TRY database (Kattge *et al.*, 2020). We compiled fruit size and drought tolerance data from multiple sources, including: *Silvics*, the USDA Fire Effects Information System (United States Department of Agriculture, Forest Service, 2026), the USA National Phenology Network (USA National Phenology Network, 2026), and The Gymnosperm Database (Earle, 2026).

We defined masting based on qualitative descriptions of seed production patterns. Although recent studies emphasize quantitative definitions of masting, many forestry manuals view masting as a species-level categorical trait, including *Silvics*. Thus, we classified species according to key phrases describing their seed production patterns. Species described as producing "a good crop almost every year" were classified as weak masting species, whereas species described as producing "a good crop every several years" or having "irregular seed production" were classified as strong masting species.

For some statistical analyses we grouped or otherwise adjusted the data on traits related to masting hypotheses. For numerical traits with a range (seed weight and fruit size), we recorded the full range from the source and took the mean value for further analyses. We took the log form of seed weight and fruit size because the range of data is large. For categorical traits (pollination mode, reproductive type, and dispersal mode), we grouped them into broader categories; for example, we classified pollination mode into wind, animal (birds, bats, or insects), or both. The complete raw and processed datasets are available through KNB.

2.2 Statistical Analysis

2.2.1 Masting pattern and single trait

To analyze the relationship between masting pattern and individual traits, we used logistic regression, treating masting as a binary response variable. Given our species-level analyses, we used a phylogenetic logistic regression approach (`phyloglm` function in the `phylolm` package; v. 2.6.2). To construct the phylogenetic tree displaying the updated phylogenetic structure of our species, we verified all species names using the `WorldFlora` package (v. 1.14-5) and updated all species names according to WCVP Backbone data (2025). We then constructed an ultrametric phylogenetic tree in R using the `phytools` (v. 2.4-4) and `ape` (v. 5.7-1) packages. To test the phylogenetic correction for each trait, we calculated the half-life using the equation $(\log(2)/\alpha)$ and compared it with the tree height (standardized to 1). This

metric represents the time required for the influence of ancestral states to decrease by half. For more interpretation, see Cooper *et al.* (2016)

2.2.2 Masting pattern with a multivariate analysis

In addition to these univariate analyses looking at the relationship between masting and each individual trait, we also conducted a multivariate analysis to test whether traits related to any single hypothesis is a stronger driver of the masting pattern, using non-metric multidimensional scaling (NMDS, implemented using `metaMDS` function in the `vegan` v. 2.7-1 package). We evaluated both two- and three-dimentional solutions $k = 2, 3$ and selected $k = 3$ based on the lower stress score. To check for potential patterns, we plotted trait distributions on two panels (NMDS1 vs. NMDS2 and NMDS2 vs. NMDS3).

2.2.3 Phylogenetic signal for individual traits

To check for the phylogenetic signal for all traits including masting, we used different metrics depending on the type of data. For traits that were binary, we calculated Fritz and Purvis' D-statistics using the function `phylo.d` in the `caper` (v. 1.0.4) package. For categorical traits, we estimated the Pagel's λ using the function `fitDiscrete` in the `geiger` (v. 2.0.11) package. For continuous traits, we calculated Pagel's λ using the function `phylosig` in the `phytools` (v. 2.4-4) package.

3 Results

3.0.1 Structure of Data

In total, our dataset included 188 species, comprising 124 angiosperm species and 64 gymnosperm species (cite PHYLOfig). Among angiosperms, 79% (98/124) of species had masting data available, with strong masting and weak masting species occurring in approximately equal proportions 52% (51/98). Among gymnosperms, 95% (61/64) of species had masting data available, and nearly all were classified as strong masting species 89% (54/61).

We were not able to find complete trait data for every single trait for all species with masting data. The traits with the most available data was seed weight. The traits with the sparse data were oil content and leaf longevity (Fig. 1 and Fig. 2).

3.0.2 Results for Each Hypothesis

We found relationships between traits related to predator satiation and pollination efficiency and strong masting for angiosperm species. Among the traits related to predator satiation, seed weight was the only trait significantly related to masting (p-value = 0.02, Table S1). Larger seed weight was associated with a greater probability of strong masting (Figure 1). For angiosperms only, increasing seed weight threefold (from a median seed weight of 90.9mg to 272.7mg) increased the predicted likelihood of strong masting from 0.44 to 0.5 (Figure 3). The effect was stronger among species with biotically dispersed seeds, where increasing seed weight threefold (from a median seed weight of 1087.15mg to 3261.45mg) increased the predicted likelihood of strong masting from 0.73 to 0.81 (Table S3; Figure 3).

Among traits related to pollination efficiency, pollination mode was the only trait significantly associated with strong masting (p-value = 0.02, Table S1) for angiosperm. Wind-pollinated species had a predicted likelihood of strong masting of 0.59, compared with 0.2 for animal-pollinated species. All gymnosperm species in our dataset were wind-pollinated, and gymnosperms contained a much higher proportion of strong masting species overall, which may also indicate that wind pollination is associated with strong masting.

3.0.3 Multivariate analysis

Strong and weak masting species were not clearly separated along any axis in the NMDS analysis (Fig. 5). We found no evidence that combinations of traits associated with any of the three hypotheses explained variation in masting pattern.

3.0.4 Phylogenetic signal

Masting pattern showed a moderate phylogenetic signal in both gymnosperms (D = 0.51) and angiosperms (D = 0.58) (Table S5), which indicated that related species are more likely to exhbit similar masting pattern.

Seed weight had strong phylogenetic signal for both angiosperm ($\lambda = 0.96$) and gymnosperm ($\lambda = 0.98$). This pattern is also evident in Fig. 4. Dispersal mode, leaf longevity, pollination mode and reproductive type also showed high phylogenetic signal in Table S4. However, these values should be interpreted cautiously because Pagel's λ is estimated differently for categorical and continuous traits, making direct comparisons inappropriate.

4 Discussion

We found that masting was associated with traits related to both pollination and seed dispersal, providing support for multiple mechanisms underlying interannual variation in seed production. The positive relationship between wind pollination and strong masting in angiosperm supports the assumption that wind-pollinated species benefit more from the synchronized flowering (Pearse *et al.*, 2015). The positive relationship between seed weight and strong masting provides evidence for the predator satiation hypothesis. These associations point to selective pressure acting at both early and late stages of the reproductive cycle, which are critical for successful reproduction and gene transmission.

Previous literature has found a positive relationship between seed mass and masting, with one explanation being that larger seeds require more resource investment and simply cannot reproduce every year (Kelly, 1994). In contrast, a more recent other study reported the opposite relationship between seed mass and masting, suggesting that small seeds are usually associated with seed bank which reduce the fitness costs associated with years of low or failed reproduction (Journé *et al.*, 2023). We found a weak but significant positive relationship between seed mass and masting, suggesting that the association between these traits may reflect processes beyond the resource costs of producing large seeds. For predator satiation to favour synchronized reproduction, seed crops must provide a sufficiently large resource to overwhelm seed consumers (Janzen, 1971; Kelly, 1994).

In some species, seed predators also function as seed dispersers. Larger seeds may provide greater rewards to animal dispersers and therefore facilitate dispersal of these species. At the same time, larger seeds generally contain greater nutritional reserves, which can increase the probability of successful germination and establishment following dispersal, even in the case when seeds experience some degree of predation or damage (Bonal *et al.*, 2007; Moles & Westoby, 2004). Thus, large seeds may provide benefits at multiple stages following seed production, from attracting dispersers to supporting seedling establishment.

Seed size and seed crop size are also often subject to a trade-off because greater investment in individual seeds can reduce the number of seeds(Castro *et al.*, 2024; Parciak, 2002). Masting may partially relieve this constraint by concentrating reproduction into episodic years, allowing a species to produce a large number of relatively costly seeds while at the same time reducing the proportion of seeds consumed (Jansen *et al.*, 2004; Bonal *et al.*, 2007). Together, these processes could generate selection favouring reproductive strategies that enhance seed dispersal and survival.

4.1 Scale Dependence in Trait–Masting Relationships

Recent work has emphasized that masting is better represented as a continuous rather than categorical trait, which is mostly available at the population level (Bogdziewicz *et al.*, 2024, 2025). The coefficient of variation (CV) is one of the most widely used continuous measures of masting. However, CV might not be a good measurement for masting in some cases, as it does not capture all characteristics of masting and can be strongly distorted by years with zero seed production (Bogdziewicz *et al.*, 2023). At the same time, evaluating the evolutionary associations between traits and masting requires comparisons across species. If a trait has a sufficiently strong and consistent association with the evolution of masting, we expect this relationship to be detectable at the species level using binary masting data.

Many of the traits considered in our analysis are phenotypically plastic and may vary among populations, potentially in response to local environmental conditions (Matesanz *et al.*, 2010). Thus, traits for which we failed to detect a species-level association with masting may still be potentially important at the population level. People have found that some species only exhibit masting under specific environmental conditions (Fleurot *et al.*, 2023; Bogdziewicz *et al.*, 2025). As a result, understanding how reproductive traits are associated with masting may require measurements of both traits and reproductive variation at the population level, considering local environmental conditions and intraspecific variation.

Masting has originally been studied primarily as interannual variation in seed production, but we can refine the definition to seed-maturation masting and flower masting (Satake & Kelly, 2021; Pearse *et al.*, 2016). These two forms of reproductive synchrony occur at different stages of the reproductive cycle: seed masting focuses on the final reproductive outcome, whereas flower masting occurs at the beginning of reproduction. Importantly, variation in flowering and seed production does not necessarily occur at the same time (Pearse *et al.*, 2015; Sork, 1993; Isagi *et al.*, 1997), suggesting that these two forms of masting may be shaped by different selective pressures. For example, we would expect mechanisms related to pollination to perform stronger selection on flowering synchrony, whereas mechanisms related to seed predation, dispersal and establishment may be more directly associated with synchrony in seed production. In our study, we have more traits related to predation, which might not allow us to reveal any relationship between the traits related to flowering masting. For example, gymnosperm are all wind-pollinated and we did not find strong relationship between any traits and masting, which could because that the masting mechanism for gymnosperm species are more limited to the pollination phase, however, we did not have many distinctive flower traits included in this study to selected that out. Separating species according to their dominant form of masting nad including more flower traits could therefore provide a more refined test of hypotheses underlying reproductive synchrony, and help understand how different selective pressures contribute to the evolution of reproductive synchrony at different stages of the reproductive cycle.

4.2 Phylogenetic Signal Limits Inference

Gymnosperms and angiosperms occupy fundamentally different functional spaces and might potentioally have different mechanisms shaping their masting behaviors. In our dataset, the proportion of species exhibiting strong masting is greater in gymnosperm. The leaf longevity in general is longer for most gymnosperm species compared to angiosperm species and wind pollination is also more prevalent. The abundance of certain traits and masting in gymnosperm might indicate this is a more important and efficient reproductive strategy for them, but not necessarily for more diverse angiosperm. There might be interests to test if masting was gradually diluted in angiosperm as they transitioned to biotic pollination and faster life histories. We found strong phylogenetic signal for many traits we selected including masting. For the analysis we did, it is hard to tease out effect of phylogeny from the correlation between single trait and masting, because there might be the possibility that the relationship between masting and these traits is because of the phylogeny structure rather than one acts as a driver for another.bAlso, many traits we included have relatively sparse data, and these data are nested in certain clade. This makes the phylogenetic bias a potentially bigger problem.

It is also important to look for the evidence of evolution separately for angiosperm and gymnosperm. Gymnosperms and angiosperms occupy fundamentally different functional spaces.

Given that the evolutionary mechanisms underlying masting may differ among lineages and operate at different biological scales, investigating evolutionary patterns within clades is an important next step. Previous studies have highlighted the potential of finer scale comparisons to investigate the evolution of masting with in clade or even between species in the same genus(Satake & Kelly, 2021; Koenig *et al.*, 1996; Bogdziewicz *et al.*, 2025). Clades that contain species exhibiting both strong masting and weak masting such as *Quercus* and *Pinus*, offer particularly valuable opportunities to investigate how traits and selective pressures are associated with variation in masting. To achieve this, we need more comprehensive data on both species-level and population-level for comparison. Such an approach would help our understanding of masting evolution by filling the gaps on how masting varies among populations in response to contemporary ecological conditions and how reproductive strategies have diverged over longer evolutionary timescales. Integrating all these perspectives together will provide a more comprehensive

and comparable understanding of how ecological conditions, life-history traits and evolutionary history jointly shape the diversity of reproductive strategies.




Figure 1: How plant traits related to different hypotheses differ in strong masting and weak masting for angiosperm species: We show data for traits related to the predator satiation hypothesis (a-e), pollination-related hypothesis (f-g) and the resource-related hypothesis (h-i) as either violin plots (a-c, i) or proportional histograms (d-h) separated by species categorized as strong masting (gold) or weak masting (teal). We show model summaries only where significant (P-value < 0.5) and also report the model sample size (N), regression coefficient (Estimate), and phylogenetic signal parameter α. Full model results are show in Table S1.




Figure 2: How plant traits related to different hypotheses differ in strong masting and weak masting for gymnosperm species: We show data for traits related to the predator satiation hypothesis (a-e), pollination-related hypothesis (f-g) and the resource-related hypothesis (h-i) as either violin plots (a-c, i) or proportional histograms (d-h) separated by species categorized as strong masting (gold) or weak masting (teal). We show model summaries only where significant (P-value < 0.5) and also report the model sample size (N), regression coefficient (Estimate), and phylogenetic signal parameter α. Full model results are show in Table S2.




Figure 3: Predicted likelihood of masting as a function of seed weight for all angiosperm species (solid black line) and the biotic-dispersed subgroup (dashed mint line), estimated using phylogenetic logistic regression. Individual species observations are plotted as binary data points at 0 (weak masting) and 1 (strong masting).



Figure 4: Phylogenetic tree for all angiosperm species included in the analysis. The colour scheme of the branch is the trait value for seed weight (log), which showed strong phylogenetic signal (Table S4). The tip labels are coloured by strong masting or weak masting. The other traits are labelled after the tip labels with different colour for category and size for magnitude.




Figure 5: Non-metric multidimensional scaling (NMDS) of trait composition for angiosperm and gymnosperm species. Each point represents a species, with position determined by NMDS axes. Points are coloured by drought tolerance, filled by strong masting or not, shaped by reproductive type, sized by seed weight. Panels are arranged as (a) angiosperm NMDS1–NMDS2, (b) angiosperm NMDS2–NMDS3, (c) gymnosperm NMDS1–NMDS2, and (d) gymnosperm NMDS2–NMDS3.

References

Adler, P.B., Salguero-Gómez, R., Compagnoni, A., Hsu, J.S., Ray-Mukherjee, J., Mbeau-Ache, C. & Franco, M. (2014) Functional traits explain variation in plant life history strategies. *Proceedings of the National Academy of Sciences* **111**, 740–745.

Bogdziewicz, M., Calama, R., Courbaud, B., Espelta, J.M., Hacket-Pain, A., Journé, V., Kunstler, G., Steele, M., Qiu, T., Zywiec, M. *et al.* (2023) How to measure mast seeding? *New Phytologist* **239**, 830–838.

Bogdziewicz, M., Kelly, D., Ascoli, D., Caignard, T., Chianucci, F., Crone, E.E., Fleurot, E., Foest, J.J., Gratzer, G., Hagiwara, T. *et al.* (2024) Evolutionary ecology of masting: mechanisms, models, and climate change. *Trends in Ecology & Evolution* **39**, 851–862.

Bogdziewicz, M., Kelly, D., Zwolak, R., Szymkowiak, J. & Hacket-Pain, A. (2025) Dynamics, mechanisms, and consequences of mast seeding. *Annual Review of Ecology, Evolution, and Systematics* **56**.

Bogdziewicz, M., Steele, M.A., Marino, S. & Crone, E.E. (2018) Correlated seed failure as an environmental veto to synchronize reproduction of masting plants. *New Phytologist* **219**, 98–108.

Bogdziewicz, M., Szymkowiak, J., Kasprzyk, I., Grewling, Ł., Borowski, Z., Borycka, K., Kantorowicz, W., Myszkowska, D., Piotrowicz, K., Ziemianin, M. *et al.* (2017) Masting in wind-pollinated trees: system-specific roles of weather and pollination dynamics in driving seed production. *Ecology* **98**, 2615–2625.

Bogdziewicz, M., Żywiec, M., Espelta, J.M., Fernández-Martinez, M., Calama, R., Ledwoń, M., McIntire, E. & Crone, E.E. (2019) Environmental veto synchronizes mast seeding in four contrasting tree species. *The American Naturalist* **194**, 246–259.

Bonal, R., Munoz, A. & Díaz, M. (2007) Satiation of predispersal seed predators: the importance of considering both plant and seed levels. *Evolutionary Ecology* **21**, 367–380.

Burns, R.M. & Honkala, B.H. (eds.) (1990a) *Silvics of North America*, vol. 1 of *Agriculture Handbook*. United States Department of Agriculture, Forest Service, Washington, DC.

Burns, R.M. & Honkala, B.H. (eds.) (1990b) *Silvics of North America*, vol. 2 of *Agriculture Handbook*. United States Department of Agriculture, Forest Service, Washington, DC.

Castro, J., Hódar, J.A. & Gómez, J.M. (2024) Seed size. *Handbook of seed science and technology*, pp. 397–427, CRC Press.

Cooper, N., Thomas, G.H., Venditti, C., Meade, A. & Freckleton, R.P. (2016) A cautionary note on the use of ornstein uhlenbeck models in macroevolutionary studies. *Biological journal of the Linnean Society* **118**, 64–77.

Crone, E.E. & Rapp, J.M. (2014) Resource depletion, pollen coupling, and the ecology of mast seeding. *Annals of the New York Academy of Sciences* **1322**, 21–34.

Earle, C.J. (2026) The gymnosperm database. `https://www.conifers.org/`, accessed 2026-05-13.

Fernández-Martínez, M., Pearse, I., Sardans, J., Sayol, F., Koenig, W.D., LaMontagne, J.M., Bogdziewicz, M., Collalti, A., Hacket-Pain, A., Vacchiano, G. *et al.* (2019) Nutrient scarcity as a selective pressure for mast seeding. *Nature Plants* **5**, 1222–1228.

Fleurot, E., Lobry, J.R., Boulanger, V., Debias, F., Mermet-Bouvier, C., Caignard, T., Delzon, S., Bel-Venner, M.C. & Venner, S. (2023) Oak masting drivers vary between populations depending on their climatic environments. *Current Biology* **33**, 1117–1124.

Isagi, Y., Sugimura, K., Sumida, A. & Ito, H. (1997) How does masting happen and synchronize? *Journal of Theoretical Biology* **187**, 231–239.

Jansen, P.A., Bongers, F. & Hemerik, L. (2004) Seed mass and mast seeding enhance dispersal by a neotropical scatter-hoarding rodent. *Ecological Monographs* **74**, 569–589.

Janzen, D.H. (1971) Seed predation by animals. *Annual review of ecology and systematics* **2**, 465–492.

Journé, V., Hacket-Pain, A. & Bogdziewicz, M. (2023) Evolution of masting in plants is linked to investment in low tissue mortality. *Nature communications* **14**, 7998.

Kattge, J., Bönisch, G., Díaz, S., Lavorel, S., Prentice, I.C., Leadley, P., Tautenhahn, S., Werner, G.D., Aakala, T., Abedi, M. *et al.* (2020) Try plant trait database–enhanced coverage and open access. *Global change biology* **26**, 119–188.

Kelly, D. (1994) The evolutionary ecology of mast seeding. *Trends in ecology & evolution* **9**, 465–470.

Kelly, D. & Sork, V.L. (2002) Mast seeding in perennial plants: why, how, where? *Annual review of ecology and systematics* **33**, 427–447.

Koenig, W.D., Knops, J.M., Carmen, W.J., Stanback, M.T. & Mumme, R.L. (1996) Acorn production by oaks in central coastal california: influence of weather at three levels. *Canadian Journal of Forest Research* **26**, 1677–1683.

Matesanz, S., Gianoli, E. & Valladares, F. (2010) Global change and the evolution of phenotypic plasticity in plants. *Annals of the new York Academy of Sciences* **1206**, 35–55.

Moles, A.T. & Westoby, M. (2004) Seedling survival and seed size: a synthesis of the literature. *Journal of Ecology* **92**, 372–383.

Moreira, X., Abdala-Roberts, L., Linhart, Y.B. & Mooney, K.A. (2015) Effects of climate on reproductive investment in a masting species: assessment of climatic predictors and underlying mechanisms. *Journal of Ecology* **103**, 1317–1324.

Parciak, W. (2002) Seed size, number, and habitat of a fleshy-fruited plant: consequences for seedling establishment. *Ecology* **83**, 794–808.

Pearse, I.S., Koenig, W.D., Funk, K.A. & Pesendorfer, M.B. (2015) Pollen limitation and flower abortion in a wind-pollinated, masting tree. *Ecology* **96**, 587–593.

Pearse, I.S., Koenig, W.D. & Kelly, D. (2016) Mechanisms of mast seeding: resources, weather, cues, and selection. *New Phytologist* **212**, 546–562.

Qiu, T., Aravena, M.C., Ascoli, D., Bergeron, Y., Bogdziewicz, M., Boivin, T., Bonal, R., Caignard, T., Cailleret, M., Calama, R. *et al.* (2023) Masting is uncommon in trees that depend on mutualist dispersers in the context of global climate and fertility gradients. *Nature Plants* **9**, 1044–1056.

Satake, A. & Iwasa, Y. (2000) Pollen coupling of forest trees: forming synchronized and periodic reproduction out of chaos. *Journal of theoretical biology* **203**, 63–84.

Satake, A. & Kelly, D. (2021) Studying the genetic basis of masting. *Philosophical Transactions of the Royal Society B: Biological Sciences* **376**.

Schauber, E.M., Kelly, D., Turchin, P., Simon, C., Lee, W.G., Allen, R.B., Payton, I.J., Wilson, P.R., Cowan, P.E. & Brockie, R. (2002) Masting by eighteen new zealand plant species: the role of temperature as a synchronizing cue. *Ecology* **83**, 1214–1225.

Silvertown, J.W. (1980) The evolutionary ecology of mast seeding in trees. *Biological journal of the Linnean Society* **14**, 235–250.

Society for Ecological Restoration, International Network for Seed-based Restoration & Royal Botanic Gardens, Kew (2023) Seed information database (sid). `https://ser-sid.org/`, accessed 2026-05-13.

Sork, V.L. (1993) Evolutionary ecology of mast-seeding in temperate and tropical oaks (quercus spp.). *Vegetatio* **107**, 133–147.

United States Department of Agriculture, Forest Service (2026) Fire effects information system (feis). `https://www.feis-crs.org/feis/`, accessed 2026-05-13.

USA National Phenology Network (2026) Usa national phenology network. `https://www.usanpn.org/`, accessed 2026-05-13.

Xiao, H., Zhang, Z., Miyamoto, Y. & Ichinose, T. (2026) Compound food failure on a deteriorated mast system drives japan's bear conflict crisis. *Science Advances* **12**, eaej8134.

A Supplementary Material

Trait	Type	Predictor	N	Estimate	SE	P	Half-life
Seed weight (log)	Numeric	Seed weight (log)	91	0.21	0.09	0.02	0.08
Fruit size (log)	Numeric	Fruit size (log)	88	0.08	0.31	0.79	0.04
Oil content (%)	Numeric	Oil content (%)	19	0.04	0.03	0.20	0.04
Seed dormancy	Categorical	Dormant vs. non-dormant	85	0.05	0.51	0.92	0.06
Dispersal mode	Categorical	Biotic vs. abiotic	93	0.53	0.53	0.32	0.09
Dispersal mode	Categorical	Both vs. abiotic	93	0.53	0.57	0.35	0.09
Pollination mode	Categorical	Wind vs. animal	52	1.72	0.72	0.02	0.01
Pollination mode	Categorical	Wind vs. animal and animals	52	1.18	1.15	0.31	0.01
Reproductive type	Categorical	Monoecious vs. dioecious	89	0.76	0.68	0.26	0.05
Reproductive type	Categorical	Monoecious vs. polygamous	89	-0.29	0.96	0.76	0.05
Drought tolerance	Categorical	Low vs. high drought tolerance	79	0.15	0.47	0.76	0.06
Drought tolerance	Categorical	Moderate vs. high drought tolerance	79	-0.15	0.45	0.74	0.06
Leaf longevity (years)	Numeric	Leaf longevity (years)	16	0.26	0.95	0.78	0.77

Table S1: Phylogenetic generalized linear model results for angiosperm species. We modelled each trait separately with masting as a binary variable and traits as either categorical or continuous (Type column). For categorical traits, we report the difference in log-odds between a given level and its designated baseline reference level; for continuous traits, we report the change in log-odds per unit increase of the predictor (Estimate column). We also report the standard error (SE), P-value (with $P < 0.05$ considered significant and marked with an asterisk next to the Predictor) and the half-life. The half-life represents the time required for the influence of ancestral states to decrease by half; where larger values suggests the residuals are more affected by phylogenetic history than the environment (and vice versa for smaller values). Given our trees are scaled to 1, a value near 1 thus means phylogenetic correlation persists across the phylogenetic depth. For more interpretation, see Cooper *et al.* (2016).

Trait	Type	Predictor	N	Estimate	SE	P	Half-life
Seed weight (log)	Numeric	Seed weight (log)	59	0.01	0.24	0.98	0.01
Fruit size (log)	Numeric	Fruit size (log)	60	0.03	0.36	0.94	0.01
Seed dispersal	Categorical	Biotic vs. abiotic	60	-0.83	0.81	0.30	0.36
Seed dispersal	Categorical	Both vs. abiotic	60	-1.11	0.67	0.10	0.36
Seed dormancy	Categorical	Dormant vs. non-dormant	59	0.69	0.69	0.32	0.01
Reproductive type	Categorical	Monoecious vs. dioecious	61	-1.13	1.01	0.26	0.78
Drought tolerance	Categorical	Low vs. high drought tolerance	52	11.36	98.16	0.91	34.66
Drought tolerance	Categorical	Moderate vs. high drought tolerance	52	-0.29	0.60	0.63	34.66
Leaf longevity (years)	Numeric	Leaf longevity (years)	38	0.15	0.35	0.67	0.01

Table S2: Phylogenetic generalized linear model results for gymnosperm species. We modelled each trait separately with masting as a binary variable and traits as either categorical or continuous (Type column). For categorical traits, we report the difference in log-odds between a given level and its designated baseline reference level; for continuous traits, we report the change in log-odds per unit increase of the predictor (Estimate column). We also report the standard error (SE), P-value (with $P < 0.05$ considered significant and marked with an asterisk next to the Predictor) and the half-life. The half-life represents the time required for the influence of ancestral states to decrease by half; where larger values suggests the residuals are more affected by phylogenetic history than the environment (and vice versa for smaller values). Given our trees are scaled to 1, a value near 1 thus means phylogenetic correlation persists across the phylogenetic depth, and values much higher suggest it persists far deeper than the root of our tree. For more interpretation, see Cooper *et al.* (2016).

Trait	Group	Predictor	N	Estimate	SE	P	Half-life
Seed weight (log)	Biotic and both	Seed weight (log)	55	0.41	0.16	0.01	0.01
Seed weight (log)	Abiotic	Seed weight (log)	36	0.12	0.13	0.36	0.01

Table S3: Phylogenetic generalized linear model results for only seed weight for biotic-dispersed subgroup (both group included) and abiotic-dispersed subgroup. We report the change in log-odds per unit increase of the predictor (Estimate column). We also report the standard error (SE), P-value (with $P < 0.05$ considered significant and marked with an asterisk next to the Predictor) and the half-life (for a description of half-life, see caption of table S1 and S2).

Group	Trait	λ
Angiosperm	Seed weight (Log)	0.95
Angiosperm	Fruit size (Log)	0.93
Angiosperm	Oil content	0.25
Angiosperm	Leaf longevity	1.02
Angiosperm	Dispersal mode	1.00
Angiosperm	Pollination mode	1.00
Angiosperm	Reproductive type	0.00
Angiosperm	Drought tolerance	0.00
Gymnosperm	Seed weight (Log)	0.98
Gymnosperm	Fruit size (Log)	0.91
Gymnosperm	Oil content %	0.00
Gymnosperm	Leaf longevity	0.90
Gymnosperm	Dispersal mode	0.00
Gymnosperm	Reproductive type	1.00
Gymnosperm	Drought tolerance	0.00

Table S4: Phylogenetic Signal (Pagel's λ) for continuous traits and multi-level categorical traits. $\lambda = 0$ indicates no phylogenetic signal, trait distributed independent of the phylogeny; $0 < \lambda < 1$ means moderate phylogenetic signal, trait is partially structured by the phylogeny; $\lambda = 1$ indicates strong phylogenetic signal under a Brownian motion of evolution.

Group	Trait	Estimated D	P(random)	P(Brownian)
Angiosperm	Masting	0.58	0	0.01
Angiosperm	Seed dormancy	0.25	0	0.16
Gymnosperm	Masting	0.51	0.02	0.10
Gymnosperm	Seed dormancy	0.98	0.41	0

Table S5: Phylogenetic Signal (D-statistic) for the phylogenetic signal of binary traits, with probabilities calculated under a random phylogenetic structure and under a Brownian motion phylogenetic structure. $D = 1$ indicates the trait is randomly distributed across the phylogeny; $D = 0$ indicates the trait follows a Brownian motion model; $0 < D < 1$ indicatesmoderate phylogenetic signal; $D < 0$ indicates the trait is more clumped than expected under Brownian motion; and $D > 1$ indicates overdispersion. P random is the p-value for the null hypothesis that the trait is randomly distributed across the phylogeny (no phylogenetic signal), and P Brownian is the p-value for the null hypothesis that the trait follows a Brownian motion model (strong phylogenetic signal). A low p-value indicates rejection of the corresponding null hypothesis.