

# A review of candidate therapies for Middle East respiratory syndrome from a molecular perspective

Ali A. Rabaan,<sup>1,\*</sup> Shamsah H. Alahmed,<sup>2</sup> Ali M. Bazzi<sup>3</sup> and Hatem M. Alhani<sup>4</sup>

## Abstract

There have been 2040 laboratory-confirmed cases of Middle East respiratory syndrome coronavirus (MERS-CoV) in 27 countries, with a mortality rate of 34.9%. There is no specific therapy. The current therapies have mainly been adapted from severe acute respiratory syndrome (SARS-CoV) treatments, including broad-spectrum antibiotics, corticosteroids, interferons, ribavirin, lopinavir–ritonavir or mycophenolate mofetil, and have not been subject to well-organized clinical trials. The development of specific therapies and vaccines is therefore urgently required. We examine existing and potential therapies and vaccines from a molecular perspective. These include viral S protein targeting; inhibitors of host proteases, including TMPRSS2, cathepsin L and furin protease, and of viral M(pro) and the PL(pro) proteases; convalescent plasma; and vaccine candidates. The Medline database was searched using combinations and variations of terms, including 'Middle East respiratory syndrome coronavirus', 'MERS-CoV', 'SARS', 'therapy', 'molecular', 'vaccine', 'prophylactic', 'S protein', 'DPP4', 'heptad repeat', 'protease', 'inhibitor', 'anti-viral', 'broad-spectrum', 'interferon', 'convalescent plasma', 'lopinavir ritonavir', 'antibodies', 'antiviral peptides' and 'live attenuated viruses'. There are many options for the development of MERS-CoV-specific therapies. Currently, MERS-CoV is not considered to have pandemic potential. However, the high mortality rate and potential for mutations that could increase transmissibility give urgency to the search for direct, effective therapies. Well-designed and controlled clinical trials are needed, both for existing therapies and for prospective direct therapies.

## INTRODUCTION

### Middle East respiratory syndrome coronavirus overview

Middle East respiratory syndrome coronavirus (MERS-CoV) was first isolated in Jeddah in the Kingdom of Saudi Arabia (KSA) from a 60-year-old male hospital patient, who died 24 June 2012, 11 days after presenting with acute pneumonia and subsequent renal failure [1]. Since then, the WHO have been notified of 2040 laboratory-confirmed cases, including 712 deaths [2]. While most cases have arisen in the Middle East, cases have also emerged in 27 countries worldwide in travellers from the Middle East and/or in their contacts [2].

Most human MERS-CoV infections are considered to be the result of multiple zoonotic transfers. Bats are the most likely MERS-CoV natural reservoir, as with other mammalian coronaviruses (CoVs), while camels are likely to be the major zoonotic source for human infections [3–5]. Secondary human-to-human transmission is considered to be limited, occurring mainly within family and healthcare settings. The first cluster of cases in humans was retrospectively identified to have occurred in a public hospital in Jordan in April 2012 [6]. Multiple healthcare facility-associated outbreaks have since occurred in the Middle East, most notably in KSA, often linked to deficiencies in infection control procedures [7–13]. Although cases outside the Middle East have mainly been isolated, a large outbreak occurred in

Received 8 May 2017; Accepted 21 July 2017

**Author affiliations:** <sup>1</sup>Molecular Diagnostic Laboratory, Johns Hopkins Aramco Healthcare, Dhahran 31311, Saudi Arabia; <sup>2</sup>Specialty Paediatric Medicine, Qatif Central Hospital, Qatif 32654, Saudi Arabia; <sup>3</sup>Microbiology Laboratory, Johns Hopkins Aramco Healthcare, Dhahran 31311, Saudi Arabia; <sup>4</sup>Maternity and Children Hospital, and Directorate of Infection Control at Eastern Province, Ministry of Health, Dammam, Saudi Arabia.

**\*Correspondence:** Ali A. Rabaan, arabaan@gmail.com or ali.rabaan@jhah.com

**Keywords:** Antiviral peptide; camostat; convalescent plasma; DPP4; GLS-5300; glycopeptide antibiotic; interferon; lopinavir; MERS-CoV; monoclonal antibodies; M(pro); PL(pro); protease; S protein; vaccine.

**Abbreviations:** 6HB, six helix bundle; 6MP, 6-mercaptopurine; 6TG, 6-thioguanine; ACE-2, angiotensin-converting enzyme 2; ARDS, acute respiratory distress syndrome; CoVs, coronaviruses; DPP4, dipeptidyl peptidase 4; E, envelope; ExoN, exoribonuclease; FDA, US Food and Drug Administration; HIVIG, hyperimmune IV immunoglobulin; ICU, intensive care unit; IRF, interferon regulatory transcription factor; IRF-3, IFN regulatory factor 3; ISARIC-WHO, International Severe Acute Respiratory and Emerging Infection Consortium; ISRE, interferon-stimulated response element; IVIG, intravenous immunoglobulin; M, membrane; mAbs, monoclonal antibodies; MERS-CoV, Middle East respiratory syndrome coronavirus; N, nucleocapsid; NF, nuclear factor; NSPs, non-structural proteins; PHE, Public Health England; RBD, receptor-binding domain; RBM, receptor-binding motif; RCTs, randomized control trials; S, spike; SARS-CoV, severe acute respiratory syndrome; TLR-3, Toll-like receptor-3; WHO, World Health Organization.

Korea in June 2015, in which human–human transmission resulted in 186 cases and 36 deaths [14]. Increased vulnerability to either cross-species or trans-human transmission could result from viral adaptations [15].

MERS-CoV infection is often accompanied by acute viral pneumonia, and sometimes gastrointestinal symptoms. Clinical severity varies from asymptomatic to death, and the extent of asymptomatic spread is unclear. The high mortality rate is mainly accounted for by acute respiratory distress syndrome (ARDS) [7, 15, 16]. Higher mortality is observed among vulnerable patients, such as older individuals and those suffering from comorbid illness, and is also associated with high viral load [7, 11, 15, 17]. In one study in a KSA hospital, intensive care unit (ICU) admission among MERS-CoV patients was associated with a mortality rate of 74.2 % [11].

While MERS-CoV is not currently considered to have pandemic potential, it is clear that human–human transmission does occur. The exact mechanisms by which MERS-CoV is transmitted from animals to humans have not been fully elucidated. In the South Korean outbreak, the virus emerged in second- and third-generation contacts, resulting in the first human case to be imported into China [18]. This raised concern that viral mutations were contributing to human–human transmission.

Given its high mortality and poor outcomes for vulnerable patients, and the potential for viral mutations, there is no room for complacency in the search for therapeutic options for MERS-CoV. There is currently no specific therapy. Many of the therapeutic options used have been adapted from approaches used to treat severe acute respiratory syndrome (SARS-CoV) during the outbreak of 2003, and/or the H1N1 influenza virus during the outbreak of 2009 [19]. However, while MERS-CoV and SARS-CoV are phylogenetically related betacoronaviruses, they differ in many important respects. MERS-CoV utilizes human dipeptidyl peptidase 4 (DPP4; CD26) receptors, with binding mediated by the viral spike (S) protein, not the angiotensin-converting enzyme 2 (ACE-2) receptors used by SARS [20–24]. MERS-CoV also has a wider cellular tropism [24–26].

Therapies currently used include broad-spectrum antibiotics, corticosteroids and anti-viral treatments, such as interferons (IFN), ribavirin, lopinavir–ritonavir, or mycophenolate mofetil [19, 22, 27–31]. However, the efficacy and/or safety of many of these therapies is unclear, and none are specific to MERS-CoV. Ribavirin monotherapy, for example, is associated with multiple side-effects in the treatment of other viral illnesses, including SARS-CoV, has uncertain efficacy, and has not been tested in animal studies or randomized control trials for MERS-CoV [22, 31, 32]. Corticosteroids have not been successful in the treatment of respiratory distress or lung fibrosis in MERS-CoV [31, 32]. Meanwhile, studies in SARS-CoV and H1N1 patients suggest that corticosteroid use may in fact increase viral

replication in airways, and SARS patient and animal studies indicate that it contributes to immunosuppression [33–35]. Mycophenolate mofetil has been associated with fatal disease and high viral loads in a marmoset model of MERS-CoV infection [36]. IFN therapy, alone or in combination with ribavirin or lopinavir–ritonavir, has shown greater promise in *in vitro*, animal and human studies [37–40]. However, clinical studies on IFNs vary with respect to factors such as time of administration and type of patient [19, 22, 40]. Overall, there is a lack of randomized control trials (RCTs) designed to test the safety and efficacy of any potential therapies specific to MERS-CoV, and much of the information available for existing therapies is based on *in vitro* and/or animal studies [22, 40, 41]. A position paper on the evidence base for specific MERS-CoV therapies, published by Public Health England (PHE) and the World Health Organization–International Severe Acute Respiratory and Emerging Infection Consortium (ISARIC–WHO), suggested that benefit was likely to exceed risk for convalescent plasma, lopinavir–ritonavir, IFNs and monoclonal/polyclonal antibodies, while, by contrast, for ribavirin monotherapy and corticosteroids it was considered that the risks would outweigh the benefits [42]. For interferon/ribavirin combination therapy, nitazoxanide and chloroquine, the available data were considered to be inadequate for assessment [42].

In this review, we consider potentially effective MERS-CoV therapies, including IFNs, lopinavir–ritonavir and inhibitors of proteases, including TMPRSS2 and cathepsin L, as well as MERS-CoV-specific potential therapies, including convalescent plasma, monoclonal antibodies (mAbs), antiviral peptides and candidate vaccines. These therapies will be considered from a molecular perspective, in the context of the infection and replication mechanisms of MERS-CoV. The therapies are summarized in Table 1.

## MERS-COV INFECTION AND REPLICATION

### MERS-CoV lineage and structure

MERS-CoV is a betacoronavirus belonging to clade c (lineage 3) of the betacoronaviruses [43]. Its closest known coronavirus relatives are the prototypic clade c betacoronaviruses, *Tylonycteris* bat virus HKU4, *Pipistrellus* bat HKU5 virus and *Neoromicia zuluensis* bat PML/2011 (NeoCoV) virus [1, 43–47]. In common with other coronaviruses, the genome of MERS-CoV is a single, positive-stranded RNA of over 30 000 nucleotides. It encodes 10 predicted open reading frames (ORFs) and genes for 4 structural proteins, namely the spike (S), nucleocapsid (N), membrane (M) and envelope (E) proteins (Figs 1 and 2) [48–50]. ORF 1a and 1b encode virus replication-related proteins (pp1a, pp1ab), which are cleaved to give 16 non-structural proteins (NSPs) involved in synthesis of viral RNA and recombination (Fig. 2) [48–50]. These include NSP-14, which contains a 39-to-59 exoribonuclease (ExoN) domain that is important in viral proofreading and in determining the sensitivity of RNA viruses to mutagens. Thus small-molecule inhibitors

**Table 1.** Summary of potential MERS-CoV therapies

Target	Type	Therapy/vaccine	In vivo/in vitro	Safety/advantages	Side-effects/disadvantages	References
S1/DPP4 binding	Antibody (mouse): S1 RBD	Mersmab	<i>In vitro</i>			[76]
	Antibody (human): S1 RBD	m336, m337, m338	<i>In vitro/in vivo</i> (mouse, rabbit – m336)			[77–79]
	Antibody (human): S1 RBD	MERS-4, MERS-27	<i>In vitro</i>			[80, 81]
	Antibody (mouse- humanized): S1 RBD	4C2	<i>In vitro/in vivo</i> (mouse)	Prophylactic and therapeutic		[82]
	Antibody (mouse- humanized): S1 RBD	hMS-1	<i>In vitro/in vivo</i> (mouse)			[83]
	Antibody (human): S1 RBD	LCA60	<i>In vitro/in vivo</i> (mouse)	Targets both NTD and RBD; stable CHO cell line; prophylactic and therapeutic		[84]
	Antibody (human): S1 RBD	3B11-N	<i>In vitro/in vivo</i> (rhesus monkeys)			[85]
	Antibody (human- anti-DPP4)	2F9, 1F7, YS110	<i>In vitro</i>			[86]
	RBD-IgG fusion vaccine candidate	RBD s377-588- Fc IgG fusion	<i>In vitro/in vivo</i> (mouse)	Humoral response in mice; potential intranasal administration; improved by adjuvant MF59; divergent strains/escape mutants		[91–95]
	Full-length S protein proprietary nanoparticles		<i>In vitro/in vivo</i> (mouse)	Use of adjuvants improves humoral response	Stable expression of abundant full-length S protein difficult	[97]
Viral S2-host membrane fusion	MVA expressing full-length S protein (vaccine candidate)		<i>In vitro/in vivo</i> (mouse, camel)	T cell and humoral response; entering human clinical trials; potential for veterinary use – camels		[98, 99]
	ad5 or ad41 adenovirus expressing full-length S (vaccine candidate)		<i>In vitro/in vivo</i> (mouse)	T cell and neutralizing antibody responses		[99]
	Measles virus expressing full-length S (vaccine candidate)		<i>In vitro/in vivo</i> (mouse)	T cell and neutralizing antibody responses		[100]
	Plasmid vaccine	GLS-5300	<i>In vitro/in vivo</i> (mouse, camels and macaques)	T cell and neutralizing antibody responses; in phase I clinical trial		[102, 103]
	Anti-HR2 viral peptide	HR2P	<i>In vitro</i>			[87]
Immune evasion response	Anti-HR2 viral peptide	HR2P-M2	<i>In vitro/in vivo</i> (mouse)	Blocks 6HB bundle formation; enhances IFN- $\beta$ effect; potential intranasal treatments		[88–90]
	IFN- $\alpha$ 2b and ribavirin		<i>In vitro/in vivo</i> (macaque)	Combination therapy allows reduced amounts of each; non-human primate model; 10 different gene pathways		[108–110]
	IFN- $\beta$ 1b and lopinavir		<i>In vitro/in vivo</i> (marmoset)	Combination therapy allows reduced amounts of each		[111]
	IFN combination therapy (ribavirin and/or lopinavir)		Case studies (human)		Needs to be used prophylactically or early for any clinical benefit; insufficient evidence of clinical efficacy as yet	[37–40]
	IFN combination therapy (ribavirin)		Retrospective cohort studies (human)	Probable benefit of early use in less vulnerable patients; safety and efficacy established for other viral illnesses	Needs to be used prophylactically or early for any clinical benefit; insufficient evidence of clinical efficacy as yet	[27, 29, 105, 107, 112, 113]
S protein host proteases	TMPRSS2 inhibitor	Camostat	<i>In vivo</i> – mouse, SARS-CoV	Already in clinical use (chronic pancreatitis)		[59]
	TMPRSS2 inhibitor	Nafamostat	Split-protein-based cell–cell fusion assay	Already in clinical use (anti-coagulant)		[118]
	Cathepsin L inhibitor	Teicoplanin dalbavancin oritavancin telavancin	High-throughput screening	Already in clinical use (antibiotic Gram-positive bacterial infections)		[119]
	PL(pro) inhibitor	6-mercaptopurine (6MP) 6-thioguanine (6TG)	<i>In vitro</i>	Potential for more MERS-specific agents		[61]

Table 1. cont.

Target	Type	Therapy/vaccine	In vivo/in vitro	Safety/advantages	Side-effects/disadvantages	References
PI(pro) inhibitor		E2124–0890	In vitro		May lose potency in physiological reducing environments	123]
Mpro		Lopinavir	In vitro/In vivo (marmosets)	High activity in low micromolar range <i>in vitro</i> ; better outcomes, reduced mortality in marmosets	Clinical efficacy not fully established in humans	[36, 47, 125]





Fig. 1. Structure of MERS-CoV. Taken from: Belouzard *et al.* [128].

of ExoN activity could be candidates for MERS-CoV and other coronavirus therapies [51]. As with other coronaviruses, the MERS-CoV S protein is critical to host cell receptor binding and cell entry, and is considered to have been under strong positive selection pressure when the virus was transmitted to humans [52, 53]. Hence the S protein is a major target for potential anti-MERS-CoV therapies [53].

MERS-CoV Spike (S) protein

The S protein of MERS-CoV is composed of S1 and S2 subunits (Fig. 2) [53]. In common with other coronaviruses, entry into host cells depends on the S1 subunit, which contains a receptor-binding domain (RBD) comprising a core subdomain and a receptor-binding motif (RBM). The MERS-CoV RBM differs from that of SARS-CoV and dictates that MERS-CoV uses the DPP4 receptor, as opposed to the ACE-2 receptor [20, 21]. The infection process is shown in Fig. 2. DPP4, which is widely expressed in tissues, including the lung and kidneys, is critical in the species tropism of MERS-CoV infection; bat, human, camel, non-human primate and swine cells, for example, are permissive for MERS-CoV infection, whereas mouse, hamster and ferret are not [54, 55]. Host species restriction has been attributed to differences in five amino acids involved in DPP4-RBD binding, with glycosylation of the mouse DPP4 also identified as being important in the inhibition of MERS-CoV infection [54–56]. The human DPP4 receptor is therefore a potential target for MERS-CoV-specific therapeutics, in particular anti-DPP4 mAbs (Fig. 2, Table 1) [53–56]. Adenosine deaminase (ADA), which is a DPP4-binding protein, competes with MERS-CoV for DPP4 binding and hence is a natural MERS-CoV antagonist; this gives potential insights for the development of therapeutic antagonists [55].

MERS-CoV binds to the permissive host cell DPP4 via the RBD of the S1 domain, one of the major targets for potential MERS-CoV therapies [53]. Similarly to other coronaviruses, MERS-CoV then uses the S2 subunit for virus–host membrane fusion (Fig. 2). Fusion results in cleavage of the S



**Fig. 2.** Replication cycle and potential therapeutic targets of MERS-CoV. Adapted from Durai *et al.* [57].

## Host cell proteases and MERS-CoV infection

serine protease TMPRSS2, endosomal cathepsins such as cathepsin L, and furin protease [23, 53, 59–61]. *In vitro* studies suggest that uncleaved MERS pseudovirus can enter host cells by cathepsin L-dependent endocytosis, but that cleavage of virus during maturation by host proteases such as TMPRSS2 results in viral entry at neutral pH and the formation of massive syncytia [58]. Host cell proteases are therefore potential molecular therapeutic targets for MERS-CoV prophylaxis and/or treatment (Fig. 2, Table 1). The TMPRSS2 inhibitor camostat, for example, has been identified as a potential therapeutic agent for coronaviruses such as SARS-CoV and MERS-CoV [59].

further potential molecular targets for therapy (Table 1). The recently described MERS-CoV Mpro crystal structure resembles other coronavirus Mpro proteases [60]. The SARS-CoV PL(pro) inhibitors, 6-mercaptopurine (6MP) and 6-thioguanine (6TG), can inhibit MERS CoV protease activity *in vitro*, as can the immunosuppressant drug mycophenolic acid [61]. However, caution is required, as the results of studies on marmosets have associated use of mycophenolate mofetil with fatal disease and high viral loads [36].

### MERS-CoV proteins and immune system circumvention

Other MERS-CoV proteins involved in helping the virus to circumvent the immune system also present potential molecular targets (Fig. 2). For example, the accessory protein products of ORF 4a, 4b and 5 are interferon (IFN) antagonists [62, 63]. The ORF 4a protein both inhibits type I IFN production via cytoplasmic and nuclear mechanisms, and interferes with the IFN-mediated interferon-stimulated response element (ISRE) promoter element signalling pathways [62, 63]. This gives a molecular level rationale for the use of IFN as a therapeutic option in MERS-CoV treatment. MERS-CoV can also infect dendritic cells and macrophages [26, 64, 65]. Endosomal uptake of MERS-CoV by dendritic cells following binding via DPP4 prompts these cells to produce abundant amounts of type I and III IFNs [65]. This gives context to the IFN antagonism exhibited by MERS-CoV accessory proteins. Recently, MERS-CoV has also been shown to infect T cells, which are rich in DPP4, both *in vitro* and in marmoset spleen [66]. This results in T cell apoptosis and could contribute significantly to viral pathogenesis and further emphasizes the potential therapeutic utility of molecular targeting of DPP4 and/or the MERS-CoV S protein. Both convalescent plasma containing virus-specific antibodies and the use of specific mAbs provide options for targeting MERS-CoV infection at a molecular level.

### CONVALESCENT PLASMA

The use of convalescent plasma [or hyperimmune IV immunoglobulin (HVIG) from the plasma of convalescent donors] for infectious disease treatments has a long history, including in the treatment of respiratory diseases [67–70]. For influenza and SARS-CoV infection, early convalescent plasma treatment within 4–5 days of symptoms is associated with decreased viral load and reduction in mortality [67–70]. However, for SARS-CoV the quality of studies has been inconsistent and the results have been inconclusive, with a lack of adequate clinical trials [69, 70]. According to the PHE and ISARIC–WHO position paper, convalescent plasma (or high neutralizing antibody titre products) is indicated for the treatment of serious MERS-CoV infection [42]. One RCT on 35 critically ill patients with H1N1 infection identified a significant reduction in viral load and mortality in patients who received HVIG within 5 days of the onset of symptoms [68]. To date, no RCTs have been completed on the use of convalescent plasma/HVIG in MERS-

CoV patients. In the light of results from SARS and influenza patients, an ongoing clinical trial on the safety and efficacy of convalescent plasma treatment for critically ill MERS-CoV patients was initiated in May 2014 and is due to report in June 2017 [75; ClinicalTrials.gov identifier: NCT02190799]. This trial is being carried out in KSA. However, as is common for convalescent plasma therapies, the trial has been affected by logistical and technical issues, including the availability of sufficient donors [71, 72]. Issues can also arise in the collection of convalescent plasma that has sufficient levels of MERS-CoV antibodies, particularly outside the Middle East [22, 72]. While there are two case reports in which intravenous immunoglobulin (IVIG) was used in treatment of MERS-CoV, it is uncertain as to whether this contributed to patient recovery [73, 74]. Thus, while convalescent plasma is a promising potential therapy for MERS-CoV, the available clinical evidence is very limited and the results of the ongoing clinical trial will be vital in guiding any future use [71]. Focused development of neutralizing monoclonal antibodies targeted against specific MERS-CoV proteins has meanwhile yielded promising *in vitro* and/or *in vivo* results.

### MONOCLONAL ANTIBODIES: S1-DPP4 BINDING

A number of mouse and human neutralizing mAbs against the S1 region of MERS-CoV have been developed and tested *in vitro* and/or in animal models [52]. Targeting of S protein for therapeutic purposes was recently comprehensively reviewed by Du *et al.* [53] [52]. In particular, the S1 RBD is a popular target, as mAbs directed against this region have the most potent neutralizing capacity. However, in terms of vaccine development, neutralizing antibodies raised by immunization with full-length S or S1 protein expression vectors may produce a more effective immunogenicity through the targeting of multiple epitopes and the reduction of the possibility of escape mutations [75]. Nevertheless, many mouse and human mAbs targeting the S1 RBD have given promising results *in vitro* and in mouse models [19]. (Table 1). A mouse monoclonal antibody, Mersmab1, blocks MERS pseudovirus cell entry *in vitro* by binding to the RBD and preventing S1 binding to DPP4 [76]. Meanwhile, the human monoclonal antibodies m336, m337 and m338, which target overlapping epitopes in the RBD, all potentially neutralize pseudovirus and live MERS-CoV *in vitro* [77]. Significantly, intraperitoneal injection of m336 has also been shown to have both prophylactic and therapeutic protective effects against MERS-CoV infection in a well-established human DPP4 (hDPP4)-expressing transgenic mouse model, and in rabbits [78, 79]. Other anti-RBD human antibodies, MERS-4 and MERS-27, which recognize distinct RBD regions and block binding to DPP4, likewise have potent *in vitro* neutralizing activity against pseudovirus and live virus infection, and they also act synergistically [80, 81]. MERS-4 has anti-syncytia formation activity [80]. The crystal structure of MERS-27 bound to the DPP4 receptor revealed two critical RBD residues [81]. The crystal



structure of another anti-RBD antibody 4C2, which was raised in mice, has also been elucidated. This has allowed the identification of an epitope that partially overlaps the RBD receptor binding unit [82]. 4C2 was consequently humanized to give an antibody with prophylactic and therapeutic properties, shown by a reduction of MERS-CoV lung viral titres in an Ad5-hCD26 (hDPP4) transgenic mouse model [82]. Another humanized anti-RBD antibody, hMS-1, similarly has potent *in vivo* protective properties against fatal MERS-CoV infection in a transgenic hDPP4 mouse model [83]. The human antibody LCA60 targets both the N-terminal domain (NTD) and the RBD of the S1 region, and was isolated from B cells of a MERS-CoV-infected human donor before being used to rapidly establish a stable CHO cell line that can be used to reliably produce clinical grade antibody [84]. This is a promising candidate for clinical development, given the antibody's potent prophylactic and therapeutic activities against MERS-CoV infection in Ad5/hDPP4 transgenic mice and type I interferon receptor (IFNAR)-KO mice [84]. The human anti-RBD antibody 3B11-N is another promising candidate that prophylactically reduces lung pathology in rhesus monkeys infected with MERS-CoV [85].

Targeting the S1-DPP4 interaction from the host side through the development of anti-DPP4 (CD26) antibodies is another possible therapeutic option. The anti-CD26 antibodies 2F9, 1F7 and YS110 target the MERS-CoV entry into cells *in vitro* [86]. The 2F9 epitope maps close to the binding site of ADA, a natural DPP4 binding protein and MERS-CoV antagonist, while the 1F7 and YS110 epitopes lie outside this region [55, 86].

Thus, targeting of the S1-DPP4 interaction by use of mAbs is a promising strategy for the clinical development of molecular therapeutics against MERS-CoV. Another molecular approach involves targeting of the S2-mediated MERS-CoV-host cell fusion element of the MERS-CoV infection cycle by use of antiviral peptides.

## ANTIVIRAL PEPTIDES: HR2 REGION OF S PROTEIN

The role of HR1 and HR2 in viral fusion makes them potentially effective molecular therapeutic targets. This has been borne out by *in vitro* and *in vivo* results obtained using HR2 peptides (Table 1). HR1 peptides are ineffective antivirals as they aggregate in physiological solutions [87–89].

A peptide named HR2P, which spans residues 1251–1286 of HR2, effectively inhibits viral replication and S protein-mediated cell fusion *in vitro* [87]. A HR2P analogue named HR2P-M2 is an even more potent fusion blocker *in vitro*, and inhibits MERS CoV-expressing pseudovirus infection [90]. HR2P-M2 interacts with an HR1 peptide to effectively block 6HB bundle formation. *In vivo* HR2P-M2 intranasal administration to Ad5/hDPP4 transgenic mice protected them from MERS-CoV infection, as evidenced by the reduction of the lung viral titres by more than 1000-fold [90]. The

addition of IFN- $\beta$  along with HR2P-M2 enhanced the protective effect [90]. Thus, S2 HR2 peptides have potential as MERS-CoV intranasal antiviral treatments.

## VACCINE CANDIDATES

### S protein targeting

The S protein is also the focus of a number of candidate vaccines (Table 1) [75, 91–93]. A fusion product combining truncated RBD and the Fc portion of human IgG can bind human DPP4 and inhibit MERS-CoV infection in an *in vitro* cell culture model [91]. Importantly, this RBD-IgG fusion product can induce a humoral response in mice vaccinated by subcutaneous injection, hence blocking RBD-DPP4 binding and inhibiting MERS-CoV infection [91]. Further *in vivo* studies have indicated that intranasal administration to mice induces similar long-term IgG humoral responses to those achieved with subcutaneous injection, but superior cellular immune responses and local mucosal responses in lungs [92, 93]. This suggests that this type of construct is both potentially effective and readily deliverable by intranasal means. Use of an adjuvant, particularly MF59, significantly improves the humoral and T cell immunogenicity of the RBD s377–588-Fc IgG fusion construct in subcutaneously immunized mice [94]. The possibility of using the S1 RBD as a vaccine molecular target for a range of divergent MERS-CoV strains and escape mutants has also been explored recently [95]. The use of five recombinant RBDs with mutations observed in different MERS-CoV outbreaks or in camel strains induced potent neutralizing antibody responses against several MERS-CoV pseudoviruses [95].

However, while the RBD of the S1 subunit is a logical and promising target for MERS-CoV vaccine development, the epitope scope is relatively limited and full-length S protein may be a preferable option [75]. Technical difficulties in stably expressing abundant quantities of full-length S protein have presented a barrier. However, studies on delivery options, including the use of adjuvants and nanoparticles, may help in overcoming such issues. One study undertaken by Novavax (Gaithersburg, Maryland, USA) showed that the inoculation of mice by intramuscular injection with full-length S protein proprietary nanoparticles produced a relatively low neutralizing antibody response after 21 days [96]. However, the addition of the adjuvants Alum or Matrix M1 resulted in a robust and sustained anti-MERS-CoV neutralizing antibody response [96].

### Viral vectors

Other potential vaccination strategies include the use of live attenuated viruses, recombinant viruses or DNA plasmids expressing MERS-CoV genes. Various types of viral vectors are currently under development for use in potential MERS-CoV vaccines, including modified vaccinia virus Ankara (MVA), ad5- or ad41-type adenoviruses and measles viruses, all of which have good safety profiles (Table 1) [97–100]. Vaccination of mice by subcutaneous or

intraperitoneal injection with MVA expressing full-length S protein induces robust and sustained MERS-CoV-specific neutralizing antibody and cytotoxic T lymphocyte responses, including in mice expressing human DPP4 [98]. These viruses are expected to enter clinical trials as a proposed prophylactic MERS-CoV vaccine. Likewise, intramuscular injection of ad5 or ad41 expressing full-length S protein induces both antigen-specific T cell and neutralizing antibody responses in mice [99]. Finally, intraperitoneal injection of measles virus expressing either membrane-anchored, full-length S protein or soluble S protein lacking the TM domain induces robust MERS-CoV antigen-specific neutralizing antibody and cytotoxic T lymphocyte in interferon- $\alpha/\beta$  receptor (IFNAR)-deficient mice [100]. Recently, an MVA-based vaccine expressing S protein has been shown to induce mucosal immunity in MERS-CoV-infected dromedary camels, with a reduction in excreted virus and viral transcripts [101]. This has potential for veterinary use and the reduction of cross-species infection of humans by camels [101].

### DNA plasmids

GLS-5300 is a DNA-plasmid vaccine that encodes MERS-CoV S protein (Table 1) [102, 103]. It was co-developed by Inovio, GeneOne Life Science, Inc. and the Walter Reed Army Institute of Research, and has become the first potential MERS-CoV vaccine to enter human testing [102, 103]. A phase I clinical trial in healthy volunteers commenced in 2016 for the evaluation of its safety and ability to generate antibody and cellular immune responses over a 1-year period, using one of three dosages in a three-injection regimen [102]. The vaccine has already undergone pre-clinical trials in mice, camels and macaques [103]. It induced robust and antigen-specific cytotoxic T lymphocyte and neutralizing antibody responses, which effectively protected animals against viral infection [103].

GLS-5300 and other potential vaccine candidates provide an opportunity to develop a prophylactic MERS-CoV vaccine. However, the barriers to development of a prophylactic vaccine include the current relatively low MERS-CoV incidence in humans, as well as sourcing suitable small animal models [75, 97, 104]. These factors complicate the definition of a target population for mass prophylactic vaccination and pre-clinical demonstration of vaccine efficacy [104]. In this context, the monoclonal antibodies described above may be invaluable resources in an outbreak situation [76–85].

## INTERFERONS: MONOTHERAPY AND COMBINATION THERAPY

### *In vitro* and animal studies

While MERS-CoV-specific therapies are offering promising pre-clinical results, and GLS-5300 has entered clinical trials, there is as yet no specific evidence-based therapy or vaccine clinically available for MERS-CoV. As described in the MERS-CoV infection and replication section, MERS-CoV

accessory protein products are IFN antagonists [62, 63]. Attenuation of the IFN response is an important MERS-CoV immune response circumvention mechanism [105]. The ORF4a in particular inhibits IFN- $\beta$  production via the inhibition of interferon regulatory transcription factor (IRF)-3 and nuclear factor (NF)- $\kappa$ B actions, and thus IRF-3-activating small molecules, for example, may be potential therapeutic agents for restoring IFN responses [62, 63]. Toll-like receptor-3 (TLR-3) is also involved in the immune response of mice to SARS-CoV and MERS-CoV, recognizing viral molecular patterns and initiating the innate response that leads to IFN production (Fig. 2) [106]. Thus, TLR-3 agonists are another possible candidate for MERS-CoV-specific anti-viral agents [106].

Therapeutically, IFN itself is particularly useful prophylactically or during the early days of viral exposure, including for coronaviruses [105, 107]. *In vitro* and animal studies have confirmed the potential efficacy of IFNs in MERS-CoV therapy, in particular in combination with other therapeutic agents such as ribavirin and/or lopinavir. *In vitro*, MERS-CoV was substantially more susceptible to IFN- $\alpha$  than SARS-CoV [107]. While MERS-CoV in Vero or LLC-MK2 cells was sensitive to both IFN- $\alpha$ 2b and ribavirin separately, relatively high concentrations were required to reduce viral replication [108]. However, combination therapy allowed the concentrations of each to be substantially reduced [108]. Combination therapy of IFN- $\alpha$ 2b and ribavirin in macaques administered 8 hours after MERS-CoV infection reduced systemic and local viral effects, and reduced viral genome copy number and gene expression levels [109]. Bioinformatics data from microarray analysis recently showed that IFN- $\alpha$ 2b and ribavirin treatment impacts on MERS-CoV gene expression in 10 different pathways, including genes involved in recognition of pathogens, immune responses and release of cytokines [110]. Both IFN- $\beta$ 1b and lopinavir treatment, alone or in combination, also protected marmosets from the adverse clinical, radiological and pathological effects of MERS-CoV infection [111].

### Clinical studies

Clinically, the use of IFN monotherapy, or IFN therapy in combination with ribavirin and/or lopinavir/ritonavir, has shown some promise (Table 1) [37–40]. However, the interpretation of clinical studies has been complicated by variability in factors such as the stage of infection at which therapy was administered. The available data are limited to case studies and retrospective cohort studies [22, 40]. In one case study on a patient who died in a Greek hospital, pegylated IFN along with ribavirin and lopinavir was administered as part of the treatment regime, but not until the thirteenth day of the illness [39]. By contrast, in another preliminary study on two patients, the first patient was treated with IFN- $\alpha$ 2b and ribavirin within a day of admission prior to MERS-CoV diagnosis, but he was also being treated with antibiotics, steroids and non-invasive ventilation [37]. Patient 2, the wife of patient 1, was treated prophylactically after developing a low-grade fever and poorly



defined lung infiltrates, but a diagnosis of MERS-CoV was not formally made [37]. Thus, while patient 1 survived and patient 2 had only a mild course of illness, it is difficult to draw any firm conclusions regarding the efficacy of the treatment. In another case study on a patient in Korea, administration of pegylated IFN- $\alpha$ 2a along with ribavirin and lopinavir 4 days after hospital admission was deemed to have been effective in viral clearance and patient survival [38]. These case studies do not overall provide firm evidence for the efficacy or otherwise of IFN combination therapy for MERS-CoV.

In one case involving a series of five patients who were critically ill with MERS-CoV infection and on mechanical ventilation and corticosteroids, IFN- $\alpha$ 2b and ribavirin was administered on average 19 days after admission [27]. All five patients died, but they may not have benefited, as they were treated late in their illness and were already critically ill [27]. The benefit of earlier treatment in less vulnerable patients was suggested in another series of six patients in which three who received IFN- $\alpha$ 2b and ribavirin early in the illness survived, while three other patients who were older and had comorbid conditions received the combination treatment later and all died [112]. However, in another study in which 20 mechanically ventilated patients with severe MERS-CoV infection who received pegylated IFN- $\alpha$ 2a and ribavirin early in treatment were compared to 24 patients who did not receive the combination therapy, the 14-day survival rate was significantly higher in the treatment group, but the 28-day survival rate was equivalently low in the two groups [113]. In another retrospective analysis of results from a series of 32 patients who received either IFN- $\alpha$ 2a or IFN- $\beta$ 1a in combination with ribavirin, no significant difference in outcome between the two types of IFN was shown, and there was no survival benefit due to use of either IFN [29]. However, most of the patients in this study were aged more than 50 years and some had comorbid conditions, including end-stage renal disease [29]. Thus the retrospective studies that have been carried out are heterogeneous in terms of type of patient, stage of disease and type of IFN used, including whether or not it was pegylated or short-acting. There is an urgent need for well-controlled clinical trials for IFN combination therapy in MERS-CoV, preferably early in the illness, as IFNs are routinely available agents whose safety and efficacy is established for other viral illnesses and whose use has a sound molecular basis for MERS-CoV treatment.

## PROTEASE INHIBITORS

### S protein proteases

Another type of therapy with a logical molecular basis for MERS-CoV treatment is the targeting of proteases, both host and viral (Table 1; Fig. 2) [19, 23, 53, 59, 60, 114–117]. Camostat, an inhibitor of TMPRSS2, is a potential therapeutic agent for coronaviruses such as SARS-CoV and MERS-CoV [59]. In a pathogenic mouse model of SARS-CoV infection, viral spread and pathogenesis was effectively

blocked by camostat, and it is likely that it would have a similar impact on MERS-CoV [59]. As camostat is already in clinical use for the treatment of chronic pancreatitis, it represents a potentially safe and effective therapeutic option. Recently, another TMPRSS2 inhibitor, nafamostat, was identified in a split protein-based cell–cell fusion assay as a potent inhibitor of MERS-CoV S protein-mediated host–viral membrane fusion *in vitro* [118]. Nafamostat is already clinically approved for use by the US Food and Drug Administration (FDA) and is used as an anticoagulant [118]. The cathepsin L inhibitor teicoplanin, a glycopeptide antibiotic, was recently shown, via high throughput screening of FDA-approved drugs, to block entry of MERS-CoV, SARS-CoV and Ebola pseudoviruses into the cytoplasm [119]. Teicoplanin is currently used clinically as an antibiotic in both prophylaxis and the treatment of serious Gram-positive bacterial infections. It also has derivatives, including dalbavancin, oritavancin and telavancin, all of which also block viral entry.

### Viral proteases

#### PL(pro) inhibitors

The viral proteases, Mpro (3CLpro) and PL(pro), also represent potential molecular therapeutic targets [57, 60]. As well as its role in viral maturation, the MERS-CoV PL(pro) causes deubiquitination of IFN regulatory factor 3 (IRF-3), and hence suppression of IFN  $\beta$  production, which contributes to viral suppression of the innate immune response (Fig. 2) [120, 121]. The X-ray 3D crystal structure of MERS-CoV PL(pro) is similar to that of SARS-CoV, and includes ubiquitin-like and catalytic core domains [120]. Thus the SARS-CoV PL(pro) inhibitors, 6-mercaptopurine (6MP) and 6-thioguanine (6TG), can inhibit MERS CoV protease activity *in vitro* [61]. However, the MERS-CoV PL<sup>pro</sup> crystal structure also has unique aspects, including the oxyanion hole, and S3 and S5 subsites, which may be viable molecular targets for antivirals specifically designed against MERS-CoV [120]. A commercial compound termed compound 4 (commercial code F2124–0890, Life Chemicals) has been identified as an inhibitor of MERS-CoV and SARS-CoV PLpro activity [122, 123]. The critical binding interactions and mode of inhibition differ between the two viral proteases, with the compound acting as a competitive inhibitor against MERS-CoV PL(pro), but an allosteric inhibitor of SARS-CoV PL[pro] [122]. However, F2124–0890 may lose potency in physiological reducing environments [123].

#### Mpro inhibitors: lopinavir/ritonavir

Lopinavir is a protease inhibitor with activity against the SARS-CoV main protease M<sup>pro</sup> [124]. In a screen of a library of 348 FDA-approved drugs to identify anti-MERS-CoV activity in cell culture, lopinavir emerged as one of four compounds that inhibited viral activity in a low micromolar range [125]. However, the clinical efficacy of lopinavir in MERS-CoV treatment has not yet been fully established. As mentioned above, it has usually been used clinically in combination with IFN and data are only available from case studies and series. However, notably,

lopinavir–ritonavir treatment resulted in better clinical, radiological and pathological outcomes and reduced mortality in marmosets infected with MERS-CoV [36]. Lopinavir has also been identified in a position paper from PHE and ISARIC–WHO as a potential MERS-CoV therapy whose benefits are likely to exceed its risks [47].

## CONCLUSIONS

Thus far, MERS-CoV has not been considered to have pandemic potential. Most cases have occurred in the Middle East, particularly in KSA. Outbreaks have been primarily linked to healthcare institutions, and shortcomings in infection control and prevention procedures [6–14]. However, potential viral mutations could facilitate expanded viral host range and enhance cross-species and human–human transmission [20, 58, 114]. The outbreak in Korea resulted in MERS-CoV emergence in second- and third-generation contacts, highlighting the potential for mutational changes that could increase the likelihood of human–human transmission [14, 18]. MERS-CoV also exacts a high mortality rate, mainly due to the development of ARDS [15–17]. These factors emphasize the importance of developing targeted therapies and/or vaccines. The most promising advances in the development of specific molecular MERS-CoV therapies relate to targeting of the viral S protein by means of anti-S1 monoclonal antibodies, HR-targeted antiviral peptides and viruses or plasmids bearing S protein as potential vaccine candidates [52, 55, 58, 88–105]. The use of IFNs, usually in combination with other therapies such as ribavirin or lopinavir, also has a logical molecular basis given that IFN antagonism is an important mechanism by which the virus circumvents the innate immune system [62, 63, 105, 107]. Targeting of host and viral proteases is also a sound molecular approach, as host proteases are important in viral–host membrane fusion, while viral proteases are key to viral maturation and are also involved in targeting IFNs [23, 53, 59–61, 114–117].

The therapies currently used for MERS-CoV have mainly been extrapolated from those used for SARS-CoV treatment, regardless of the important differences in receptor usage and cellular tropism between the viruses [20–26]. None of these therapies have been subject to well-controlled trials, and in some cases the risks are likely to outweigh any poorly defined benefits [19, 22, 27–42]. In general, the clinical research response to MERS-CoV may have been too slow [126]. Thus, while there are many promising lines of research in terms of specific molecular targeting of MERS-CoV, no potential therapies have yet been subject to well-designed clinical trials, and none have been approved for clinical use, apart from the GLS-5300 DNA-plasmid vaccine [102, 105]. Continuing outbreaks of MERS-CoV, with possible increases in human–human transmission, are likely to galvanize the research community to push ahead with the design and performance of clinical trials for some of the available monoclonal antibodies and/or antiviral peptides for use in outbreak situations.

There are various challenges inherent in the development of specific MERS-CoV therapies. These include the difficulty of identifying a target population for potential prophylactic vaccines, limited small animal model availability and dependence on transgenic mouse models, and the current relatively low incidence of infection, which complicates the performance of adequate clinical trials [75, 96, 97, 104]. For example, one currently ongoing trial on convalescent plasma therapy has been affected by logistical and technical issues, including insufficient available donors and difficulty in collecting convalescent plasma containing sufficient MERS-CoV antibody levels [71, 72]. Thus, while numerous monoclonal antibodies have been raised with anti-MERS activity, in particular against the S protein [76–85], and promising antiviral HR2 peptides have been synthesized [87, 90], the available data are thus far limited to *in vitro* and animal studies.

Despite these issues, there is cause for optimism, given the many candidate antibody and peptide therapies. There is also some promising *in vitro* and animal model evidence suggesting that use of IFNs, which are well-established therapies in other viral illnesses, may be of benefit if used sufficiently early in MERS-CoV treatment, or as a prophylactic, especially in combination with other therapies, including ribavirin or lopinavir–ritonavir [108–111]. Likewise, other drugs that are currently in clinical use for other conditions have been shown to be potentially useful for MERS-CoV treatment, including camostat and nafamostat; teicoplanin and its derivatives dalbavancin, oritavancin and telavancin; and the SARS-CoV PL(pro) inhibitors, 6-mercaptopurine (6MP) and 6-thioguanine (6TG) [59, 61, 118–121]. These drugs have already been shown to be safe and well-tolerated by humans. Repurposing of existing drugs may therefore prove to be the most viable option in MERS-CoV therapy. For example, 1 screen of 290 approved drugs uncovered 27 candidates with *in vitro* activity against both MERS-CoV and SARS-CoV, including oestrogen receptor inhibitors and dopamine receptor inhibitors [127]. Thus, there are many options available on a molecular level for the development of new MERS-CoV-specific therapies, as well as the adoption of drugs that are currently in use for other purposes, which should assist in more effective and reliable prevention and treatment of this virus.

### Funding information

The authors received no specific grant from any funding agency.

### Conflicts of interest

The authors declare that there are no conflicts of interest.

### Ethical statement

This is a review article and no experimental work with humans was performed.

### References

1. Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA *et al.* Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* 2012;367:1814–1820.

2. WHO. Middle East Respiratory Syndrome Coronavirus (MERS-CoV) [Internet]. Geneva: World Health Organization (WHO); 2017. [www.who.int/emergencies/mers-cov/en/](http://www.who.int/emergencies/mers-cov/en/) [Cited 11 July 2017].
3. Drexler JF, Corman VM, Drosten C. Ecology, evolution and classification of bat coronaviruses in the aftermath of SARS. *Antiviral Res* 2014;101:45–56.
4. Haagmans BL, Al Dhahiry SH, Reusken CB, Raj VS, Galiano M et al. Middle East respiratory syndrome coronavirus in dromedary camels: an outbreak investigation. *Lancet Infect Dis* 2014;14:140–145.
5. Reusken CB, Haagmans BL, Müller MA, Gutierrez C, Godeke GJ et al. Middle East respiratory syndrome coronavirus neutralising serum antibodies in dromedary camels: a comparative serological study. *Lancet Infect Dis* 2013;13:859–866.
6. Hijawi B, Abdallat M, Sayaydeh A, Alqasrawi S, Haddadin A et al. Novel coronavirus infections in Jordan, April 2012: epidemiological findings from a retrospective investigation/Infections par le nouveau coronavirus en Jordanie, avril 2012: résultats épidémiologiques d'une étude rétrospective. *East Mediterr Health J* 2013;19:S12–S18.
7. Assiri A, McGeer A, Perl TM, Price CS, Al Rabeeah AA et al. Hospital outbreak of Middle East respiratory syndrome coronavirus. *N Engl J Med* 2013;369:407–416.
8. Oboho IK, Tomczyk SM, Al-Asmari AM, Banjar AA, Al-Mugti H et al. 2014 MERS-CoV outbreak in Jeddah – a link to health care facilities. *N Engl J Med* 2015;372:846–854.
9. Fagbo SF, Skakni L, Chu DK, Garbati MA, Joseph M et al. Molecular epidemiology of hospital outbreak of Middle East respiratory syndrome, Riyadh, Saudi Arabia, 2014. *Emerg Infect Dis* 2015;21:1981–1988.
10. Drosten C, Muth D, Corman VM, Hussain R, Al Masri M et al. An observational, laboratory-based study of outbreaks of middle East respiratory syndrome coronavirus in Jeddah and Riyadh, Kingdom of Saudi Arabia, 2014. *Clin Infect Dis* 2015;60:369–377.
11. Almekhlafi GA, Albarrak MM, Mandourah Y, Hassan S, Alwan A et al. Presentation and outcome of Middle East respiratory syndrome in Saudi intensive care unit patients. *Crit Care* 2016;20:123.
12. Balkhy HH, Alenazi TH, Alshamrani MM, Baffoe-Bonnie H, Al-Abdely HM et al. Notes from the field: nosocomial outbreak of Middle East respiratory syndrome in a large Tertiary Care Hospital - Riyadh, Saudi Arabia, 2015. *MMWR Morb Mortal Wkly Rep* 2016;65:163–164.
13. Balkhy HH, Perl TM, Arabi YM. Preventing healthcare-associated transmission of the Middle East respiratory syndrome (MERS): our Achilles heel. *J Infect Public Health* 2016;9:208–212.
14. World Health Organization (WHO). 2017. Middle East respiratory syndrome coronavirus (MERS-CoV). MERS-CoV in Republic of Korea at a glance. [www.wpro.who.int/outbreaks\\_emergencies/wpro\\_coronavirus/en/](http://www.wpro.who.int/outbreaks_emergencies/wpro_coronavirus/en/). [Accessed 10 January 2017].
15. World Health Organization (WHO). 2015. Middle East respiratory syndrome coronavirus (MERS-CoV): Summary of Current Situation, Literature Update and Risk Assessment. [www.who.int/csr/disease/coronavirus\\_infections/risk-assessment-7july2015/en/](http://www.who.int/csr/disease/coronavirus_infections/risk-assessment-7july2015/en/) [Accessed 10 January 2017].
16. Assiri A, Al-Tawfiq JA, Al-Rabeeah AA, Al-Rabiah FA, Al-Hajjar S et al. Epidemiological, demographic, and clinical characteristics of 47 cases of Middle East respiratory syndrome coronavirus disease from Saudi Arabia: a descriptive study. *Lancet Infect Dis* 2013;13:752–761.
17. Min CK, Cheon S, Ha NY, Sohn KM, Kim Y et al. Comparative and kinetic analysis of viral shedding and immunological responses in MERS patients representing a broad spectrum of disease severity. *Sci Rep* 2016;6:25359.
18. Wang Y, Liu D, Shi W, Lu R, Wang W et al. Origin and possible genetic recombination of the Middle East respiratory syndrome coronavirus from the first imported case in China: phylogenetics and coalescence analysis. *MBio* 2015;6:e01280-15.
19. Zumla A, Chan JF, Azhar EI, Hui DS, Yuen KY. Coronaviruses – drug discovery and therapeutic options. *Nat Rev Drug Discov* 2016;15:327–347.
20. Raj VS, Mou H, Smits SL, Dekkers DH, Müller MA et al. Dipeptidyl peptidase 4 is a functional receptor for the emerging human coronavirus-EMC. *Nature* 2013;495:251–254.
21. Wang N, Shi X, Jiang L, Zhang S, Wang D et al. Structure of MERS-CoV spike receptor-binding domain complexed with human receptor DPP4. *Cell Res* 2013;23:986–993.
22. Mo Y, Fisher D. A review of treatment modalities for Middle East respiratory syndrome. *J Antimicrob Chemother* 2016;71:3340–3350.
23. Shirato K, Kawase M, Matsuyama S. Middle East respiratory syndrome coronavirus infection mediated by the transmembrane serine protease TMPRSS2. *J Virol* 2013;87:12552–12561.
24. Müller MA, Raj VS, Muth D, Meyer B, Kallies S et al. Human coronavirus EMC does not require the SARS-coronavirus receptor and maintains broad replicative capability in mammalian cell lines. *mBio* 2012;3:e00515-12.
25. Chan RW, Hemida MG, Kayali G, Chu DK, Poon LL et al. Tropism and replication of Middle East respiratory syndrome coronavirus from dromedary camels in the human respiratory tract: an *in-vitro* and *ex-vivo* study. *Lancet Respir Med* 2014;2:813–822.
26. Zhou J, Chu H, Li C, Wong BH, Cheng ZS et al. Active replication of Middle East respiratory syndrome coronavirus and aberrant induction of inflammatory cytokines and chemokines in human macrophages: implications for pathogenesis. *J Infect Dis* 2014;209:1331–1342.
27. Al-Tawfiq JA, Momattin H, Dib J, Memish ZA. Ribavirin and interferon therapy in patients infected with the Middle East respiratory syndrome coronavirus: an observational study. *Int J Infect Dis* 2014;20:42–46.
28. Momattin H, Mohammed K, Zumla A, Memish ZA, Al-Tawfiq JA et al. Therapeutic options for Middle East respiratory syndrome coronavirus (MERS-CoV) – possible lessons from a systematic review of SARS-CoV therapy. *Int J Infect Dis* 2013;17:e792–e798.
29. Shalhoub S, Farahat F, Al-Jiffri A, Simhairi R, Shamma O et al. IFN- $\alpha$ 2a or IFN- $\beta$ 1a in combination with ribavirin to treat Middle East respiratory syndrome coronavirus pneumonia: a retrospective study. *J Antimicrob Chemother* 2015;70:2129–2132.
30. Al Ghamdi M, Alghamdi KM, Ghandoor A, Alzahrani A, Salah F et al. Treatment outcomes for patients with Middle Eastern Respiratory Syndrome Coronavirus (MERS CoV) infection at a coronavirus referral center in the Kingdom of Saudi Arabia. *BMC Infect Dis* 2016;16:174.
31. Who Mers-Cov Research Group. State of Knowledge and data gaps of Middle East respiratory syndrome coronavirus (MERS-CoV) in humans. *PLoS Curr* 2013;5.
32. Jones BM, Ma ES, Peiris JS, Wong PC, Ho JC et al. Prolonged disturbances of *in vitro* cytokine production in patients with severe acute respiratory syndrome (SARS) treated with ribavirin and steroids. *Clin Exp Immunol* 2004;135:467–473.
33. Zhang X, Alekseev K, Jung K, Vlasova A, Hadya N et al. Cytokine responses in porcine respiratory coronavirus-infected pigs treated with corticosteroids as a model for severe acute respiratory syndrome. *J Virol* 2008;82:4420–4428.
34. Lee N, Allen Chan KC, Hui DS, Ng EK, Wu A et al. Effects of early corticosteroid treatment on plasma SARS-associated Coronavirus RNA concentrations in adult patients. *J Clin Virol* 2004;31:304–309.
35. Kim SH, Hong SB, Yun SC, Choi WI, Ahn JJ et al. Corticosteroid treatment in critically ill patients with pandemic influenza A/H1N1 2009 infection: analytic strategy using propensity scores. *Am J Respir Crit Care Med* 2011;183:1207–1214.
36. Chan JF, Yao Y, Yeung ML, Deng W, Bao L et al. Treatment with lopinavir/ritonavir or interferon- $\beta$ 1b improves outcome of MERS-CoV infection in a nonhuman primate model of common marmoset. *J Infect Dis* 2015;212:1904–1913.

37. Khalid M, Al Rabiah F, Khan B, Al Mobeireek A, Butt TS *et al*. Ribavirin and interferon- $\alpha$ 2b as primary and preventive treatment for Middle East respiratory syndrome coronavirus: a preliminary report of two cases. *Antivir Ther* 2015;20:87–91.
38. Kim UJ, Won EJ, Kee SJ, Jung SI, Jang HC *et al*. Combination therapy with lopinavir/ritonavir, ribavirin and interferon- $\alpha$  for Middle East respiratory syndrome. *Antivir Ther* 2016;21:455–459.
39. Spanakis N, Tsiodras S, Haagmans BL, Raj VS, Pontikis K *et al*. Virological and serological analysis of a recent Middle East respiratory syndrome coronavirus infection case on a triple combination antiviral regimen. *Int J Antimicrob Agents* 2014;44:528–532.
40. Strayer DR, Dickey R, Carter WA. Sensitivity of SARS/MERS CoV to interferons and other drugs based on achievable serum concentrations in humans. *Infect Disord Drug Targets* 2014;14:37–43.
41. Al-Tawfiq JA, Memish ZA. Update on therapeutic options for Middle East Respiratory Syndrome Coronavirus (MERS-CoV). *Expert Rev Anti Infect Ther* 2017;15:269–275.
42. Public Health England/ISARIC. 2015. Treatment of MERS-CoV; information for clinicians. Clinical decision-making support for treatment of MERS-CoV patients. [www.google.ie/?gws\\_rd=ssl#q=public+health+england+treatment+mers-cov](http://www.google.ie/?gws_rd=ssl#q=public+health+england+treatment+mers-cov) [Accessed 21 October 2016].
43. de Groot RJ, Baker SC, Baric R, Enjuanes L, Gorbalenya AE *et al*. Family Coronaviridae. In: King AMQ, Adams MJ, Carstens EB and Lefkowitz EJ (editors). *Virus Taxonomy: Classification and Nomenclature of Viruses. Ninth Report of the International Committee on Taxonomy of Viruses*. London, United Kingdom: Academic Press; 2012. pp. 806–820.
44. Reusken CB, Raj VS, Koopmans MP, Haagmans BL. Cross host transmission in the emergence of MERS coronavirus. *Curr Opin Virol* 2016;16:55–62.
45. Corman VM, Ithete NL, Richards LR, Schoeman MC, Preiser W *et al*. Rooting the phylogenetic tree of middle East respiratory syndrome coronavirus by characterization of a conspecific virus from an African bat. *J Virol* 2014;88:11297–11303.
46. Ithete NL, Stoffberg S, Corman VM, Cottontail VM, Richards LR *et al*. Close relative of human Middle East respiratory syndrome coronavirus in bat, South Africa. *Emerg Infect Dis* 2013;19:1697–1699.
47. Rambaut A. 2014. MERS-Coronavirus molecular epidemiology and genetic analysis – Origin and evolution. [http://epidemic.bio.ed.ac.uk/coronavirus\\_analysis](http://epidemic.bio.ed.ac.uk/coronavirus_analysis) [Accessed 14 June 2016].
48. Mackay IM, Arden KE. MERS coronavirus: diagnostics, epidemiology and transmission. *Virol J* 2015;12:222.
49. van Boheemen S, de Graaf M, Lauber C, Bestebroer TM, Raj VS *et al*. Genomic characterization of a newly discovered coronavirus associated with acute respiratory distress syndrome in humans. *mBio* 2012;3:e00473–12.
50. Scobey T, Yount BL, Sims AC, Donaldson EF, Agnihothram SS *et al*. Reverse genetics with a full-length infectious cDNA of the Middle East respiratory syndrome coronavirus. *Proc Natl Acad Sci USA* 2013;110:16157–16162.
51. Smith EC, Blanc H, Surdel MC, Vignuzzi M, Denison MR. Coronaviruses lacking exoribonuclease activity are susceptible to lethal mutagenesis: evidence for proofreading and potential therapeutics. *PLoS Pathog* 2013;9:e1003565.
52. Zhang Z, Shen L, Gu X. Evolutionary dynamics of MERS-CoV: potential recombination, positive selection and transmission. *Sci Rep* 2016;6:25049.
53. Du L, Yang Y, Zhou Y, Lu L, Li F *et al*. MERS-CoV spike protein: a key target for antivirals. *Expert Opin Ther Targets* 2017;21:131–143.
54. van Doremalen N, Miazgowiec KL, Milne-Price S, Bushmaker T, Robertson S *et al*. Host species restriction of Middle East respiratory syndrome coronavirus through its receptor, dipeptidyl peptidase 4. *J Virol* 2014;88:9220–9232.
55. Raj VS, Smits SL, Provacia LB, van den Brand JM, Wiersma L *et al*. Adenosine deaminase acts as a natural antagonist for dipeptidyl peptidase 4-mediated entry of the Middle East respiratory syndrome coronavirus. *J Virol* 2014;88:1834–1838.
56. Peck KM, Cockrell AS, Yount BL, Scobey T, Baric RS *et al*. Glycosylation of mouse DPP4 plays a role in inhibiting Middle East respiratory syndrome coronavirus infection. *J Virol* 2015;89:4696–4699.
57. Durai P, Batool M, Shah M, Choi S. Middle East respiratory syndrome coronavirus: transmission, virology and therapeutic targeting to aid in outbreak control. *Exp Mol Med* 2015;47:e181.
58. Graham RL, Baric RS. Recombination, reservoirs, and the modular spike: mechanisms of coronavirus cross-species transmission. *J Virol* 2010;84:3134–3146.
59. Zhou Y, Vedantham P, Lu K, Agudelo J, Carrion R *et al*. Protease inhibitors targeting coronavirus and filovirus entry. *Antiviral Res* 2015;116:76–84.
60. Ho BL, Cheng SC, Shi L, Wang TY, Ho KI *et al*. Critical assessment of the important residues involved in the dimerization and catalysis of MERS Coronavirus Main protease. *PLoS One* 2015;10:e0144865.
61. Cheng KW, Cheng SC, Chen WY, Lin MH, Chuang SJ *et al*. Thio-purine analogs and mycophenolic acid synergistically inhibit the papain-like protease of Middle East respiratory syndrome coronavirus. *Antiviral Res* 2015;115:9–16.
62. Yang Y, Zhang L, Geng H, Deng Y, Huang B *et al*. The structural and accessory proteins M, ORF 4a, ORF 4b, and ORF 5 of Middle East respiratory syndrome coronavirus (MERS-CoV) are potent interferon antagonists. *Protein Cell* 2013;4:951–961.
63. Yang Y, Ye F, Zhu N, Wang W, Deng Y *et al*. Middle East respiratory syndrome coronavirus ORF4b protein inhibits type I interferon production through both cytoplasmic and nuclear targets. *Sci Rep* 2015;5:17554.
64. Chu H, Zhou J, Wong BH, Li C, Cheng ZS *et al*. Productive replication of Middle East respiratory syndrome coronavirus in monocyte-derived dendritic cells modulates innate immune response. *Virology* 2014;454–455:197–205.
65. Scheuplein VA, Seifried J, Malczyk AH, Miller L, Höcker L *et al*. High secretion of interferons by human plasmacytoid dendritic cells upon recognition of Middle East respiratory syndrome coronavirus. *J Virol* 2015;89:3859–3869.
66. Chu H, Zhou J, Wong BH, Li C, Chan JF *et al*. Middle East respiratory syndrome Coronavirus efficiently infects human primary T lymphocytes and activates the extrinsic and intrinsic apoptosis pathways. *J Infect Dis* 2016;213:904–914.
67. Hui DS, Lee N. Adjunctive therapies and immunomodulating agents for severe influenza. *Influenza Other Respir Viruses* 2013;7:52–59.
68. Hung IFN, To KKW, Lee CK, Lee KL, Yan WW *et al*. Hyperimmune IV immunoglobulin treatment: a multicenter double-blind randomized controlled trial for patients with severe 2009 influenza A(H1N1) infection. *Chest* 2013;144:464–473.
69. Mair-Jenkins J, Saavedra-Campos M, Baillie JK, Cleary P, Khaw FM *et al*. The effectiveness of convalescent plasma and hyperimmune immunoglobulin for the treatment of severe acute respiratory infections of viral etiology: a systematic review and exploratory meta-analysis. *J Infect Dis* 2015;211:80–90.
70. Stockman LJ, Bellamy R, Garner P. SARS: systematic review of treatment effects. *PLoS Med* 2006;3:e343.
71. Arabi Y, Balkhy H, Hajeer AH, Bouchama A, Hayden FG *et al*. Feasibility, safety, clinical, and laboratory effects of convalescent plasma therapy for patients with Middle East respiratory syndrome coronavirus infection: a study protocol. *Springerplus* 2015;4:709.
72. Modjarrad K. Treatment strategies for Middle East respiratory syndrome coronavirus. *J Virus Erad* 2016;2:1–4.

73. Arabi YM, Arifi AA, Balkhy HH, Najm H, Aldawood AS *et al*. Clinical course and outcomes of critically ill patients with Middle East respiratory syndrome coronavirus infection. *Ann Intern Med* 2014;160:389–397.
74. Kapoor M, Pringle K, Kumar A, Dearth S, Liu L *et al*. Clinical and laboratory findings of the first imported case of Middle East respiratory syndrome coronavirus to the United States. *Clin Infect Dis* 2014;59:1511–1518.
75. Wang L, Shi W, Joyce MG, Modjarrad K, Zhang Y *et al*. Evaluation of candidate vaccine approaches for MERS-CoV. *Nat Commun* 2015;6:7712.
76. Du L, Zhao G, Yang Y, Qiu H, Wang L *et al*. A conformation-dependent neutralizing monoclonal antibody specifically targeting receptor-binding domain in Middle East respiratory syndrome coronavirus spike protein. *J Virol* 2014;88:7045–7053.
77. Ying T, Du L, Ju TW, Prabakaran P, Lau CC *et al*. Exceptionally potent neutralization of Middle East respiratory syndrome coronavirus by human monoclonal antibodies. *J Virol* 2014;88:7796–7805.
78. Agrawal AS, Ying T, Tao X, Garron T, Algaissi A *et al*. Passive transfer of A Germline-like neutralizing human monoclonal antibody protects transgenic mice against Lethal Middle East respiratory syndrome coronavirus infection. *Sci Rep* 2016;6:31629.
79. Houser KV, Gretebeck L, Ying T, Wang Y, Vogel L *et al*. Prophylaxis with a Middle East respiratory syndrome coronavirus (MERS-CoV)-Specific human monoclonal antibody protects rabbits from MERS-CoV infection. *J Infect Dis* 2016;213:1557–1561.
80. Jiang L, Wang N, Zuo T, Shi X, Poon KM *et al*. Potent neutralization of MERS-CoV by human neutralizing monoclonal antibodies to the viral spike glycoprotein. *Sci Transl Med* 2014;6:234ra59.
81. Yu X, Zhang S, Jiang L, Cui Y, Li D *et al*. Structural basis for the neutralization of MERS-CoV by a human monoclonal antibody MERS-27. *Sci Rep* 2015;5:13133.
82. Li Y, Wan Y, Liu P, Zhao J, Lu G *et al*. A humanized neutralizing antibody against MERS-CoV targeting the receptor-binding domain of the spike protein. *Cell Res* 2015;25:1237–1249.
83. Qiu H, Sun S, Xiao H, Feng J, Guo Y *et al*. Single-dose treatment with a humanized neutralizing antibody affords full protection of a human transgenic mouse model from lethal Middle East respiratory syndrome (MERS)-coronavirus infection. *Antiviral Res* 2016;132:141–148.
84. Corti D, Passini N, Lanzavecchia A, Zambon M. Rapid generation of a human monoclonal antibody to combat Middle East respiratory syndrome. *J Infect Public Health* 2016;9:231–235.
85. Johnson RF, Bagci U, Keith L, Tang X, Mollura DJ *et al*. 3B11-N, a monoclonal antibody against MERS-CoV, reduces lung pathology in rhesus monkeys following intratracheal inoculation of MERS-CoV Jordan-n3/2012. *Virology* 2016;490:49–58.
86. Ohnuma K, Haagmans BL, Hatano R, Raj VS, Mou H *et al*. Inhibition of Middle East respiratory syndrome coronavirus infection by anti-CD26 monoclonal antibody. *J Virol* 2013;87:13892–13899.
87. Lu L, Liu Q, Zhu Y, Chan KH, Qin L *et al*. Structure-based discovery of Middle East respiratory syndrome coronavirus fusion inhibitor. *Nat Commun* 2014;5:3067.
88. Liu S, Xiao G, Chen Y, He Y, Niu J *et al*. Interaction between heptad repeat 1 and 2 regions in spike protein of SARS-associated coronavirus: implications for virus fusogenic mechanism and identification of fusion inhibitors. *Lancet* 2004;363:938–947.
89. Bosch BJ, Martina BE, van der Zee R, Lepault J, Haijema BJ *et al*. Severe acute respiratory syndrome coronavirus (SARS-CoV) infection inhibition using spike protein heptad repeat-derived peptides. *Proc Natl Acad Sci USA* 2004;101:8455–8460.
90. Channappanavar R, Lu L, Xia S, Du L, Meyerholz DK *et al*. Protective effect of intranasal regimens containing peptidic Middle East respiratory syndrome coronavirus fusion inhibitor against MERS-CoV infection. *J Infect Dis* 2015;212:1894–1903.
91. Du L, Kou Z, Ma C, Tao X, Wang L *et al*. A truncated receptor-binding domain of MERS-CoV spike protein potently inhibits MERS-CoV infection and induces strong neutralizing antibody responses: implication for developing therapeutics and vaccines. *PLoS One* 2013;8:e81587.
92. Ma C, Li Y, Wang L, Zhao G, Tao X *et al*. Intranasal vaccination with recombinant receptor-binding domain of MERS-CoV spike protein induces much stronger local mucosal immune responses than subcutaneous immunization: implication for designing novel mucosal MERS vaccines. *Vaccine* 2014;32:2100–2108.
93. Zhang N, Tang J, Lu L, Jiang S, Du L *et al*. Receptor-binding domain-based subunit vaccines against MERS-CoV. *Virus Res* 2015;202:151–159.
94. Zhang N, Channappanavar R, Ma C, Wang L, Tang J *et al*. Identification of an ideal adjuvant for receptor-binding domain-based subunit vaccines against Middle East respiratory syndrome coronavirus. *Cell Mol Immunol* 2016;13:180–190.
95. Tai W, Wang Y, Fett CA, Zhao G, Li F *et al*. Recombinant receptor-binding domains of multiple Middle East respiratory syndrome coronaviruses (MERS-CoVs) induce cross-neutralizing antibodies against divergent human and camel MERS-CoVs and antibody escape mutants. *J Virol* 2017;91:e01651–16.
96. Coleman CM, Liu YV, Mu H, Taylor JK, Massare M *et al*. Purified coronavirus spike protein nanoparticles induce coronavirus neutralizing antibodies in mice. *Vaccine* 2014;32:3169–3174.
97. Excler JL, Delvecchio CJ, Wiley RE, Williams M, Yoon IK *et al*. Toward developing a preventive MERS-CoV Vaccine-Report from a workshop organized by the Saudi Arabia Ministry of Health and the International Vaccine Institute, Riyadh, Saudi Arabia, November 14–15, 2015. *Emerg Infect Dis* 2016;22:e1–e7.
98. Volz A, Kupke A, Song F, Jany S, Fux R *et al*. Protective efficacy of recombinant modified vaccinia virus Ankara delivering Middle East respiratory syndrome coronavirus spike glycoprotein. *J Virol* 2015;89:8651–8656.
99. Guo X, Deng Y, Chen H, Lan J, Wang W *et al*. Systemic and mucosal immunity in mice elicited by a single immunization with human adenovirus type 5 or 41 vector-based vaccines carrying the spike protein of Middle East respiratory syndrome coronavirus. *Immunology* 2015;145:476–484.
100. Malczyk AH, Kupke A, Prüfer S, Scheuplein VA, Hutzler S *et al*. A highly immunogenic and protective Middle East respiratory syndrome coronavirus vaccine based on a recombinant measles virus vaccine platform. *J Virol* 2015;89:11654–11667.
101. Haagmans BL, van den Brand JM, Raj VS, Volz A, Wohlsein P *et al*. An orthopoxvirus-based vaccine reduces virus excretion after MERS-CoV infection in dromedary camels. *Science* 2016;351:77–81.
102. Inovio. 2016. GLS-5300 SynCon® immunotherapy targeting Middle East Respiratory Syndrome. [www.inovio.com/products/infectious-disease-vaccines/mers/](http://www.inovio.com/products/infectious-disease-vaccines/mers/) [Accessed 18 January 2017].
103. Muthumani K, Falzarano D, Reuschel EL, Tingey C, Flingai S *et al*. A synthetic consensus anti-spike protein DNA vaccine induces protective immunity against Middle East respiratory syndrome coronavirus in nonhuman primates. *Sci Transl Med* 2015;7:301ra132.
104. Modjarrad K. MERS-CoV vaccine candidates in development: the current landscape. *Vaccine* 2016;34:2982–2987.
105. Lau SK, Lau CC, Chan KH, Li CP, Chen H, Skp L, Ccy L *et al*. Delayed induction of proinflammatory cytokines and suppression of innate antiviral response by the novel Middle East respiratory syndrome coronavirus: implications for pathogenesis and treatment. *J Gen Virol* 2013;94:2679–2690.
106. Totura AL, Whitmore A, Agnihothram S, Schäfer A, Katze MG *et al*. Toll-Like receptor 3 signaling via TRIF contributes to a protective innate immune response to severe acute respiratory syndrome coronavirus infection. *MBio* 2015;6:e00638–15–15.

107. de Wilde AH, Raj VS, Oudshoorn D, Bestebroer TM, van Nieuwkoop S *et al*. MERS-coronavirus replication induces severe *in vitro* cytopathology and is strongly inhibited by cyclosporin A or interferon- $\alpha$  treatment. *J Gen Virol* 2013;94:1749–1760.
108. Falzarano D, de Wit E, Martellaro C, Callison J, Munster VJ *et al*. Inhibition of novel  $\beta$  coronavirus replication by a combination of interferon- $\alpha$ 2b and ribavirin. *Sci Rep* 2013;3:1686.
109. Falzarano D, de Wit E, Rasmussen AL, Feldmann F, Okumura A *et al*. Treatment with interferon- $\alpha$ 2b and ribavirin improves outcome in MERS-CoV-infected rhesus macaques. *Nat Med* 2013;19:1313–1317.
110. Zheng Y, Wang QY. [Bioinformatics analysis on molecular mechanism of Ribavirin and interferon- $\alpha$  in treating MERS-CoV]. *Zhonghua Liu Xing Bing Xue Za Zhi* 2016;37:291–293.
111. Chan JF-W, Yao Y, Yeung M-L, Deng W, Bao L *et al*. Treatment with lopinavir/ritonavir or interferon- $\beta$ 1b improves outcome of MERS-CoV infection in a nonhuman primate model of common marmoset. *J Infect Dis* 2015;212:1904–1913.
112. Khalid M, Khan B, Al Rabiah F, Alismaili R, Saleemi S *et al*. Middle Eastern Respiratory syndrome Corona virus (MERS CoV): case reports from a tertiary care hospital in Saudi Arabia. *Ann Saudi Med* 2014;34:396–400.
113. Omrani AS, Saad MM, Baig K, Bahloul A, Abdul-Matin M *et al*. Ribavirin and interferon alfa-2a for severe Middle East respiratory syndrome coronavirus infection: a retrospective cohort study. *Lancet Infect Dis* 2014;14:1090–1095.
114. Qian Z, Dominguez SR, Holmes KV. Role of the spike glycoprotein of human Middle East respiratory syndrome coronavirus (MERS-CoV) in virus entry and syncytia formation. *PLoS One* 2013;8:e76469.
115. Gierer S, Bertram S, Kaup F, Wrensch F, Heurich A *et al*. The spike protein of the emerging betacoronavirus EMC uses a novel coronavirus receptor for entry, can be activated by TMPRSS2, and is targeted by neutralizing antibodies. *J Virol* 2013;87:5502–5511.
116. Millet JK, Whittaker GR. Host cell entry of Middle East respiratory syndrome coronavirus after two-step, furin-mediated activation of the spike protein. *Proc Natl Acad Sci USA* 2014;111:15214–15219.
117. Forni D, Filippi G, Cagliani R, de Gioia L, Pozzoli U *et al*. The heptad repeat region is a major selection target in MERS-CoV and related coronaviruses. *Sci Rep* 2015;5:14480.
118. Yamamoto M, Matsuyama S, Li X, Takeda M, Kawaguchi Y *et al*. Identification of nafamostat as a potent inhibitor of Middle East respiratory syndrome coronavirus S protein-mediated membrane fusion using the split-protein-based cell-cell fusion assay. *Antimicrob Agents Chemother* 2016;60:6532–6539.
119. Zhou N, Pan T, Zhang J, Li Q, Zhang X *et al*. Glycopeptide antibiotics potentially inhibit cathepsin L in the late endosome/lysosome and block the entry of Ebola virus, Middle East respiratory syndrome coronavirus (MERS-CoV), and severe acute respiratory syndrome coronavirus (SARS-CoV). *J Biol Chem* 2016;291:9218–9232.
120. Lei J, Mesters JR, Drosten C, Anemüller S, Ma Q *et al*. Crystal structure of the papain-like protease of MERS coronavirus reveals unusual, potentially druggable active-site features. *Antiviral Res* 2014;109:72–82.
121. Yang X, Chen X, Bian G, Tu J, Xing Y *et al*. Proteolytic processing, deubiquitinase and interferon antagonist activities of Middle East respiratory syndrome coronavirus papain-like protease. *J Gen Virol* 2014;95:614–626.
122. Lee H, Lei H, Santarsiero BD, Gatuz JL, Cao S *et al*. Inhibitor recognition specificity of MERS-CoV papain-like protease may differ from that of SARS-CoV. *ACS Chem Biol* 2015;10:1456–1465.
123. Clasman JR, Báez-Santos YM, Mettelman RC, O'Brien A, Baker SC *et al*. X-ray Structure and Enzymatic Activity Profile of a Core Papain-like Protease of MERS Coronavirus with utility for structure-based drug design. *Sci Rep* 2017;7:40292.
124. Wu CY, Jan JT, Ma SH, Kuo CJ, Juan HF *et al*. Small molecules targeting severe acute respiratory syndrome human coronavirus. *Proc Natl Acad Sci USA* 2004;101:10012–10017.
125. de Wilde AH, Jochmans D, Posthuma CC, Zevenhoven-Dobbe JC, van Nieuwkoop S *et al*. Screening of an FDA-approved compound library identifies four small-molecule inhibitors of Middle East respiratory syndrome coronavirus replication in cell culture. *Antimicrob Agents Chemother* 2014;58:4875–4884.
126. Hayden FG, Farrar J, Peiris JS. Towards improving clinical management of Middle East respiratory syndrome coronavirus infection. *Lancet Infect Dis* 2014;14:544–546.
127. Dyall J, Coleman CM, Hart BJ, Venkataraman T, Holbrook MR *et al*. Repurposing of clinically developed drugs for treatment of Middle East respiratory syndrome coronavirus infection. *Antimicrob Agents Chemother* 2014;58:4885–4893.
128. Belouzard S, Millet JK, Licitra BN, Whittaker GR. Mechanisms of coronavirus cell entry mediated by the viral spike protein. *Viruses* 2012;4:1011–1033.

### Five reasons to publish your next article with a Microbiology Society journal

1. The Microbiology Society is a not-for-profit organization.
2. We offer fast and rigorous peer review – average time to first decision is 4–6 weeks.
3. Our journals have a global readership with subscriptions held in research institutions around the world.
4. 80% of our authors rate our submission process as 'excellent' or 'very good'.
5. Your article will be published on an interactive journal platform with advanced metrics.

Find out more and submit your article at [microbiologyresearch.org](http://microbiologyresearch.org).