

Two-Dimensional Partition Theorem: Math Derivation

matousek-partition-tree / docs

Contents

1	Two-Dimensional Partition Theorem: Math Derivation	1
1.1	Goal	1
1.2	Cutting Inputs	2
1.3	Step 1: Build the Finite Test Set Q	2
1.4	Step 2: Test Set Lemma	5
1.5	Step 3: Construct Π for Fixed Q	9
1.6	Step 4: Weight Recurrence	11
1.7	Step 5: Product Estimate	13
1.8	Step 6: Bound K_Q	14
1.9	Step 7: Extend from Q to All Lines	15
1.10	Appendix: Proof of the 2D Cutting Lemma	15
1.11	Final Statement	18
1.12	Preprocessing Time: Building the Partition Tree	18
1.13	References	20

Two-Dimensional Partition Theorem: Math Derivation

Source: planar specialization of Matousek's Partition Theorem from *Efficient Partition Trees*. The cutting theorem used as a black box is the Chazelle-Friedman cutting machinery.

Goal

Input:

$$P = \{p_1, \dots, p_n\} \subset \mathbb{R}^2$$

Choose:

$$2 \leq s < n$$

Set:

$$r = \frac{n}{s}$$

Construct:

$$\Pi = \{(P_1, \Delta_1), \dots, (P_m, \Delta_m)\}$$

such that:

$$P_i \cap P_j = \emptyset \quad (i \neq j)$$

$$\bigcup_i P_i = P$$

$$P_i \subseteq \Delta_i$$

$$s \leq |P_i| < 2s$$

and:

$$\text{cr}(\Pi) = O(\sqrt{r})$$

For a line h :

$$\text{cr}_{\Pi}(h) = |\{i : h \text{ crosses } \Delta_i\}|$$

and:

$$\text{cr}(\Pi) = \max_h \text{cr}_{\Pi}(h)$$

Cutting Inputs

For n lines in the plane, a $(1/t)$ -cutting has:

$$O(t^2)$$

cells, and each cell is crossed by:

$$O\left(\frac{n}{t}\right)$$

lines.

Weighted version:

For weighted lines (Q, w) , define:

$$W = \sum_{q \in Q} w(q)$$

A weighted $(1/t)$ -cutting satisfies for every cell C :

$$\sum_{q \in Q(C)} w(q) \leq \frac{W}{t}$$

where:

$$Q(C) = \{q \in Q : q \text{ crosses } C\}$$

In 2D, for some absolute constant A:

$$\# \text{faces} \leq At^2$$

Step 1: Build the Finite Test Set Q

Use duality:

$$p = (a, b) \longleftrightarrow p^* : y = ax - b$$

$$\ell : y = ax - b \longleftrightarrow \ell^* = (a, b)$$

Incidence:

$$p \in \ell \iff \ell^* \in p^*$$

Above-below:

$$p \text{ above } \ell \iff \ell^* \text{ above } p^*$$

Verification:

Let:

$$p = (u, v)$$

and:

$$\ell : y = ax - b$$

Then:

$$p \text{ above } \ell \iff v > au - b$$

Also:

$$p^* : y = ux - v$$

and:

$$\ell^* = (a, b)$$

The dual point ℓ^* is above the dual line p^* iff:

$$b > ua - v$$

which is equivalent to:

$$v > au - b$$

Dualize P:

$$P^* = \{p_1^*, \dots, p_n^*\}$$

Choose:

$$t = \beta\sqrt{r}$$

with beta small enough that the cutting has at most r vertices.

Since:

$$O(t^2) = O(\beta^2 r) \leq r$$

the vertex set:

$$V = \{v_1, \dots, v_N\}$$

satisfies:

$$N \leq r$$

Dualize vertices back:

$$q_j = v_j^*$$

Define:

$$Q = \{q_1, \dots, q_N\}$$

Thus:

$$|Q| = N \leq r$$

Important object distinction:

$$R \subseteq P^*$$

is the sampled set of about t dual lines used to build the cutting.

$$V$$

is the set of cutting vertices, which are points in the dual plane.

$$Q = V^*$$

is a new set of primal lines.

Thus:

$$Q \not\subseteq P^*$$

The cutting parameter is:

$$t = \Theta(\sqrt{r})$$

so every dual cutting cell A is crossed by at most:

$$O\left(\frac{n}{\sqrt{r}}\right)$$

dual lines from P^* .

Step 2: Test Set Lemma

Assume Π is any simplicial partition with:

$$|P_i| \geq s$$

Define:

$$K_Q = \max_{q \in Q} cr_{\Pi}(q)$$

Claim:

$$\forall h, \quad cr_{\Pi}(h) \leq 3K_Q + O\left(\frac{n}{s\sqrt{r}}\right)$$

Proof skeleton:

Take any nonvertical line h. Its dual point is:

$$h^*$$

Let A be the triangular dual cutting cell with:

$$h^* \in \text{int}(A)$$

Let:

$$A = \text{conv}\{v_1, v_2, v_3\}$$

Define:

$$q_j = v_j^*$$

Then:

$$q_1, q_2, q_3 \in Q$$

Cells crossed by at least one of q_1, q_2, q_3 are at most:

$$3K_Q$$

Detailed count:

$$\text{cr}_\Pi(q_1) \leq K_Q$$

$$\text{cr}_\Pi(q_2) \leq K_Q$$

$$\text{cr}_\Pi(q_3) \leq K_Q$$

Therefore:

$$|\{\Delta_i : \Delta_i \text{ crossed by at least one of } q_1, q_2, q_3\}| \leq K_Q + K_Q + K_Q = 3K_Q$$

Call Δ_i bad if:

$$h \text{ crosses } \Delta_i$$

but:

$$q_1, q_2, q_3 \text{ do not cross } \Delta_i$$

Key geometric implication:

$$\Delta_i \text{ bad and } p \in P_i \implies p^* \cap A \neq \emptyset$$

Reason in sign language:

The three primal lines q_1, q_2, q_3 decompose the primal plane into sign cells.

The all-above cell is:

$$\{x : x \text{ is above } q_1, q_2, q_3\}$$

The all-below cell is:

$$\{x : x \text{ is below } q_1, q_2, q_3\}$$

If h crossed the all-above cell, there would be:

$$x \in h$$

with:

x above q_1, q_2, q_3

By duality:

$$q_1^* = v_1, \quad q_2^* = v_2, \quad q_3^* = v_3$$

all lie on the same side of x^* .

Since:

$$x \in h$$

we have:

$$h^* \in x^*$$

So x^* is a line through an interior point of triangle A.

But a line through an interior point of a triangle cannot put all three triangle vertices on the same side. Contradiction.

The same applies to the all-below cell.

Therefore every sign cell crossed by h is mixed-sign.

Now let:

$$p \in P_i \subseteq \Delta_i$$

where Δ_i is bad.

Because q_1, q_2, q_3 do not cross Δ_i , the simplex Δ_i lies inside one sign cell of their arrangement.

Since h crosses Δ_i , that sign cell is crossed by h , hence is mixed-sign.

So p has mixed signs relative to:

$$q_1, q_2, q_3$$

By above-below duality, the vertices:

$$v_1, v_2, v_3$$

are not all on the same side of p^* .

If p^* did not cross A, then the convex triangle:

$$A = \text{conv}\{v_1, v_2, v_3\}$$

would lie entirely on one side of p^* , so all three vertices would be on the same side.

Contradiction. Thus:

$$p^* \cap A \neq \emptyset$$

Equivalently:

bad primal simplex $\implies$ dual conflict with A

Since A is a $(1/\sqrt{r})$ -cutting cell:

$$|\{p \in P : p^* \cap A \neq \emptyset\}| = O\left(\frac{n}{\sqrt{r}}\right)$$

Bad groups are disjoint and each has at least s points, so:

$$\#\text{bad simplices} \leq O\left(\frac{n/\sqrt{r}}{s}\right)$$

Detailed division:

Let B be the number of bad simplices.

Each bad simplex contributes at least s original points:

$$|P_i| \geq s$$

The bad groups are disjoint, so total bad points are at least:

$$Bs$$

But the dual conflict bound gives total bad points at most:

$$O\left(\frac{n}{\sqrt{r}}\right)$$

Thus:

$$Bs \leq O\left(\frac{n}{\sqrt{r}}\right)$$

and:

$$B \leq O\left(\frac{n}{s\sqrt{r}}\right)$$

Thus:

$$\#\text{bad simplices} = O\left(\frac{n}{s\sqrt{r}}\right)$$

Therefore:

$$\text{cr}_{\Pi}(h) \leq 3K_Q + O\left(\frac{n}{s\sqrt{r}}\right)$$

Since:

$$r = \frac{n}{s}$$

we have:

$$\frac{n}{s\sqrt{r}} = \frac{r}{\sqrt{r}} = \sqrt{r}$$

So:

$$K_Q = O(\sqrt{r}) \implies \text{cr}_\Pi(h) = O(\sqrt{r})$$

Step 3: Construct Π for Fixed Q

At round i :

$$P^{(i)} = P \setminus (P_1 \cup \dots \cup P_i)$$

$$n_i = |P^{(i)}|$$

For test line q :

$$\kappa_i(q) = |\{j \leq i : q \text{ crosses } \Delta_j\}|$$

Define exponential weight:

$$w_i(q) = 2^{\kappa_i(q)}$$

Total weight:

$$W_i = \sum_{q \in Q} w_i(q)$$

Initially:

$$\kappa_0(q) = 0$$

$$w_0(q) = 1$$

$$W_0 = |Q|$$

If:

$$n_i < 2s$$

make the terminal group:

$$P_{i+1} = P^{(i)}$$

and stop.

Otherwise choose:

$$t_i = \alpha \sqrt{\frac{n_i}{s}}$$

where alpha is small enough that:

$$A\alpha^2 \leq 1$$

Then:

$$At_i^2 = A\alpha^2 \frac{n_i}{s} \leq \frac{n_i}{s}$$

Known and chosen quantities in this formula:

$$n_i = |P^{(i)}|$$

is known at the start of round i.

$$s$$

is the fixed group-size parameter.

$$A$$

is the absolute constant from the weighted cutting theorem.

$$t_i$$

is chosen from these quantities.

The reason is the pigeonhole constraint:

$$\# \text{faces} \leq At_i^2 \leq \frac{n_i}{s}$$

so the average number of remaining points per face is at least:

$$\frac{n_i}{n_i/s} = s$$

Build a weighted $(1/t_i)$ -cutting for (Q, w_i) .

For every face C:

$$w_i(Q(C)) \leq \frac{W_i}{t_i}$$

Number of faces:

$$\leq \frac{n_i}{s}$$

Pigeonhole:

$$\exists F_{i+1} \quad |P^{(i)} \cap F_{i+1}| \geq s$$

Set:

$$\Delta_{i+1} = F_{i+1}$$

Choose:

$$P_{i+1} \subseteq P^{(i)} \cap \Delta_{i+1}$$

with:

$$|P_{i+1}| = s$$

Remove:

$$P^{(i+1)} = P^{(i)} \setminus P_{i+1}$$

Step 4: Weight Recurrence

Let:

$$Q_{i+1} = \{q \in Q : q \text{ crosses } \Delta_{i+1}\}$$

For $q \in Q_{i+1}$:

$$w_{i+1}(q) = 2w_i(q)$$

For $q \notin Q_{i+1}$:

$$w_{i+1}(q) = w_i(q)$$

Therefore:

$$W_{i+1} = W_i + w_i(Q_{i+1})$$

No-step derivation:

Split Q into crossing and noncrossing test lines:

$$Q = Q_{i+1} \cup (Q \setminus Q_{i+1})$$

Then:

$$W_{i+1} = \sum_{q \notin Q_{i+1}} w_{i+1}(q) + \sum_{q \in Q_{i+1}} w_{i+1}(q)$$

For noncrossing lines:

$$w_{i+1}(q) = w_i(q)$$

For crossing lines:

$$w_{i+1}(q) = 2w_i(q)$$

So:

$$W_{i+1} = \sum_{q \notin Q_{i+1}} w_i(q) + \sum_{q \in Q_{i+1}} 2w_i(q)$$

Rewrite:

$$W_{i+1} = \sum_{q \in Q} w_i(q) + \sum_{q \in Q_{i+1}} w_i(q)$$

Therefore:

$$W_{i+1} = W_i + w_i(Q_{i+1})$$

Since Δ_{i+1} is a cutting face:

$$w_i(Q_{i+1}) \leq \frac{W_i}{t_i}$$

Hence:

$$W_{i+1} \leq W_i \left(1 + \frac{1}{t_i}\right)$$

In nonterminal rounds:

$$n_i = n - is$$

and:

$$\frac{n_i}{s} = r - i$$

Thus:

$$t_i = \Theta(\sqrt{r - i})$$

so for a constant C:

$$\frac{1}{t_i} \leq \frac{C}{\sqrt{r-i}}$$

Therefore:

$$W_{i+1} \leq W_i \left(1 + \frac{C}{\sqrt{r-i}}\right)$$

The terminal round can at most double every weight:

$$W_m \leq 2W_L$$

where L is the number of nonterminal rounds.

Step 5: Product Estimate

Since:

$$L \leq r$$

we get:

$$W_m \leq 2|Q| \prod_{i=0}^{L-1} \left(1 + \frac{C}{\sqrt{r-i}}\right)$$

Taking logs:

$$\log W_m \leq O(1) + \log |Q| + \sum_{i=0}^{L-1} \log \left(1 + \frac{C}{\sqrt{r-i}}\right)$$

Using:

$$\log(1+x) \leq x$$

gives:

$$\log W_m \leq O(1) + \log |Q| + C \sum_{i=0}^{L-1} \frac{1}{\sqrt{r-i}}$$

Since:

$$\sum_{i=0}^{L-1} \frac{1}{\sqrt{r-i}} \leq \sum_{j=1}^r \frac{1}{\sqrt{j}}$$

and:

$$\sum_{j=1}^r j^{-1/2} \leq 1 + \int_1^r x^{-1/2} dx$$

with:

$$\int_1^r x^{-1/2} dx = 2(\sqrt{r} - 1)$$

we obtain:

$$\sum_{j=1}^r j^{-1/2} = O(\sqrt{r})$$

Therefore:

$$\log W_m = O(\log |Q| + \sqrt{r})$$

Step 6: Bound K_Q

For any $q \in Q$:

$$w_m(q) = 2^{\kappa_m(q)}$$

and:

$$w_m(q) \leq W_m$$

Thus:

$$2^{\kappa_m(q)} \leq W_m$$

Taking logs:

$$\kappa_m(q) \leq \log_2 W_m$$

Hence:

$$\kappa_m(q) = O(\log |Q| + \sqrt{r})$$

Therefore:

$$K_Q = \max_{q \in Q} \kappa_m(q) = O(\log |Q| + \sqrt{r})$$

Since:

$$|Q| \leq r$$

we have:

$$\log |Q| \leq \log r$$

and:

$$\log r = O(\sqrt{r})$$

Thus:

$$K_Q = O(\sqrt{r})$$

Here:

$$\kappa_i(q)$$

is cumulative for one q up to round i .

$$K_Q$$

is the final maximum over all q in Q .

At the end:

$$K_Q = \max_{q \in Q} \kappa_m(q)$$

Step 7: Extend from Q to All Lines

From the Test Set Lemma:

$$\text{cr}_\Pi(h) \leq 3K_Q + O\left(\frac{n}{s\sqrt{r}}\right)$$

Substitute:

$$K_Q = O(\sqrt{r})$$

and:

$$\frac{n}{s\sqrt{r}} = \sqrt{r}$$

Then:

$$\text{cr}_\Pi(h) \leq 3O(\sqrt{r}) + O(\sqrt{r})$$

so:

$$\text{cr}_\Pi(h) = O(\sqrt{r})$$

for every line h .

Therefore:

$$\text{cr}(\Pi) = O(\sqrt{r})$$

Appendix: Proof of the 2D Cutting Lemma

Statement:

Given n lines H in the plane and a parameter t , there exists a cutting with:

$$O(t^2)$$

cells, such that every cell is crossed by at most:

$$O\left(\frac{n}{t}\right)$$

lines of H .

The weighted version replaces counts by total weights.

Proof skeleton:

Choose a random sample R from H of size:

$$O(t)$$

Build the arrangement of R and triangulate its cells. The arrangement of $O(t)$ lines has:

$$O(t^2)$$

faces.

For a triangle Δ , define its conflict list:

$$H_{\Delta} = \{h \in H : h \text{ crosses } \Delta\}$$

and its excess:

$$e(\Delta) = \frac{|H_{\Delta}|}{n/t} = \frac{|H_{\Delta}|t}{n}$$

If:

$$e(\Delta) \leq C$$

for a sufficiently large constant C , then Δ is already good, because:

$$|H_{\Delta}| \leq C \frac{n}{t}$$

If:

$$e(\Delta) > C$$

then refine Δ locally. Let:

$$n_\Delta = |H_\Delta|$$

To make each refined cell crossed by at most $O(n/t)$ original lines, use a local cutting for H_Δ with parameter:

$$u_\Delta = \frac{n_\Delta}{n/t} = \frac{n_\Delta t}{n} = e(\Delta)$$

This local refinement has:

$$O(u_\Delta^2) = O(e(\Delta)^2)$$

subcells, and each subcell is crossed by at most:

$$O\left(\frac{n_\Delta}{u_\Delta}\right) = O\left(\frac{n}{t}\right)$$

lines.

Thus the final cutting size is bounded by:

$$\sum_{\Delta} O(\max\{1, e(\Delta)^2\})$$

The key random-sampling estimate is:

$$\mathbb{E} [|\{\Delta : e(\Delta) \geq u\}|] \leq C_1 t^2 e^{-C_2 u}$$

for constants $C_1, C_2 > 0$.

Using dyadic levels:

$$\sum_{\Delta} \max\{1, e(\Delta)^2\} \leq O(t^2) + \sum_{j \geq 0} O(2^{2j}) \cdot |\{\Delta : 2^j \leq e(\Delta) < 2^{j+1}\}|$$

Taking expectations and applying the exponential tail bound:

$$\mathbb{E} \left[\sum_{\Delta} \max\{1, e(\Delta)^2\} \right] \leq O(t^2) + \sum_{j \geq 0} O(2^{2j}) \cdot C_1 t^2 e^{-C_2 2^j}$$

The series:

$$\sum_{j \geq 0} 2^{2j} e^{-C_2 2^j}$$

converges, so:

$$\mathbb{E} \left[\sum_{\Delta} \max\{1, e(\Delta)^2\} \right] = O(t^2)$$

Therefore some random sample produces a cutting of size:

$$O(t^2)$$

with every final cell crossed by at most:

$$O\left(\frac{n}{t}\right)$$

lines.

Weighted form:

For weighted lines (H, w) , let:

$$W = w(H)$$

Sample lines with probability proportional to weight, or equivalently replace each line h by $w(h)$ copies after scaling rational weights to integers. The same argument gives cells C satisfying:

$$w(H_C) \leq O\left(\frac{W}{t}\right)$$

After absorbing constants into the cutting parameter, this is the weighted cutting form used in the partition proof:

$$w(H_C) \leq \frac{W}{t}$$

Scope of this appendix:

This appendix explains why the cutting tool is valid and why the two-dimensional complexity is $O(t^2)$. The sharp optimal version relies on the exponential decay estimate above, which is the Chazelle-Friedman sampling argument cited in the references.

Final Statement

For every finite point set:

$$P \subset \mathbb{R}^2$$

and every:

$$2 \leq s < n$$

with:

$$r = \frac{n}{s}$$

there exists a simplicial partition:

$$\Pi = \{(P_1, \Delta_1), \dots, (P_m, \Delta_m)\}$$

such that:

$$s \leq |P_i| < 2s$$

and:

$$\text{cr}(\Pi) = O(\sqrt{r})$$

Preprocessing Time: Building the Partition Tree

This is a theorem-level bound for the *construction* of the whole tree, separate from the query recurrence. It is the theorem's guarantee, **not** a proved wall-clock bound for this Python implementation (which uses exact `Fraction` arithmetic and randomized cutting retries — see the caveat at the end).

Let $B(m)$ be the time to build a partition tree on m points. For any fixed $\delta > 0$, one simplicial partition at a node is constructed in $O(m^{1+\delta})$ time, so:

$$B(m) = O(m^{1+\delta}) + \sum_i B(m_i),$$

where the child classes are disjoint and each is a constant factor smaller:

$$\sum_i m_i = m, \quad m_i \leq \frac{2m}{r}.$$

Cost at a single depth. Fix a depth j of the recursion tree, and let V_j be the nodes at that depth. Their point sets are disjoint, so their sizes sum to at most m , and by applying the size shrinkage j times the largest one is bounded:

$$\sum_{v \in V_j} m_v \leq m, \quad \max_{v \in V_j} m_v \leq m \left(\frac{2}{r}\right)^j.$$

Split each per-node cost as $m_v^{1+\delta} = m_v m_v^\delta$ and pull out the max on the δ factor:

$$\begin{aligned} C_j &= \sum_{v \in V_j} O(m_v^{1+\delta}) = O\left(\sum_{v \in V_j} m_v m_v^\delta\right) \\ &\leq O\left(\left(\max_{v \in V_j} m_v\right)^\delta \sum_{v \in V_j} m_v\right) \leq O\left(m^{1+\delta} \left(\frac{2}{r}\right)^{j\delta}\right). \end{aligned}$$

Sum over all depths. For fixed $r > 2$ the per-level factor is below one, so the level costs form a convergent geometric series:

$$\left(\frac{2}{r}\right)^\delta < 1 \implies \sum_{j \geq 0} \left(\frac{2}{r}\right)^{j\delta} = \frac{1}{1 - \left(\frac{2}{r}\right)^\delta} = O(1).$$

Therefore:

$$B(m) = O\left(m^{1+\delta} \sum_{j \geq 0} \left(\frac{2}{r}\right)^{j\delta}\right) = O(m^{1+\delta}).$$

Matching the stated bound. Taking $\delta = \varepsilon/2$ gives the stronger intermediate bound $O(n^{1+\varepsilon/2})$, and hence the standard stated preprocessing bound:

$$B(n) = O(n^{1+\varepsilon}).$$

Note that $O(n \log n) \subsetneq O(n^{1+\varepsilon})$ strictly: for every fixed $\varepsilon > 0$ we have $\log n = O(n^\varepsilon)$ but not the reverse. Matoušek's tighter $O(n \log n)$ preprocessing is a separate result that charges construction against the partition structure rather than this level-by-level geometric series.

Caveat for this implementation. The bound above is the theoretical guarantee. It does **not** claim this Python code runs in $O(n^{1+\varepsilon})$ wall-clock time: that would additionally require a bit-complexity analysis of exact `Fraction` arithmetic and a runtime analysis of the randomized cutting retries and runtime verification. This code trades all of that away for exact arithmetic and verifiable preconditions, and is correspondingly slow.

References

- Jiri Matousek, **Efficient Partition Trees**, *Discrete & Computational Geometry* 8, 315-334, 1992.
- Bernard Chazelle and Joel Friedman, **A deterministic view of random sampling and its use in geometry**, *Combinatorica* 10, 229-249, 1990.