

# Last Time

---

Gauss' Law

# Today

---

Gauss' Law: Examples  
"Magnetic Gauss Law"  
Ampere's Law

# Gauss' Law for Point Charge


$$\oint \vec{E} \cdot \hat{n} dA = \frac{1}{\epsilon_o} \sum q_{\text{inside}} \quad \begin{array}{l} \text{Gauss'} \\ \text{Law} \end{array}$$

$$\vec{E} = \frac{1}{4\pi\epsilon_o} \frac{q}{r^2} \hat{r} \quad \begin{array}{l} \text{E-field} \\ \text{Point Charge} \end{array}$$

Gauss' Law for Point Charge:

$$\oint \vec{E} \cdot \hat{n} dA = \frac{1}{4\pi\epsilon_o} \frac{q}{r^2} (4\pi r^2) = \frac{q}{\epsilon_o}$$

Works for any size sphere because r cancels



On each sphere:

$$\text{Field } E \propto \frac{1}{r^2}$$

$$\text{Surface Area } A \propto r^2$$




# Gauss' Law for Point Charge

$$\oint \vec{E} \cdot \hat{n} dA = \frac{1}{\epsilon_o} \sum q_{\text{inside}} \quad \text{GAUSS' LAW}$$

Gauss' Law for Point Charge:

$$\oint \vec{E} \cdot \hat{n} dA = \frac{1}{4\pi\epsilon_o} \frac{q}{r^2} (4\pi r^2) = \frac{q}{\epsilon_o}$$

Works for any size sphere because r cancels



On each sphere:

$$\text{Field } E \propto \frac{1}{r^2}$$

$$\text{Surface Area } A \propto r^2$$


→ In any segment, the contribution from any "shell" is the **same**. (Like a flashlight.)

# Gauss' Law for Point Charge

$$\oint \vec{E} \cdot \hat{n} dA = \frac{1}{\epsilon_o} \sum q_{\text{inside}} \quad \text{GAUSS' LAW}$$

---

In any segment, the contribution from any "shell" is the **same**. (Like a flashlight.)



On each sphere:

$$\text{Field } E \propto \frac{1}{r^2}$$


$$\text{Surface Area } A \propto r^2$$

# Gauss' Law for Point Charge

$$\oint \vec{E} \cdot \hat{n} dA = \frac{1}{\epsilon_0} \sum q_{\text{inside}} \quad \text{GAUSS' LAW}$$

In any segment, the contribution from any "shell" is the **same**. (Like a flashlight.)

Surround the charge with any shape by following different spheres in different places.



On each sphere:

$$\text{Field } E \propto \frac{1}{r^2}$$

$$\text{Surface Area } A \propto r^2$$


# Gauss' Law for Point Charge

$$\oint \vec{E} \cdot \hat{n} dA = \frac{1}{\epsilon_0} \sum q_{\text{inside}} \quad \text{GAUSS' LAW}$$

In any segment, the contribution from any "shell" is the **same**. (Like a flashlight.)

Surround the charge with any shape by following different spheres in different places.

Flux through outer surface is always the **same**.




On each sphere:

$$\text{Field } E \propto \frac{1}{r^2}$$

$$\text{Surface Area } A \propto r^2$$

# Gauss' Law for Point Charge

$$\oint \vec{E} \cdot \hat{n} dA = \frac{1}{\epsilon_0} \sum q_{\text{inside}} \quad \text{GAUSS' LAW}$$



On each sphere:

$$\text{Field } E \propto \frac{1}{r^2}$$

$$\text{Surface Area } A \propto r^2$$

In any segment, the contribution from any "shell" is the **same**. (Like a flashlight.)

Surround the charge with any shape by following different spheres in different places.

Flux through outer surface is always the **same**.

Limit of small segments

→ works for any smooth shape.



# Example: Gauss' Law for Plane

$$\oint \vec{E} \cdot \hat{n} dA = \frac{1}{\epsilon_0} \sum q_{\text{inside}} \quad \text{GAUSS' LAW}$$

GIVEN: Infinite plane.

Surface charge density  $\sigma = [Q/A]$


FIND: E-field due to plane

Due to geometry of plane:

E-field is perpendicular to plane

E-field has same magnitude everywhere

→ USE Gauss' Law to find magnitude.



# Example: Gauss' Law for Plane

$$\oint \vec{E} \cdot \hat{n} dA = \frac{1}{\epsilon_o} \sum q_{\text{inside}} \quad \text{GAUSS' LAW}$$

**GIVEN:** Infinite plane.  
Surface charge density  $\sigma = [Q/A]$


**FIND:** E-field due to plane

Due to geometry of plane:

E-field is perpendicular to plane

E-field has same magnitude everywhere

→ USE Gauss' Law to find magnitude.



Choose "Gaussian box" wisely!

Same area A on ends of cylinders

E is constant and perpendicular

to the surface A of the "Gaussian box":

$$\oint \vec{E} \cdot \hat{n} dA = \frac{1}{\epsilon_o} q_{\text{inside}}$$
$$2EA = \frac{1}{\epsilon_o} \sigma A$$

$q_{\text{inside}} = [Q/A] * A = \sigma A$

$$E = \frac{1}{2\epsilon_o} \sigma$$

**ELECTRIC FIELD  
of a PLANE**



# Gauss' Law for Magnetism?

$$\oint \vec{E} \cdot \hat{n} dA = \frac{1}{\epsilon_0} \sum q_{\text{inside}} \quad \text{GAUSS' LAW for charge}$$

---

So far, no experiment has found a "magnetic charge"  
(*a.k.a. magnetic monopole*)

$$q_{\text{magnet}} = 0 \quad \text{Big fat ZERO!}$$

→ Gauss' Law for Magnetism is simpler:



$$\oint \vec{B} \cdot \hat{n} dA = 0$$

**GAUSS' LAW  
FOR MAGNETISM**

---

## **Next Up: Ampere's Law**

First review Biot-Savart Law

---


# Very Close to the Wire

$$\vec{B} = \left( \frac{\mu_o}{4\pi} \right) \frac{IL}{r \sqrt{r^2 + (L/2)^2}} \hat{\theta}$$

Very close to the wire:  $r \ll L$      $\sqrt{r^2 + (L/2)^2} \approx L/2$

$$\Rightarrow \vec{B} = \left( \frac{\mu_o}{4\pi} \right) \frac{IL}{r(L/2)} \hat{\theta} = \left( \frac{\mu_o}{4\pi} \right) \frac{2I}{r} \hat{\theta} = \vec{B}$$

**CLOSE TO  
THE WIRE**



<http://physick.wikispaces.com/Electric+Current>

Blast from the Past  
Lecture 13


# Very Long Wire

$$\vec{B} = \left( \frac{\mu_o}{4\pi} \right) \frac{IL}{r \sqrt{r^2 + (L/2)^2}} \hat{\theta}$$

Very Long wire:  $L \gg r$        $\sqrt{r^2 + (L/2)^2} \approx L/2$

$$\Rightarrow \vec{B} = \left( \frac{\mu_o}{4\pi} \right) \frac{IL}{r(L/2)} \hat{\theta} = \boxed{\left( \frac{\mu_o}{4\pi} \right) \frac{2I}{r} \hat{\theta} = \vec{B}}$$

**VERY LONG  
WIRE**




<http://physick.wikispaces.com/Electric+Current>

Blast from the Past  
Lecture 13

# Very Long Wire

$$\vec{B} = \left( \frac{\mu_o}{4\pi} \right) \frac{2I}{r} \hat{\theta}$$


---



Viewed from the end  
Current coming out of board

# Very Long Wire

$$\vec{B} = \left( \frac{\mu_o}{4\pi} \right) \frac{2I}{r} \hat{\theta}$$



Viewed from the end  
Current coming out of board

Cylindrical pattern of B-field

→ Let's take a line integral along one circle

$$\oint \vec{B} \cdot d\vec{l} = \oint \left( \frac{\mu_o}{4\pi} \right) \frac{2I}{r} \hat{\theta} \cdot d\vec{l}$$

$$= \left( \frac{\mu_o}{4\pi} \right) \frac{2I}{r} \oint \hat{\theta} \cdot d\vec{l}$$

$$= \left( \frac{\mu_o}{4\pi} \right) \frac{2I}{r} (2\pi r) = \mu_o I$$

$d\vec{l}$  is along our circle

$$d\vec{l} = r d\hat{\theta}$$

$$\oint \hat{\theta} \cdot d\vec{l} = r \oint \hat{\theta} d\hat{\theta} = 2\pi r$$

$$\oint \vec{B} \cdot d\vec{l} = \mu_o I$$

**AMPERE'S  
LAW**





# Gauss' Law for Point Charge

$$\oint \vec{E} \cdot \hat{n} dA = \frac{1}{\epsilon_o} \sum q_{\text{inside}} \quad \begin{array}{l} \text{Gauss'} \\ \text{Law} \end{array}$$

$$\vec{E} = \frac{1}{4\pi\epsilon_o} \frac{q}{r^2} \hat{r} \quad \begin{array}{l} \text{E-field} \\ \text{Point Charge} \end{array}$$


Gauss' Law for Point Charge:

$$\oint \vec{E} \cdot \hat{n} dA = \frac{1}{4\pi\epsilon_o} \frac{q}{r^2} (4\pi r^2) = \frac{q}{\epsilon_o}$$

Works for any size sphere because r cancels

Something similar is going to happen for B of a wire

Remember Me?  
Slide 3



On each sphere:

$$\text{Field } E \propto \frac{1}{r^2}$$


$$\text{Surface Area } A \propto r^2$$

# Ampere's Law for Long Wire

$$\vec{B} = \left( \frac{\mu_o}{4\pi} \right) \frac{2I}{r} \hat{\theta} \quad \begin{array}{l} \text{Biot-Savart} \\ \text{Law} \end{array}$$

$$\oint \vec{B} \cdot d\vec{l} = \mu_o I \quad \begin{array}{l} \text{B-field} \\ \text{Long Wire} \end{array}$$

Current coming out of board



On each circle:

$$\text{Field } B \propto \frac{1}{r}$$

$$\text{Circumference } C \propto r$$

Ampere's Law for Long Wire:

$$\oint \vec{B} \cdot d\vec{l} = \left( \frac{\mu_o}{4\pi} \right) \frac{2I}{r} (2\pi r) = \mu_o I$$

Works for any size circle because r cancels




# Ampere's Law for Long Wire

$$\vec{B} = \left( \frac{\mu_o}{4\pi} \right) \frac{2I}{r} \hat{\theta} \quad \begin{array}{l} \text{Biot-Savart} \\ \text{Law} \end{array}$$

$$\oint \vec{B} \cdot d\vec{l} = \mu_o I \quad \begin{array}{l} \text{B-field} \\ \text{Long Wire} \end{array}$$

Current coming out of board



On each circle:

$$\text{Field } B \propto \frac{1}{r}$$

$$\text{Circumference } C \propto r$$

Ampere's Law for Long Wire:

$$\oint \vec{B} \cdot d\vec{l} = \left( \frac{\mu_o}{4\pi} \right) \frac{2I}{r} (2\pi r) = \mu_o I$$

Works for any size circle because r cancels


→ In any segment, the contribution from any **circle** is the **same**. (Like a flashlight.)

# Ampere's Law for Long Wire

$$\vec{B} = \left( \frac{\mu_o}{4\pi} \right) \frac{2I}{r} \hat{\theta} \quad \begin{array}{l} \text{Biot-Savart} \\ \text{Law} \end{array}$$

$$\oint \vec{B} \cdot d\vec{l} = \mu_o I \quad \begin{array}{l} \text{B-field} \\ \text{Long Wire} \end{array}$$

Current coming out of board



In any segment, the contribution from any "circle" is the **same**. (Like a flashlight.)

On each circle:

$$\text{Field } B \propto \frac{1}{r}$$


$$\text{Circumference } C \propto r$$

# Ampere's Law for Long Wire

$$\vec{B} = \left( \frac{\mu_o}{4\pi} \right) \frac{2I}{r} \hat{\theta} \quad \begin{array}{l} \text{Biot-Savart} \\ \text{Law} \end{array}$$

$$\oint \vec{B} \cdot d\vec{l} = \mu_o I \quad \begin{array}{l} \text{B-field} \\ \text{Long Wire} \end{array}$$

Current coming out of board



On each circle:

$$\text{Field } B \propto \frac{1}{r}$$

$$\text{Circumference } C \propto r$$

In any segment, the contribution from any **circle** is the **same**. (Like a flashlight.)


Surround the **wire** with any shape by following different **circles** in different places.

# Ampere's Law for Long Wire

$$\vec{B} = \left( \frac{\mu_o}{4\pi} \right) \frac{2I}{r} \hat{\theta} \quad \begin{array}{l} \text{Biot-Savart} \\ \text{Law} \end{array}$$

$$\oint \vec{B} \cdot d\vec{l} = \mu_o I \quad \begin{array}{l} \text{B-field} \\ \text{Long Wire} \end{array}$$

Current coming out of board



On each circle:

$$\text{Field } B \propto \frac{1}{r}$$

$$\text{Circumference } C \propto r$$

In any segment, the contribution from any **circle** is the **same**. (Like a flashlight.)

Surround the **wire** with any path by following different **circles** in different places.


Line integral around outer surface is always **same**.

# Ampere's Law for Long Wire

$$\vec{B} = \left( \frac{\mu_o}{4\pi} \right) \frac{2I}{r} \hat{\theta} \quad \begin{array}{l} \text{Biot-Savart} \\ \text{Law} \end{array}$$

$$\oint \vec{B} \cdot d\vec{l} = \mu_o I \quad \begin{array}{l} \text{B-field} \\ \text{Long Wire} \end{array}$$

Current coming out of board



On each circle:

$$\text{Field } B \propto \frac{1}{r}$$

$$\text{Circumference } C \propto r$$

In any segment, the contribution from any **circle** is the **same**. (Like a flashlight.)

Surround the **wire** with any path by following different **circles** in different places.


Line integral around outer surface is always **same**.

Limit of small segments

→ works for any smooth path


# Electric Current is like Water Flow

---



Water flows through a net  
at a certain rate.

Molecules/second  
through each square



The net can deform,  
but the flow rate is the same.


Same number of  
molecules/second  
through each square



# Ampere's Law


Water flows through a net  
at a certain rate.

Charges/second  
through each square




The net can deform,  
but the flow rate is the same.

Same number of  
charges/second  
through each square



Likewise, Ampere's Law works  
even when the surface through which  
current  $I$  "pokes" is deformed,  
even if it means the wire "pokes"  
through in more places.



# Today

---

Gauss' Law: Examples  
"Magnetic Gauss Law"  
Ampere's Law