

A STACKY p -ADIC RIEMANN–HILBERT CORRESPONDENCE ON HITCHIN-SMALL LOCUS

YUDONG LIU, XIECHENG NIE, XIAOYU QU, AND YUPENG WANG

ABSTRACT. Let C be a algebraically closed perfectoid field over \mathbb{Q}_p with the ring of integer \mathcal{O}_C and the infinitesimal thickening \mathbf{A}_{inf} . Let \mathfrak{X} be a smooth formal scheme over \mathcal{O}_C with a fixed smooth lifting $\tilde{\mathfrak{X}}$ over \mathbf{A}_{inf} . Let X be the generic fiber of \mathfrak{X} and \tilde{X} be its lifting over \mathbf{B}_{dR}^+ induced by $\tilde{\mathfrak{X}}$. Let $\text{MIC}_r(\tilde{X})^{\text{H-small}}$ and $\text{LS}_r(X, \mathbb{B}_{\text{dR}}^+)^{\text{H-small}}$ be the v -stacks of rank- r Hitchin-small integrable connections on $X_{\text{ét}}$ and \mathbb{B}_{dR}^+ -local systems on X_v , respectively. In this paper, we establish an equivalence between this two stacks by introducing a new period sheaf with connection $(\mathcal{O}\mathbb{B}_{\text{dR,pd}}^+, d)$ on X_v .

CONTENTS

1.	Introduction	1
2.	The period sheaf with connection $(\mathcal{O}\mathbb{B}_{\text{dR,pd}}^+, d)$	6
3.	Local small Riemann–Hilbert correspondence	13
4.	The stacky Riemann–Hilbert correspondence: Proof of Theorem 1.1	22
	References	25

1. INTRODUCTION

1.1. Overview. For a projective smooth variety over the the field \mathbb{C} of complex numbers, the Riemann–Hilbert correspondence describes an equivalence between the category \mathbb{C} -local systems and the category of integrable connections on X . This equivalence upgrades to the moduli level; that is, there is an homeomorphism between the moduli space \mathbf{M}_B of \mathbb{C} -local systems and the moduli space \mathbf{M}_{dR} of integrable connections.

The p -adic Riemann–Hilbert correspondence aims to give an analogue of the above correspondence for rigid spaces; that is, it suggests an equivalence between the category of certain local systems and the category of certain integrable connections on a rigid spaces over a complete p -adic field. The first step to question is due to Scholze [Sch13]. In *loc.cit.*, for a smooth rigid variety X over a complete discrete valuation field K of mixed characteristic $(0, p)$ with the perfect residue field, he introduced so-call \mathbb{B}_{dR}^+ -local systems on the pro-étale site $X_{\text{proét}}$, constructed a period sheaf with connection $(\mathcal{O}\mathbb{B}_{\text{dR}}^+, d)$ on $X_{\text{proét}}$ whose de Rham complex gives a resolution of \mathbb{B}_{dR}^+ , and established an equivalence between the category of *de Rham* \mathbb{B}_{dR}^+ -local systems and the category of *filtered integrable connections* on $X_{\text{ét}}$. Based on his work, Liu and Zhu constructed two functors \mathcal{RH} and D_{dR} from the category of \mathbb{Q}_p -local systems on $X_{\text{ét}}$ to the category of G_K -equivariant filtered integrable connections on “ $(X \hat{\otimes}_K \mathbf{B}_{\text{dR}}^+)_{\text{ét}}$ ” and the category of filtered integrable connections on $X_{\text{ét}}$, respectively. Using this, they proved the rigidity for “being de Rham” of \mathbb{Q}_p -local systems [LZ17]. Their approach also works in the theory of log-geometry [DLLZ23]. Also, the functor \mathcal{RH} can be also defined for any \mathbb{B}_{dR}^+ -local systems. Indeed, in [GMW], one can establish an equivalence between the category of \mathbb{B}_{dR}^+ -local systems on $X_{\text{proét}}$ and the category of G_K -equivariant integrable connections on “ $X \hat{\otimes}_K \mathbf{B}_{\text{dR}}^+$ ”. This is used to classify \mathbb{B}_{dR}^+ -local systems with prismatic source (see [MW22], [AHLB23b] for relevant results in “mod t ” case). On the other hand, in the lecture note of his ICM talk, Bhatt described his joint work with Lurie [Bha21, Th. 5.4], a Riemann–Hilbert functor from the category of certain \mathbb{Q}_p -sheaves to the category of certain D -modules. Parts of their results are also obtained by Li [Li] independently by using a certain variant of $\mathcal{O}\mathbb{B}_{\text{dR}}^+$ and applying the approach of Liu–Zhu [LZ17]. All

2020 *Mathematics Subject Classification.* Primary .

Key words and phrases. p -adic Riemman-Hilbert correspondence, period sheaf, \mathbb{B}_{dR}^+ -local systems, flat connections, Hitchin-small locus.

results above require X is defined over a complete discrete valuation field K containing \mathbb{Q}_p with the perfect residue field.

Now, let C be a algebraically closed perfectoid field containing \mathbb{Q}_p with ring of integers \mathcal{O}_C and the infinitesimal thickening \mathbf{A}_{inf} . Very recently, starting with a smooth formal scheme \mathfrak{X} over \mathcal{O}_C admitting a smooth lifting $\tilde{\mathfrak{X}}$ over \mathbf{A}_{inf} , Chen, Liu, Zhu and the last author [CLWZ24] can constructed an overconvergent de Rham sheaf with connection $(\mathcal{O}\mathbb{B}_{\text{dR}}^{+,+}, d)$ whose de Rham complex is again a resolution of \mathbb{B}_{dR}^+ . Using this, they are able to establish an equivalence between the category of *small* \mathbb{B}_{dR}^+ -local systems on $X_{\text{proét}}$ and the category of *small* integrable connections on \tilde{X} , where X is the generic fiber of \mathfrak{X} and \tilde{X} is the lifting of X over \mathbf{B}_{dR}^+ induced by $\tilde{\mathfrak{X}}$. This generalizes Faltings' p -adic Simpson correspondence [Fal05]. Indeed, in *loc.cit.*, they even constructed a period sheaf with connection $(\mathcal{O}\tilde{\mathbb{C}}^I, d)$ which induces a resolution of relative Robba ring $\tilde{\mathbb{C}}^I$ and established a Riemann–Hilbert correspondence for small $\tilde{\mathbb{C}}^I$ -local systems.

In this paper, we give a slightly generalisation of the p -adic Riemann–Hilbert correspondence in [CLWZ24]. Again, let \mathfrak{X} be a smooth formal scheme over \mathcal{O}_C with a fixed smooth lifting $\tilde{\mathfrak{X}}$ over \mathbf{A}_{inf} . We shall establish an equivalence between the category of *Hitchin-small* \mathbb{B}_{dR}^+ -local systems on X_v , the v -site of X introduced in [Sch17], and the category of *Hitchin-small* integrable connections on \tilde{X} . Indeed, we shall prove this on the moduli level; that is, we shall give an equivalence between the moduli stacks of Hitchin-small \mathbb{B}_{dR}^+ -local systems and Hitchin-small integrable connections by constructing a new period sheaf with connection $(\mathcal{O}\mathbb{B}_{\text{dR,pd}}^+, d)$. This generalises the stacky p -adic Simpson correspondence in [AHLB23b].

1.2. Main results. Now let us state our main theorem. From now on, we always let C be a algebraically closed perfectoid field containing \mathbb{Q}_p with ring of integers \mathcal{O}_C , maximal ideal \mathfrak{m}_C of \mathcal{O}_C and the infinitesimal thickening \mathbf{A}_{inf} . Fix a compatible sequence $\{\zeta_{p^n}\}_{n \geq 0}$ of primitive p^n -th roots of unity in C . We also let \mathfrak{X} be a liftable smooth formal scheme over \mathcal{O}_C with a fixed lifting $\tilde{\mathfrak{X}}$ over \mathbf{A}_{inf} as before. Let X be the generic fiber of \mathfrak{X} and \tilde{X} be its lifting over \mathbf{B}_{dR}^+ induced by $\tilde{\mathfrak{X}}$. For any $n \geq 1$, let \tilde{X}_n be the reduction of \tilde{X} modulo t^n . Let Perfd be the v -site of all affinoid perfectoid spaces over $\text{Spa}(C, \mathcal{O}_C)$.

Theorem 1.1. *For any $n \geq 1$, there is an equivalence*

$$\rho_{\tilde{\mathfrak{X}}} : \text{LS}_r(X, \mathbb{B}_{\text{dR},n}^+)^{\text{H-small}} \simeq \text{MIC}_r(\tilde{X}_n)^{\text{H-small}}$$

between the moduli stack $\text{LS}_r(X, \mathbb{B}_{\text{dR},n}^+)^{\text{H-small}}$ of Hitchin-small $\mathbb{B}_{\text{dR},n}^+$ -local systems of rank r on $X_{\text{proét}}$ and the moduli stack $\text{MIC}_r(\tilde{X}_n)^{\text{H-small}}$ of Hitchin-small integrable connections of rank r on \tilde{X}_n .

Remark 1.2. The $n = 1$ case for above theorem was already obtained by Anschütz, Heuer and Le Bras [AHLB23b], where the functor $\rho_{\tilde{\mathfrak{X}}}$ (in the $n = 1$ case) is denoted by $S_{\tilde{\mathfrak{X}}}$. We remark the injectivity of $S_{\tilde{\mathfrak{X}}}$ (that is, the full faithfulness of its evaluation at every $\text{Spa}(A, A^+) \in \text{Perfd}$) was proved in [AHLB23a], and the surjectivity was established in [AHLB23b] by checking this locally and then gluing together. Compared with theirs, our construction provides an *explicit* and *global* description of $\rho_{\tilde{\mathfrak{X}}}^{-1}$.

Now, we explain some notions appearing in above Theorem. Let $S = \text{Spa}(A, A^+) \in \text{Perfd}$ be an affinoid perfectoid space. Let \mathfrak{Z} be a smooth formal scheme over $\text{Spf}(A^+)$ with a smooth lifting $\tilde{\mathfrak{Z}}$ over $\mathbf{A}_{\text{inf}}(A, A^+)$. Let Z be the generic fiber of \mathfrak{Z} with the lifting \tilde{Z} over \mathbf{B}_{dR}^+ induced by $\tilde{\mathfrak{Z}}$. For any $n \geq 1$, let \tilde{Z}_n be the reduction of \tilde{Z} modulo t^n and $\mathbb{B}_{\text{dR},n}^+ := \mathbb{B}_{\text{dR}}^+/t^n$.

By a $\mathbb{B}_{\text{dR},n}^+$ -local system of rank r on Z_v , the v -site of Z in the sense of [Sch17], we mean a locally finite free $\mathbb{B}_{\text{dR},n}^+$ -module of rank r . When $n = 1$, the $\mathbb{B}_{\text{dR},1}^+$ is the structure sheaf $\hat{\mathcal{O}}_Z$, and the $\mathbb{B}_{\text{dR},1}^+$ -local systems reduce to the v -vector bundles on Z_v . One can similarly define $\mathbb{B}_{\text{dR},n}^+$ -local systems of rank r on Z_v .

Let $\tilde{d} : \tilde{\mathfrak{Z}} \rightarrow \Omega_{\tilde{\mathfrak{Z}}}^1$ be the usual (p, ξ) -complete derivation on $\tilde{\mathfrak{Z}}$ and let d be the following composite

$$d : \mathcal{O}_{\tilde{\mathfrak{Z}}} \xrightarrow{\tilde{d}} \Omega_{\tilde{\mathfrak{Z}}}^1 \hookrightarrow \Omega_{\tilde{\mathfrak{Z}}}^1\{-1\}.$$

Here $\Omega_{\mathbb{Z}}^1\{-1\}$ denotes the Breuil–Kisin–Fargues twist of $\Omega_{\mathbb{Z}}^1$ (cf. §1.4). By an *integrable connection* of rank r on \tilde{Z}_n , we mean a pair (\mathcal{M}, ∇) consisting of a locally finite free $\mathcal{O}_{\tilde{Z}_n}$ -module \mathcal{M} together with a $\mathbb{B}_{\text{dR},n}^+(A, A^+)$ -linear map

$$\nabla : \mathcal{M} \rightarrow \mathcal{M} \otimes_{\mathcal{O}_{\tilde{Z}_n}} \Omega_{\mathbb{Z}}^1\{-1\}$$

satisfying the Leibniz rule with respect to d such that $\nabla \wedge \nabla = 0$. One can similarly define integrable connections of rank r on \tilde{Z} . We remark that if we trivialize the Tate twist $\Omega_{\tilde{Z}}^1(-1)$ by $\Omega_{\tilde{Z}}^1 \cdot t^{-1}$, then via the identification

$$\mathcal{M} \otimes_{\mathcal{O}_{\tilde{Z}_n}} \Omega_{\mathbb{Z}}^1\{-1\} = \mathcal{M} \otimes_{\mathcal{O}_{\tilde{Z}_n}} \Omega_{\mathbb{Z}}^1(-1),$$

locally on \tilde{Z}_n , ∇ behaves like a *t-connection* in the sense of [Yu24]. In particular, when $n = 1$, we see that $d = 0$ and the integrable connections on $\tilde{Z}_1 = Z$ reduce to the Higgs bundles on Z .

For our \mathfrak{X} with the fixed lifting $\tilde{\mathfrak{X}}$, for any $S = \text{Spa}(A, A^+) \in \text{Perfd}$, let \mathfrak{X}_A , $\tilde{\mathfrak{X}}_A$, X_A and \tilde{X}_A be the base-changes of \mathfrak{X} , $\tilde{\mathfrak{X}}$, X and \tilde{X} to A^+ , $\mathbb{A}_{\text{inf}}(A, A^+)$, A and $\mathbb{B}_{\text{dR}}^+(A, A^+)$, respectively. Consider the following two functors:

$$\text{LS}_r(X, \mathbb{B}_{\text{dR},n}^+) : \text{Perfd} \rightarrow \text{Groupoid}, \quad S \mapsto \{ \mathbb{B}_{\text{dR},n}^+\text{-local systems on } (X_A)_v \text{ of rank } r \}$$

and

$$\text{MIC}_r(\tilde{X}_n) : \text{Perfd} \rightarrow \text{Groupoid}, \quad S \mapsto \{ \text{integrable connections on } \tilde{X}_{A,n} \text{ of rank } r \}.$$

Remark 1.3. For $n = 1$, the above functors were first considered by Heuer [Heu22] without assume X has a good reduction. In *loc.cit.*, he introduced the functors $v\text{Bun}_r(X)$ (resp. $\text{HIG}_r(X)$) of v -vector bundles (resp. Higgs bundles) of rank r on $X_{\text{proét}}$ (resp. $X_{\text{ét}}$), which is exactly the $\text{LS}_r(X, \mathbb{B}_{\text{dR},1}^+)$ (resp. $\text{MIC}_r(\tilde{X}_1)$) above. He proved these two stacks are small v -stacks on Perfd and isomorphic after taking étale sheafifications. For general n , the above stacks are introduced by Yu [Yu24]. In *loc.cit.*, Yu also proved all stacks above are small v -stacks, and $\text{LS}_r(X, \mathbb{B}_{\text{dR},n}^+)$ is isomorphic to $\text{MIC}_r(\tilde{X}_n)$ after taking étale sheafifications.

Clearly, we have the following diagram:

(1.1)

$$\begin{array}{ccc}
 \begin{array}{c} \vdots \\ \downarrow \\ \text{LS}(X, \mathbb{B}_{\text{dR},n+1}^+) \\ \downarrow \\ \text{LS}(X, \mathbb{B}_{\text{dR},n}^+) \\ \downarrow \\ \vdots \\ \downarrow \\ v\text{Bun}_r(X) = \text{LS}(X, \mathbb{B}_{\text{dR},1}^+) \end{array} & & \begin{array}{c} \vdots \\ \downarrow \\ \text{MIC}_r(\tilde{X}_{n+1}) \\ \downarrow \\ \text{MIC}_r(\tilde{X}_n) \\ \downarrow \\ \vdots \\ \downarrow \\ \text{MIC}_r(\tilde{X}_1) = \text{HIG}_r(X) \end{array} \\
 & \searrow \tilde{h} & \swarrow h \\
 & \mathcal{A}_r &
 \end{array}$$

where all vertical maps are induced by taking the obvious reduction, h and \tilde{h} denote the Hitchin fibrations introduced in [Heu22], and \mathcal{A}_r is the Hitchin base, which is defined as a functor

$$\mathcal{A}_r : \text{Perfd} \rightarrow \text{Sets}, \quad \text{Spa}(A, A^+) \mapsto \oplus_{i=1}^r H^0(X_A, (\Omega_{X_A/A}^1\{-1\})^{\otimes r}).$$

One can define the *Hitchin-small locus* $\mathcal{A}_r^{\text{H-small}} \subset \mathcal{A}_r$ by

$$\mathcal{A}_r^{\text{H-small}} : \text{Perfd} \rightarrow \text{Sets}, \quad S = \text{Spa}(A, A^+) \mapsto \oplus_{i=1}^r p^{< \frac{i}{p-1}} H^0(\mathfrak{X}_A, (\Omega_{\mathfrak{X}_A/A^+}^1\{-1\})^{\otimes r})$$

as in [AHLB23b], where $p^{\leq \frac{i}{p-1}} := (\zeta_p - 1)^i \mathfrak{m}_C \subset \mathcal{O}_C$. The \mathcal{A}_r and $\mathcal{A}_r^{\text{H-small}}$ both make sense as v -sheaves. For any stacks Z lying over \mathcal{A}_r , define its Hitchin-small locus $Z^{\text{H-small}}$ as the sub-stack

$$Z^{\text{H-small}} := Z \times_{\mathcal{A}_r} \mathcal{A}_r^{\text{H-small}}.$$

Now, we have explained all notions involved in Theorem 1.1. It is still worth pointing out

Remark 1.4. A $\mathbb{B}_{\text{dR},n}^+$ -local system (resp. integrable connection on \tilde{X}_n) is Hitchin-small *if and only if* its reduction modulo t is a Hitchin-small v -bundle on $X_{\text{proét}}$ (resp. Higgs bundle on $X_{\text{ét}}$).

We list some immediate corollaries. Again, let \mathfrak{X} be a liftable smooth formal scheme over \mathcal{O}_C with a fixed lifting $\tilde{\mathfrak{X}}$ over \mathbf{A}_{inf} . First, by letting n go to ∞ , we obtain the following equivalence of stacks:

Corollary 1.5. *There exists an equivalence*

$$\rho_{\tilde{\mathfrak{X}}} : \text{LS}_r(X, \mathbb{B}_{\text{dR}}^+)^{\text{H-small}} \simeq \text{MIC}_r(\tilde{X})^{\text{H-small}}$$

between the moduli stack $\text{LS}_r(X, \mathbb{B}_{\text{dR}}^+)^{\text{H-small}}$ of Hitchin-small \mathbb{B}_{dR}^+ -local systems of rank r and the moduli stack $\text{MIC}_r(\tilde{X})^{\text{H-small}}$ of Hitchin-small integrable connections of rank r .

To simplify the notations, set $\mathbb{B}_{\text{dR}}^+ := \mathbb{B}_{\text{dR},\infty}^+$ and $\tilde{X}_{\infty} := \tilde{X}$. The next corollary follows from by taking C -points in Theorem 1.1 immediately.

Corollary 1.6. *For any $1 \leq n \leq \infty$, there exists an equivalence*

$$\text{LS}(X, \mathbb{B}_{\text{dR}}^+)^{\text{H-small}}(C) \simeq \text{MIC}(\tilde{X})^{\text{H-small}}(C)$$

between the category $\text{LS}(X, \mathbb{B}_{\text{dR}}^+)^{\text{H-small}}(C)$ of Hitchin-small $\mathbb{B}_{\text{dR},n}^+$ -local systems on X_v and the category $\text{MIC}(\tilde{X})^{\text{H-small}}(C)$ of Hitchin-small integrable connections on \tilde{X}_n .

This generalizes the equivalence between the categories of (Faltings-)small objects in [CLWZ24, Th.7.11]. See also Theorem 1.10 for a more explicit statement.

Corollary 1.7. *Let \mathbb{P}^d be the projective space over C of dimension d . For any $1 \leq n \leq \infty$, there exists an equivalence of the whole stacks*

$$\text{LS}_r(\mathbb{P}^d, \mathbb{B}_{\text{dR},n}^+) \simeq \text{MIC}_r(\tilde{\mathbb{P}}_n^d).$$

Note that \mathbb{P}^d always admits a liftable smooth formal model \mathfrak{P}^d over \mathcal{O}_C , and $\tilde{\mathbb{P}}_n^d$ is the lifting of \mathbb{P}^d coming from a lifting of \mathfrak{P}^d .

Proof. This holds true as any $\mathbb{B}_{\text{dR},n}^+$ -local systems (resp. integrable connections on $\tilde{\mathbb{P}}_n^d$) are automatically Hitchin-small (and even belonging to the fiber of the origin $0 \in \mathcal{A}_r$). See [AHLB23b, Cor. 3.26] for details. \square

Remark 1.8. In Corollary 1.7, it is not necessary to work with the smooth integral model of \mathbb{P}^n . More generally, for any smooth rigid variety X over C with a smooth lifting \tilde{X} over \mathbf{B}_{dR}^+ , using the period sheaf with connection $(\mathcal{O}_{\mathbb{B}_{\text{dR}}}, d)$ in [Yu24, Def. 2.36] and the same argument in this paper, we can give an equivalence of stacks

$$\text{LS}_r(X, \mathbb{B}_{\text{dR},n}^+)^0 \simeq \text{MIC}_r(\tilde{X}_n)^0$$

where Z^0 denotes the fiber at the origin $0 \in \mathcal{A}$ for any stack Z over \mathcal{A} . See [LQN24] for more details.

Remark 1.9. For general X , it is still a question if there exists an equivalence of the whole stacks

$$(1.2) \quad \text{LS}_r(X, \mathbb{B}_{\text{dR},n}^+) \simeq \text{MIC}_r(\tilde{X}_n).$$

Note that for $n = 1$; that is, in the case for v -vector bundles and Higgs bundles, it was proved by Heuer and Xu [HX24] that if X is a smooth curve, such an equivalence (1.2) exists up to a certain normalisation. Beyond curves, we only get an equivalence between the category of v -vector bundles on X_v and the category Higgs bundles on $X_{\text{ét}}$ for *proper* smooth X [Heu23]. It seems the problem (in the $n = 1$ case) might be solved using the Simpson gerbe claimed by Bhargav Bhatt and Mingjia Zhang. However, for general $n > 1$, we know *nothing* on (1.2) besides results in this paper!

1.3. Strategy for the proof and organization. Here, we sketch the idea for the proof of Theorem 1.1 and explain how the paper is organized.

Fix a smooth \mathfrak{X} with the fixed lifting $\tilde{\mathfrak{X}}$. For any $S = \mathrm{Spa}(A, A^+) \in \mathrm{Perfd}$, let \mathfrak{X}_A , $\tilde{\mathfrak{X}}_A$, X_A and \tilde{X}_A be the base-changes of \mathfrak{X} , $\tilde{\mathfrak{X}}$, X and \tilde{X} to A^+ , $\mathbb{A}_{\mathrm{inf}}(A, A^+)$, A and $\mathbb{B}_{\mathrm{dR}}^+(A, A^+)$, respectively. Recall that in order to get the stacky Simpson correspondence [AHLB23b, Th. 1.1], one have to constructed an equivalence of categories

$$S_{\tilde{\mathfrak{X}}_A} : \mathrm{HIG}_r^{\mathrm{H-small}}(X)(A) \xrightarrow{\sim} v\mathrm{Bun}_r(X, \hat{\mathcal{O}})^{\mathrm{H-small}}(A)$$

which is functorial in A . To define $S_{\tilde{\mathfrak{X}}_A}$, one have to use the period sheaf with Higgs field $(\mathcal{B}_{\tilde{X}_A}, \Theta)$ constructed in [AHLB23a] (depending on the lifting of \mathfrak{X} over $\mathbb{A}_{\mathrm{inf}}/\xi^2$ functorial in A and coming from prismatic theory), which locally behaves like a pd-polynomial ring over $\hat{\mathcal{O}}_X$ with the usual derivation on the coordinates.

The key observation is that if one could constructed a period sheaf with connection $(\mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}, A}^+, d)$ for each \mathfrak{X}_A functorial in A , of which the reduction modulo t behaves (at least locally) like $(\mathcal{B}_{\tilde{X}_A}, \Theta)$, then it seems possible to generalize the stacky small Simpson correspondence to a stacky small Riemann–Hilbert correspondence. The §2 is devoted do this. We also remark that our construction of $(\mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}, A}^+, d)$ is *self-contained*; that is, it is different with the construction in [AHLB23a], and in particular does not use any prismatic theory. It is indeed inspired of the constructions in [MW23, §2] and [CLWZ24, §3].

Using $(\mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}, A}^+, d)$, one can prove the following Riemann–Hilbert correspondence:

Theorem 1.10 (Theorem 4.1). *Fix a $\mathrm{Spa}(A, A^+) \in \mathrm{Perfd}$. Let $\nu : X_{A, v} \rightarrow X_{A, \mathrm{ét}}$ be the natural morphism of sites. Let $1 \leq n \leq \infty$.*

- (1) *For any $\mathbb{L} \in \mathrm{LS}^{\mathrm{H-small}}(X, \mathbb{B}_{\mathrm{dR}, n}^+)(A)$ of rank r , we have*

$$R^n \nu_* (\mathbb{L} \otimes_{\mathbb{B}_{\mathrm{dR}}^+} \mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}, A}^+) = \begin{cases} \mathcal{D}(\mathbb{L}), & n = 0 \\ 0, & n \geq 1 \end{cases}$$

where $\mathcal{D}(\mathbb{L})$ is a locally finite free $(\mathcal{O}_{\tilde{X}_{A, n}})$ -module of rank r on $X_{A, \mathrm{ét}}$ such that the

$$\mathrm{id}_{\mathbb{L}} \otimes d : \mathbb{L} \otimes_{\mathbb{B}_{\mathrm{dR}}^+} \mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}, A}^+ \rightarrow \mathbb{L} \otimes_{\mathbb{B}_{\mathrm{dR}}^+} \mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}, A}^+ \otimes_{\mathcal{O}_{\tilde{\mathfrak{X}}_A}} \Omega_{\tilde{\mathfrak{X}}_A}^1 \{-1\}$$

induces a flat connection $\nabla_{\mathbb{L}}$ on $\mathcal{D}(\mathbb{L})$ making $(\mathcal{D}(\mathbb{L}), \nabla_{\mathbb{L}})$ an object in $\mathrm{MIC}^{\mathrm{H-small}}(\tilde{X}_n)(A)$.

- (2) *For any $(\mathcal{D}, \nabla) \in \mathrm{MIC}^{\mathrm{H-small}}(\tilde{X}_n)(A)$ of rank r , define*

$$\nabla_{\mathcal{D}} := \nabla \otimes \mathrm{id} + \mathrm{id} \otimes d : \mathcal{D} \otimes_{\mathcal{O}_{\tilde{\mathfrak{X}}_A}} \mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}, A}^+ \rightarrow \mathcal{D} \otimes_{\mathcal{O}_{\tilde{\mathfrak{X}}_A}} \mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}, A}^+ \otimes_{\mathcal{O}_{\tilde{\mathfrak{X}}_A}} \Omega_{\tilde{\mathfrak{X}}_A}^1 \{-1\}.$$

Then

$$\mathbb{L}(\mathcal{D}, \nabla) := (\mathcal{D} \otimes_{\mathcal{O}_{\tilde{\mathfrak{X}}_A}} \mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}, A}^+)^{\nabla_{\mathcal{D}}=0}$$

is an object of rank r in $\mathrm{LS}^{\mathrm{H-small}}(X, \mathbb{B}_{\mathrm{dR}, n}^+)(A)$.

- (3) *The functors $\mathbb{L} \mapsto (\mathcal{D}(\mathbb{L}), \nabla_{\mathbb{L}})$ and $(\mathcal{D}, \nabla) \mapsto \mathbb{L}(\mathcal{D}, \nabla)$ in Items (1) and (2) respectively define an equivalence of categories*

$$\rho_{\tilde{\mathfrak{X}}_A} : \mathrm{LS}^{\mathrm{H-small}}(X, \mathbb{B}_{\mathrm{dR}, n}^+)(A) \xrightarrow{\sim} \mathrm{MIC}^{\mathrm{H-small}}(\tilde{X}_n)(A)$$

which preserves ranks, tensor products and dualities. Moreover, for any $\mathbb{L} \in \mathrm{LS}^{\mathrm{H-small}}(X, \mathbb{B}_{\mathrm{dR}, n}^+)(A)$ with corresponding $(\mathcal{D}, \nabla) \in \mathrm{MIC}^{\mathrm{H-small}}(\tilde{X}_n)(A)$, there exists a quasi-isomorphism

$$R\nu_* \mathbb{L} \simeq \mathrm{DR}(\mathcal{D}, \nabla).$$

In particular, we have a quasi-isomorphism

$$\mathrm{R}\Gamma(X_{A, v}, \mathbb{L}) \simeq \mathrm{R}\Gamma(X_{A, \mathrm{ét}}, \mathrm{DR}(\mathcal{D}, \nabla)).$$

It is clearly the construction of $\rho_{\tilde{\mathfrak{X}}_A}$ is functorial in A and thus Theorem 1.1 follows immediately. The §4 is devoted to proving 1.10. The approach is standard: We reduce the proof to the case where \mathfrak{X} admits a chart and then check everything locally. We shall do the latter in §3.

Remark 1.11. The construction of $(\mathcal{O}\mathbb{B}_{\mathrm{dR},\mathrm{pd}}^+, \mathrm{d})$ is no doubt under the influence of the constructions in [CLWZ24, §3]. All results in this paper should be viewed as a continuation of the Riemann–Hilbert correspondence (for *Faltings-small* objects) [CLWZ24, Th. 7.11]. It is worth pointing out Theorem 1.10 is also compatible with [CLWZ24, Th. 7.11] as we can compare period sheaves used here and in *loc.cit.* (cf. Remark 2.11).

1.4. Notations. Throughout this paper, let C be a algebraically closed perfectoid field containing \mathbb{Q}_p with the ring of integers \mathcal{O}_C . Let $\mathbf{A}_{\mathrm{inf}}$ and $\mathbf{B}_{\mathrm{dR}}^+$ be the corresponding infinitesimal and de Rham period ring. Fix an embedding $p^\mathbb{Q} \subset C^\times$, which induces an embedding $\varpi^\mathbb{Q} \subset C^{\flat^\times}$, where $\varpi = (p, p^{1/p}, p^{1/p^2}, \dots) \in C^\flat$. Fix a coherent system $\{\zeta_{p^n}\}_{n \geq 0}$ of primitive p^n -th roots of unity in C , and let $\epsilon := (1, \zeta_p, \zeta_p^2, \dots) \in C^\flat$. Let $u = [\epsilon^{\frac{1}{p}}] - 1 \in \mathbf{A}_{\mathrm{inf}}$, and then the canonical surjection $\theta : \mathbf{A}_{\mathrm{inf}} \rightarrow \mathcal{O}_C$ is principally generated by $\xi := \frac{\phi(u)}{u}$. Let $t = \log([\epsilon]) \in \mathbf{B}_{\mathrm{dR}}^+$ be the Fontaine’s p -adic analogue of “ $2\pi i$ ”. For a (sheaf of) $\mathbf{A}_{\mathrm{inf}}$ -module M and any $n \in \mathbb{Z}$, denote by $M\{n\}$ its Breuil–Kisin–Fargues twist

$$M\{n\} := M \otimes_{\mathbf{A}_{\mathrm{inf}}} \mathrm{Ker}(\theta)^{\otimes n},$$

which can be trivialized by ξ^n ; that is, we have the identification $M\{n\} = M \cdots \xi^n$. Using this, we may regard M as a sub- $\mathbf{A}_{\mathrm{inf}}$ -module of $M\{-1\}$ via the identification $M = \xi M\{-1\}$.

Fix a ring R . If an element $x \in R$ admits arbitrary pd-powers, we denote by $x^{[n]}$ its n -th pd-power (i.e. analogue of $\frac{x^n}{n!}$) in R . Put $E_i = (0, \dots, 1, \dots, 0) \in \mathbb{N}^d$ with 1 appearing at exactly the i -th component. For any $J = (j_1, \dots, j_d) \in \mathbb{N}^d$ and any $x_1, \dots, x_d \in R$, we put

$$\underline{x}^J := x_1^{j_1} \cdots x_d^{j_d}$$

and if moreover x_i admits arbitrary pd-powers in A for all i , we put

$$\underline{x}^{[J]} := x_1^{[j_1]} \cdots x_d^{[j_d]}.$$

For any $\alpha \in \mathbb{N}[1/p] \cap [0, 1)$, we put

$$\zeta^\alpha = \zeta_{p^n}^m$$

if $\alpha = \frac{m}{p^n}$ with p, m are coprime in \mathbb{N} . If $x \in A$ admits compatible p^n -th roots $x^{\frac{1}{p^n}}$, we put

$$x^\alpha = x^{\frac{m}{p^n}}$$

for $\alpha = \frac{m}{p^n}$ as above. In general, for any $\underline{\alpha} := (\alpha_1, \dots, \alpha_d) \in (\mathbb{N}[1/p] \cap [0, 1))^d$ and any x_1, \dots, x_d admitting compatible p^n -th roots, we put

$$\underline{x}^{\underline{\alpha}} := x_1^{\alpha_1} \cdots x_d^{\alpha_d}.$$

Acknowledgement. The work started as a research project of 2024 summer school on algebra and number theory at Peking university, held by School of Mathematical Science, Peking University and Academy of Mathematics and System Science, Chinese Academy of Science. The authors would like to thank the organizers and institutes for providing with a great environment for our collaborating. We would like to thank Jiahong Yu for explaining his paper [Yu24] and thank Yue Chen, Chenglong Ma and Wenrong Zou for their interests in this project. The idea for constructing $(\mathcal{O}\mathbb{B}_{\mathrm{dR},\mathrm{pd}}^+, \mathrm{d})$ occurs during the period of the last author’s collaboration [CLWZ24] with Zekun Chen, Ruochuan Liu and Xinwen Zhu, and we would like to thank them for allowing us to use some results in *loc.cit.* and exhibit Theorem 1.1 here.

Y.W. is partially supported by CAS Project for Young Scientists in Basic Research, Grant No. YSBR-032.

2. THE PERIOD SHEAF WITH CONNECTION $(\mathcal{O}\mathbb{B}_{\mathrm{dR},\mathrm{pd}}^+, \mathrm{d})$

Fix a $\mathrm{Spa}(A, A^+) \in \mathrm{Perfd}$ and put $\mathbf{A}_{\mathrm{inf}} := \mathbb{A}_{\mathrm{inf}}(A, A^+)$ and $\mathbf{B}_{\mathrm{dR}}^+ := \mathbb{B}_{\mathrm{dR}}^+(A, A^+)$ for short in what follows. Let \mathfrak{X} be a smooth formal scheme over A^+ with the generic fiber X . We say \mathfrak{X} is *liftable* if it admits a smooth lifting $\tilde{\mathfrak{X}}$ over $\mathbf{A}_{\mathrm{inf}}$ along the canonical map $\theta : \mathbf{A}_{\mathrm{inf}} \rightarrow A^+$. From now on, we assume \mathfrak{X} is liftable and fix such a lifting $\tilde{\mathfrak{X}}$ and a fortiori a lifting \tilde{X} of X over $\mathbf{B}_{\mathrm{dR}}^+$.

For any affinoid perfectoid $U = \mathrm{Spa}(S, S^+) \in X_v$, let Σ_U be the set of all étale morphisms

$$i_{\mathfrak{Y}} : \mathfrak{Y} = \mathrm{Spf}(\mathcal{R}) \rightarrow \mathfrak{X}$$

over A^+ such that the natural map $U \rightarrow X$ factors through the generic fiber of $i_{\mathfrak{Y}}$. By the étaleness of $i_{\mathfrak{Y}}$, the \mathfrak{Y} admits a unique lifting $\tilde{\mathfrak{Y}} = \mathrm{Spf}(\tilde{\mathcal{R}})$ over $\tilde{\mathfrak{X}}$ such that $i_{\mathfrak{Y}}$ uniquely lifts to a morphism

$$i_{\tilde{\mathfrak{Y}}} : \tilde{\mathfrak{Y}} = \mathrm{Spf}(\tilde{\mathcal{R}}) \rightarrow \tilde{\mathfrak{X}}$$

over $\mathbf{A}_{\mathrm{inf}}$. Then we have a natural morphism

$$\theta_{\mathfrak{Y}} : \tilde{\mathcal{R}} \hat{\otimes}_{\mathbf{A}_{\mathrm{inf}}} \mathbf{A}_{\mathrm{inf}}(U) \rightarrow S^+$$

lifting the natural map $\mathcal{R} \hat{\otimes}_{A^+} S^+ \rightarrow S^+$ along θ .

Lemma 2.1. *The kernel $\mathrm{Ker}(\theta_{\mathfrak{Y}})$ of $\theta_{\mathfrak{Y}}$ is a finite generated ideal of $\tilde{\mathcal{R}} \hat{\otimes}_{\mathbf{A}_{\mathrm{inf}}} \mathbf{A}_{\mathrm{inf}}(U)$.*

Proof. Similar to the proof of [CLWZ24, Lem. 2.24]. \square

Let $A_{\mathfrak{Y}}$ be the $(p, \mathrm{Ker}(\theta_{\mathfrak{Y}}))$ -adic completion of $\tilde{\mathcal{R}} \hat{\otimes}_{\mathbf{A}_{\mathrm{inf}}} \mathbf{A}_{\mathrm{inf}}(U)$. Denote by $B_{\mathfrak{Y}} = A_{\mathfrak{Y}}[u \cdot \mathrm{Ker}(\theta_{\mathfrak{Y}})\{-1\}] \subset A_{\mathfrak{Y}}[\frac{1}{\xi}]$ the blow-up algebra

$$B_{\mathfrak{Y}} := A_{\mathfrak{Y}}\left[\frac{u \cdot \mathrm{Ker}(\theta_{\mathfrak{Y}})}{\xi}\right]$$

and define

$$\Gamma_{\mathfrak{Y}} := (B_{\mathfrak{Y}}[u \cdot \mathrm{Ker}(\theta_{\mathfrak{Y}})\{-1\}]_{\mathrm{pd}})^{\wedge} = (B_{\mathfrak{Y}}[\frac{u \cdot \mathrm{Ker}(\theta_{\mathfrak{Y}})}{\xi}]_{\mathrm{pd}})^{\wedge}$$

the $(p, \mathrm{Ker}(\theta_{\mathfrak{Y}}))$ -adic completion of the pd-envelope of $B_{\mathfrak{Y}}$ with respect to the ideal generated by $\frac{u \cdot \mathrm{Ker}(\theta_{\mathfrak{Y}})}{\xi}$. Clearly, it is an algebra over $\mathbf{A}_{\mathrm{inf}}(U)[u]_{\mathrm{pd}}^{\wedge}$, the (p, ξ) -adic completion of the pd-envelope of $\mathbf{A}_{\mathrm{inf}}(U)$ with respect to the ideal (u) .

Before we move on, let us give a quick review of the relative Robba rings $\tilde{\mathbb{C}}^1(U)$. Let

$$\mathrm{Spa}(\tilde{\mathbb{C}}^1(U), \tilde{\mathbb{C}}^{1,+}(U)) \rightarrow \mathrm{Spa}(\mathbf{A}_{\mathrm{inf}}(U)[\frac{1}{p}], \mathbf{A}_{\mathrm{inf}}(U))$$

be the rational localisation with respect to the condition $||[\varpi]|| = p^{-1} = |p|$. Let $|\cdot|_1$ be the norm on $\mathbf{A}_{\mathrm{inf}}(U)[\frac{1}{p[\varpi]}]$ such that for any $x = \sum_{n \geq 0} [x_n]p^n$ with $x_n \in S^{b+}$,

$$|x|_1 := \sup_{n \geq 0} \|x_n\| p^{-n},$$

where $\|\cdot\|$ denotes the spectral norm on S^b . Then we have the following result:

Lemma 2.2. (1) *The $\tilde{\mathbb{C}}^{1,+}(U) = \mathbf{A}_{\mathrm{inf}}(U)[(\frac{[\varpi]}{p})^{\pm 1}]^{\wedge}$ is the p -adic completion of $\mathbf{A}_{\mathrm{inf}}(U)[(\frac{[\varpi]}{p})^{\pm 1}]$ and $\tilde{\mathbb{C}}^1(U) = \tilde{\mathbb{C}}^{1,+}(U)[\frac{1}{p}]$.*

(2) *The $\tilde{\mathbb{C}}^1(U)$ is the completion of $\mathbf{A}_{\mathrm{inf}}(U)[\frac{1}{p[\varpi]}]$ with respect to the norm $|\cdot|_1$, which is thus a Banach \mathbb{Q}_p -algebra with the closed unit ball*

$$\tilde{\mathbb{C}}^{1,+}(U) = \{x \in \tilde{\mathbb{C}}^1(U) \mid |x|_1 \leq 1\}.$$

(3) *The $t = \log[\epsilon] = \sum_{n \geq 1} \frac{(1-\epsilon)^n}{n}$ converges in $\tilde{\mathbb{C}}^{1,+}(U)$ and is a unit-multiple of $u\xi$.*

(4) *The u is a unit-multiple of $[\epsilon^{1/p} - 1]$ in $\tilde{\mathbb{C}}^{1,+}(U)$, and thus admits arbitrary pd-powers in $\tilde{\mathbb{C}}^{1,+}(U)$.*

(5) *The natural map $\theta : \mathbf{A}_{\mathrm{inf}}(U) \rightarrow S^+$ extends to a surjection $\tilde{\mathbb{C}}^{1,+}(U) \rightarrow S^+$ with kernel principally generated by $\frac{\xi}{p}$, which induces a canonical bounded morphism*

$$\tilde{\mathbb{C}}^1(U) \rightarrow \mathbb{B}_{\mathrm{dR}}^+(U)$$

identifying the target with the ξ -adic completion of the source.

Proof. Item (1) follows from the definition. Item (2) is [CLWZ24, Lem. 2.7(1)]. Item (3) and the first part of Item (4) is [CLWZ24, Prop. 3.12]. For the second part of Item (4), just note that for any $n \geq 1$,

$$|\frac{u^n}{n!}|_1 = |[\frac{\epsilon^{1/p} - 1}{n!}]^n|_1 = p^{\nu_p(n!) - \frac{n}{p-1}} \leq p^{-\frac{1}{p-1}}.$$

Here, we use the well-known fact that $\nu_p(n!) \geq \frac{n-1}{p-1}$ for any $n \geq 1$. All statements in Item (5) follows from [CLWZ24, Lem. 2.12], except that

$$\text{Ker}(\tilde{\mathbb{C}}^{1,+}(U) \rightarrow S^+) = \left(\frac{\xi}{p}\right).$$

To see this, note that ξ is a unit-multiple of $\beta := p - [\varpi]$ in $\mathbb{A}_{\text{inf}}(U)$. It suffices to show that

$$\text{Ker}(\tilde{\mathbb{C}}^{1,+}(U) \rightarrow S^+) = \left(1 - \frac{[\varpi]}{p}\right).$$

This follows from Item (1) easily. \square

By Lemma 2.2(4), we see the natural inclusion $\mathbb{A}_{\text{inf}}(U) \rightarrow \tilde{\mathbb{C}}^{1,+}(U)$ uniquely extends to the inclusion

$$\mathbb{A}_{\text{inf}}(U)[u]_{\text{pd}}^\wedge \rightarrow \tilde{\mathbb{C}}^{1,+}(U).$$

Now, for any $? \in \{+, \emptyset\}$, we put

$$C_{\mathfrak{Y}}^? := \Gamma_{\mathfrak{Y}} \hat{\otimes}_{\mathbb{A}_{\text{inf}}(U)[u]_{\text{pd}}^\wedge} \tilde{\mathbb{C}}^{1,?}(U),$$

and then we have $C_{\mathfrak{Y}} = C_{\mathfrak{Y}}^+[\frac{1}{p}]$. Define

$$B_{\text{dR}, \text{pd}, \mathfrak{Y}}^+ := \xi\text{-adic completion of } C_{\mathfrak{Y}}.$$

Then $B_{\text{dR}, \text{pd}, \mathfrak{Y}}$ is a $\mathbb{B}_{\text{dR}}^+(U)$ -algebra by Lemma 2.2(5).

Let $d : \tilde{\mathcal{R}} \rightarrow \Omega_{\tilde{\mathcal{R}}}^1$ be the usual derivation on $\tilde{\mathcal{R}}$, where $\Omega_{\tilde{\mathcal{R}}}^1$ denotes the module of (p, ξ) -adically continuous differentials over $\tilde{\mathcal{R}}$ over \mathbf{A}_{inf} . It extends uniquely to an $\mathbb{A}_{\text{inf}}(U)$ -linear derivation

$$d : \tilde{\mathcal{R}} \hat{\otimes}_{\mathbf{A}_{\text{inf}}} \mathbb{A}_{\text{inf}}(U) \rightarrow \Omega_{\tilde{\mathcal{R}}}^1 \hat{\otimes}_{\mathbf{A}_{\text{inf}}} \mathbb{A}_{\text{inf}}(U)$$

such that for any $n \geq 1$,

$$d(\text{Ker}(\theta_{\mathfrak{Y}})^n) \subset \text{Ker}(\theta_{\mathfrak{Y}})^{n-1}.$$

Thus it extends uniquely to an $\mathbb{A}_{\text{inf}}(U)$ -linear derivation

$$d : A_{\mathfrak{Y}} \rightarrow A_{\mathfrak{Y}} \otimes_{\tilde{\mathcal{R}}} \Omega_{\tilde{\mathcal{R}}}^1,$$

and then an $\mathbb{A}_{\text{inf}}(U)$ -linear derivation

$$d : B_{\mathfrak{Y}} \rightarrow B_{\mathfrak{Y}} \otimes_{\tilde{\mathcal{R}}} \Omega_{\tilde{\mathcal{R}}}^1\{-1\}$$

where $\Omega_{\tilde{\mathcal{R}}}^1\{-1\} = \Omega_{\tilde{\mathcal{R}}}^1 \cdot \xi^{-1}$. Similar to the above, it extends uniquely to an $\mathbb{A}_{\text{inf}}(U)[u]_{\text{pd}}^\wedge$ -linear derivation

$$d : \Gamma_{\mathfrak{Y}} \rightarrow u\Gamma_{\mathfrak{Y}} \otimes_{\tilde{\mathcal{R}}} \Omega_{\tilde{\mathcal{R}}}^1\{-1\}.$$

By base-change to $\tilde{\mathbb{C}}^{1,+}(U)$ and taking ξ -adic completion, it uniquely extends to a $\tilde{\mathbb{C}}^{1,+}(U)$ -linear (resp. $\tilde{\mathbb{C}}^1(U)$ -linear, $\mathbb{B}_{\text{dR}}^+(U)$ -linear) derivation

$$d : C_{\mathfrak{Y}}^+ \rightarrow uC_{\mathfrak{Y}}^+ \otimes_{\tilde{\mathcal{R}}} \Omega_{\tilde{\mathcal{R}}}^1\{-1\} \text{ (resp. } d : C_{\mathfrak{Y}} \rightarrow C_{\mathfrak{Y}} \otimes_{\tilde{\mathcal{R}}} \Omega_{\tilde{\mathcal{R}}}^1\{-1\}, d : B_{\text{dR}, \text{pd}, \mathfrak{Y}}^+ \rightarrow B_{\text{dR}, \text{pd}, \mathfrak{Y}}^+ \otimes_{\tilde{\mathcal{R}}} \Omega_{\tilde{\mathcal{R}}}^1\{-1\}).$$

Recall that u is invertible in $\tilde{\mathbb{C}}^1(U)$ and $\mathbb{B}_{\text{dR}}^+(U)$.

Definition 2.3. Let $\mathcal{O}\tilde{\mathbb{C}}_{\text{pd}}^+$, $\mathcal{O}\tilde{\mathbb{C}}_{\text{pd}}$ and $\mathcal{O}\mathbb{B}_{\text{dR}, \text{pd}}^+$ be the sheaves corresponding to the sheafification of the presheaves sending each affinoid perfectoid $U \in X_{\text{proét}}$ to

$$\text{colim}_{\mathfrak{Y} \in \Sigma_U} C_{\mathfrak{Y}}^+, \text{ colim}_{\mathfrak{Y} \in \Sigma_U} C_{\mathfrak{Y}}, \text{ and } \text{colim}_{\mathfrak{Y} \in \Sigma_U} B_{\text{dR}, \text{pd}, \mathfrak{Y}}^+$$

respectively. For any $\mathcal{P} \in \{\mathcal{O}\tilde{\mathbb{C}}_{\text{pd}}^+, \mathcal{O}\tilde{\mathbb{C}}_{\text{pd}}, \mathcal{O}\mathbb{B}_{\text{dR}, \text{pd}}^+\}$, let

$$d : \mathcal{P} \rightarrow \mathcal{P} \otimes_{\mathcal{O}_{\tilde{\mathfrak{X}}}} \Omega_{\tilde{\mathfrak{X}}}^1\{-1\}$$

be the derivation inducing from the usual (continuous) derivation on $\tilde{\mathfrak{X}}$ as constructed above. Denote by $\text{DR}(\mathcal{P}, d)$ the corresponding de Rham complex

$$\mathcal{P} \xrightarrow{d} u\mathcal{P} \otimes_{\mathcal{O}_{\tilde{\mathfrak{X}}}} \Omega_{\tilde{\mathfrak{X}}}^1\{-1\} \xrightarrow{d} u^2\mathcal{P} \otimes_{\mathcal{O}_{\tilde{\mathfrak{X}}}} \Omega_{\tilde{\mathfrak{X}}}^2\{-2\} \rightarrow \cdots \xrightarrow{d} u^d\mathcal{P} \otimes_{\mathcal{O}_{\tilde{\mathfrak{X}}}} \Omega_{\tilde{\mathfrak{X}}}^d\{-d\}.$$

We remark that u is invertible for $\mathcal{P} \in \{\mathcal{O}\tilde{\mathbb{C}}_{\text{pd}}, \mathcal{O}\mathbb{B}_{\text{dR}, \text{pd}}^+\}$.

Then we have the following Poincaré's Lemma:

Theorem 2.4 (Poincaré's Lemma). *For any $(\mathbb{P}, \mathcal{P}) \in \{(\tilde{\mathbb{C}}^{1,+}, \mathcal{O}\tilde{\mathbb{C}}_{\text{pd}}^+), (\tilde{\mathbb{C}}^1, \mathcal{O}\tilde{\mathbb{C}}_{\text{pd}}), (\mathbb{B}_{\text{dR}}^+, \mathcal{O}\mathbb{B}_{\text{dR,pd}}^+)\}$, in the case $p \geq 3$, the natural morphism $\mathbb{P} \rightarrow \mathcal{P}$ induces an exact sequence*

$$0 \rightarrow \mathbb{P} \rightarrow \text{DR}(\mathcal{P}, d).$$

If $(\mathbb{P}, \mathcal{P}) = (\mathbb{B}_{\text{dR}}^+, \mathcal{O}\mathbb{B}_{\text{dR,pd}}^+)$, the above sequence is still exact for $p = 2$.

We shall prove the above theorem later by giving the local description of $\mathcal{P} \in \{\mathcal{O}\tilde{\mathbb{C}}_{\text{pd}}^+, \mathcal{O}\tilde{\mathbb{C}}_{\text{pd}}, \mathcal{O}\mathbb{B}_{\text{dR,pd}}^+\}$.

Suppose that $\mathfrak{X} = \text{Spf}(\mathcal{R})$ is *small affine*; that is, there is a (p -completely) étale morphism of formal schemes over A^+

$$\psi : \mathfrak{X} = \text{Spf}(\mathcal{R}) \rightarrow \text{Spf}(A^+ \langle T_1^{\pm 1}, \dots, T_d^{\pm 1} \rangle) = \text{Spf}(A^+ \langle \underline{T}^{\pm 1} \rangle).$$

Then ψ lifts uniquely to a $((p, \xi)$ -completely) étale map

$$\tilde{\psi} : \tilde{\mathfrak{X}} = \text{Spf}(\tilde{\mathcal{R}}) \rightarrow \mathbf{A}_{\text{inf}} \langle \underline{T}^{\pm 1} \rangle.$$

Let $X_{\infty} := \text{Spa}(\widehat{R}_{\infty}, \widehat{R}_{\infty}^+)$ be the base-change of $X = \text{Spa}(R = \mathcal{R}_{[1/p]}, R^+ = \mathcal{R}) \rightarrow \text{Spa}(A \langle \underline{T}^{\pm 1} \rangle, A^+ \langle \underline{T}^{\pm 1} \rangle)$ along the pro-étale Γ -torsor

$$\text{Spa}(A \langle \underline{T}^{\pm 1/p^{\infty}} \rangle, A^+ \langle \underline{T}^{\pm 1/p^{\infty}} \rangle) \rightarrow \text{Spa}(A \langle \underline{T} \rangle, A^+ \langle \underline{T} \rangle),$$

where $\Gamma = \bigoplus_{i=1}^d \mathbb{Z}_p \gamma_i$ acts on $A \langle \underline{T}^{1/p^{\infty}} \rangle$ via

$$\gamma_i(T_j^{\frac{1}{p^n}}) = \zeta_{p^n}^{\delta_{ij}} T_j^{\frac{1}{p^n}}, \quad \forall 1 \leq i, j \leq d \text{ \& } n \geq 0.$$

Put $T_i^{\flat} := (T_i, T_i^{1/p}, \dots) \in \widehat{R}_{\infty}^+$. Then for any $1 \leq i, j \leq d$ and any $n \geq 0$, we have

$$\gamma_i((T_j^{\flat})^{\frac{1}{p^n}}) = \epsilon^{\delta_{ij} \frac{1}{p^n}} (T_j^{\flat})^{\frac{1}{p^n}}.$$

Lemma 2.5. *For any affinoid perfectoid $U = \text{Spa}(S, S^+) \in X_{\infty, v}$ and any $\mathfrak{Y} \in \Sigma_U$, the map*

$$\iota : \mathbf{A}_{\text{inf}}(U)[[V_1, \dots, V_d]] \rightarrow A_{\mathfrak{Y}}$$

sending each V_i to $\frac{T_i - [T_i^{\flat}]}{T_i}$ is well-defined and an isomorphism of $\mathbf{A}_{\text{inf}}(U)$ -algebras. Here, $\mathbf{A}_{\text{inf}}(U)[[V_1, \dots, V_d]]$ is the ring of formal power series over $\mathbf{A}_{\text{inf}}(U)$ freely generated by V_1, \dots, V_d . Moreover, the isomorphism carries the ideal (ξ, V_1, \dots, V_d) onto the ideal $\text{Ker}(\theta_{\mathfrak{Y}})$.

Proof. Without loss of generality, we may assume $\mathfrak{Y} = \mathfrak{X} = \text{Spf}(\mathcal{R})$. To see ι is well-defined, consider the map

$$i : \mathbf{A}_{\text{inf}}(U)[V_1, \dots, V_d] \rightarrow A_{\mathfrak{Y}}$$

sending V_i to $\frac{T_i - [T_i^{\flat}]}{T_i}$. As $i(V_j) \in \text{Ker}(\theta_{\mathfrak{Y}})$ and $A_{\mathfrak{Y}}$ is $\text{Ker}(\theta_{\mathfrak{Y}})$ -complete, the i uniquely extends to a map of $\mathbf{A}_{\text{inf}}(U)$ -algebras

$$i : \mathbf{A}_{\text{inf}}(U)[[V_1, \dots, V_d]] \rightarrow A_{\mathfrak{Y}},$$

which is nothing but ι . To conclude, it remains to construct a map

$$j : A_{\mathfrak{Y}} \rightarrow \mathbf{A}_{\text{inf}}(U)[[V_1, \dots, V_d]]$$

which is the inverse of ι .

Consider the morphism of \mathbf{A}_{inf} -algebras

$$j : \mathbf{A}_{\text{inf}} \langle \underline{T}^{\pm 1} \rangle \rightarrow \mathbf{A}_{\text{inf}}(U)[[V_1, \dots, V_d]]$$

sending each T_i to $[T_i^{\flat}](1 - V_i)^{-1}$. Then the composite

$$\mathbf{A}_{\text{inf}} \langle \underline{T}^{\pm 1} \rangle \xrightarrow{j} \mathbf{A}_{\text{inf}}(U)[[V_1, \dots, V_d]] \xrightarrow{\text{mod } (\xi, V_1, \dots, V_d)} S^+$$

coincides with the natural morphism $A^+ \langle \underline{T}^{\pm 1} \rangle \rightarrow \mathcal{R} \rightarrow S^+$. By the étaleness of $\tilde{\psi}$, the j uniquely lifts to a morphism of \mathbf{A}_{inf} -algebras

$$\tilde{\mathcal{R}} \rightarrow \mathbf{A}_{\text{inf}}(U)[[V_1, \dots, V_d]]$$

and thus by scalar extension, uniquely extends to a morphism of $\mathbf{A}_{\text{inf}}(U)$ -algebras (still denoted by)

$$j : \tilde{\mathcal{R}} \widehat{\otimes}_{\mathbf{A}_{\text{inf}}} \mathbf{A}_{\text{inf}}(U) \rightarrow \mathbf{A}_{\text{inf}}(U)[[V_1, \dots, V_d]]$$

such that the composite

$$\tilde{\mathcal{R}} \hat{\otimes}_{\mathbf{A}_{\text{inf}}} \mathbf{A}_{\text{inf}}(U) \xrightarrow{j} \mathbf{A}_{\text{inf}}(U)[[V_1, \dots, V_d]] \xrightarrow{\text{mod } (\xi, V_1, \dots, V_d)} S^+$$

coincides with the composite

$$\tilde{\mathcal{R}} \hat{\otimes}_{\mathbf{A}_{\text{inf}}} \mathbf{A}_{\text{inf}}(U) \xrightarrow{\text{mod } \xi} \mathcal{R} \hat{\otimes}_{A^+} S^+ \rightarrow S^+,$$

which is nothing but $\theta_{\mathfrak{y}}$. In particular, j carries $\text{Ker}(\theta_{\mathfrak{y}})$ into the ideal (ξ, V_1, \dots, V_d) . Thus, as $\mathbf{A}_{\text{inf}}(U)[[V_1, \dots, V_d]]$ is $(p, \xi, V_1, \dots, V_d)$ -adically complete, j uniquely extends to a morphism of $\mathbf{A}_{\text{inf}}(U)$ -algebras (still denoted by)

$$j : A_{\mathfrak{y}} \rightarrow \mathbf{A}_{\text{inf}}(U)[[V_1, \dots, V_d]][u, \frac{uV_1}{\xi}, \dots, \frac{uV_d}{\xi}]_{\text{pd}}^{\wedge}.$$

To complete the proof, we only need to check j is the inverse of ι . Granting this, it follows from the above construction that

$$\iota((\xi, V_1, \dots, V_d)) \subset \text{Ker}(\theta_{\mathfrak{y}}) \text{ and } j(\text{Ker}(\theta_{\mathfrak{y}})) \subset (\xi, V_1, \dots, V_d),$$

yielding the “moreover” part of the result.

It is clear from the construction that $j \circ \iota = \text{id}$. We have to show $\iota \circ j = \text{id}$. Note that by construction, the composite

$$\mathbf{A}_{\text{inf}}\langle \underline{T}^{\pm 1} \rangle \xrightarrow{i \circ j} A_{\mathfrak{y}} \xrightarrow{\text{mod } \text{Ker}(\theta_{\mathfrak{y}})} S^+$$

coincides with natural map $A^+\langle \underline{T}^{\pm 1} \rangle \rightarrow \mathcal{R} \rightarrow S^+$, and thus by the étaleness of $\tilde{\psi}$, the $i \circ j$ unique extends to a morphism of \mathbf{A}_{inf} -algebras

$$\tilde{\mathcal{R}} \rightarrow A_{\mathfrak{y}}.$$

However, as $i \circ j$ coincides with natural map

$$\mathbf{A}_{\text{inf}}\langle \underline{T}^{\pm 1} \rangle \rightarrow \tilde{\mathcal{R}} \rightarrow \tilde{\mathcal{R}} \hat{\otimes}_{\mathbf{A}_{\text{inf}}} \mathbf{A}_{\text{inf}}(U) \rightarrow A_{\mathfrak{y}},$$

by the uniqueness, the above map $\tilde{\mathcal{R}} \rightarrow A_{\mathfrak{y}}$ is also the natural map

$$\tilde{\mathcal{R}} \rightarrow \tilde{\mathcal{R}} \hat{\otimes}_{\mathbf{A}_{\text{inf}}} \mathbf{A}_{\text{inf}}(U) \rightarrow A_{\mathfrak{y}}.$$

As argued in the second paragraph, this map uniquely extends to a morphism $\mathbf{A}_{\text{inf}}(U)$ -algebras $A_{\mathfrak{y}} \rightarrow A_{\mathfrak{y}}$, which is nothing but $\iota \circ j$ by construction. Then the uniqueness criterion forces $\iota \circ j = \text{id}$ as desired. \square

Proposition 2.6. *The map of $\tilde{\mathcal{C}}^{1,+}(U)$ -algebras*

$$\iota : \tilde{\mathcal{C}}^{1,+}[U_1, \dots, U_d]_{\text{pd}}^{\wedge} \rightarrow C_{\mathfrak{y}}^+ \text{ (resp. } \iota_n : \tilde{\mathcal{C}}^{1,+}[U_1, \dots, U_d]_{\text{pd}}^{\wedge} / (\xi p^{-1})^n \rightarrow C_{\mathfrak{y}}^+ / (\xi p^{-1})^n)$$

sending each U_i to $\frac{u(T_i - [T_i^b])}{\xi}$ is a well-defined isomorphism for $p \geq 3$ (resp. for arbitrary p and for any $n \geq 1$) such that the derivation d on $C_{\mathfrak{y}}^+$ reads

$$d = \sum_{i=1}^d u \frac{\partial}{\partial U_i} \otimes \frac{dT_i}{\xi} : \tilde{\mathcal{C}}^{1,+}[U_1, \dots, U_d]_{\text{pd}}^{\wedge} \rightarrow \oplus_{i=1}^d \tilde{\mathcal{C}}^{1,+}[U_1, \dots, U_d]_{\text{pd}}^{\wedge} \cdot \frac{dT_i}{\xi}$$

$$\text{(resp. } d = \sum_{i=1}^d u \frac{\partial}{\partial U_i} \otimes \frac{dT_i}{\xi} : \tilde{\mathcal{C}}^{1,+}[U_1, \dots, U_d]_{\text{pd}}^{\wedge} / (\xi p^{-1})^n \rightarrow \oplus_{i=1}^d \tilde{\mathcal{C}}^{1,+}[U_1, \dots, U_d]_{\text{pd}}^{\wedge} / (\xi p^{-1})^n \cdot \frac{dT_i}{\xi})$$

via the above isomorphism and the isomorphism $\Omega_{\tilde{\mathcal{R}}}^1\{-1\} \cong \oplus_{i=1}^d \tilde{\mathcal{R}} \cdot \frac{dT_i}{\xi}$.

Proof. It suffices to show that ι is a well-defined isomorphism, and the rest follows directly. As T_i is invertible in $\tilde{\mathcal{R}}$, by Lemma 2.5, we have the isomorphism

$$\mathbf{A}_{\text{inf}}(U)[[V_1, \dots, V_d]] \xrightarrow{V_i \mapsto (T_i - [T_i^b])} A_{\mathfrak{y}}.$$

Thus, by construction of $C_{\mathfrak{y}}^+$, we have an isomorphism

$$\mathbf{A}_{\text{inf}}(U)[[V_1, \dots, V_d]][u, \frac{uV_1}{\xi}, \dots, \frac{uV_d}{\xi}]_{\text{pd}}^{\wedge} \hat{\otimes}_{\mathbf{A}_{\text{inf}}(U)} \tilde{\mathcal{C}}^{1,+}(U) \xrightarrow{V_i \mapsto (T_i - [T_i^b])} C_{\mathfrak{y}}^+,$$

where $\mathbb{A}_{\text{inf}}(U)[[V_1, \dots, V_d]][u, \frac{uV_1}{\xi}, \dots, \frac{uV_d}{\xi}]_{\text{pd}}^\wedge$ is the $(p, \xi, V_1, \dots, V_d)$ -adic completion of the pd-algebra $\mathbb{A}_{\text{inf}}(U)[[V_1, \dots, V_d]][u, \frac{uV_1}{\xi}, \dots, \frac{uV_d}{\xi}]_{\text{pd}}$. Thus, to show ι is an isomorphism, it suffices to show that the map

$$i : \tilde{\mathbb{C}}^{1,+}(U)[U_1, \dots, U_d]_{\text{pd}}^\wedge \rightarrow \mathbb{A}_{\text{inf}}(U)[[V_1, \dots, V_d]][u, \frac{uV_1}{\xi}, \dots, \frac{uV_d}{\xi}]_{\text{pd}}^\wedge \hat{\otimes}_{\mathbb{A}_{\text{inf}}(U)} \tilde{\mathbb{C}}^{1,+}(U)$$

sending U_i to $\frac{uV_i}{\xi}$ is a well-defined isomorphism.

The well-definedness of i is trivial as $\mathbb{A}_{\text{inf}}(U)[[V_1, \dots, V_d]][u, \frac{uV_1}{\xi}, \dots, \frac{uV_d}{\xi}]_{\text{pd}}^\wedge \hat{\otimes}_{\mathbb{A}_{\text{inf}}(U)} \tilde{\mathbb{C}}^{1,+}(U)$ is a p -adic complete $\tilde{\mathbb{C}}^{1,+}$ -algebra and $\frac{uV_i}{\xi}$ admits arbitrary pd-powers. Now, we construct the inverse of i as follows:

As $|\frac{\xi}{u}|_1 = p^{\frac{1}{p-1}-1} < 1$ (this is where we use $p \geq 3$), we see from Lemma 2.2(2) that $\frac{\xi}{u} \in \tilde{\mathbb{C}}^{1,+}(U)$ is topologically nilpotent. Thus, the map of $\mathbb{A}_{\text{inf}}(U)$ -algebras

$$j : \mathbb{A}_{\text{inf}}(U)[V_1, \dots, V_d] \rightarrow \tilde{\mathbb{C}}^{1,+}(U)[U_1, \dots, U_d]_{\text{pd}}^\wedge$$

sending V_i to $\frac{\xi U_i}{u}$ uniquely extends to a map (still denoted by)

$$j : \mathbb{A}_{\text{inf}}(U)[[V_1, \dots, V_d]] \rightarrow \tilde{\mathbb{C}}^{1,+}(U)[U_1, \dots, U_d]_{\text{pd}}^\wedge.$$

Since u, U_1, \dots, U_d admit arbitrary pd-powers in $\tilde{\mathbb{C}}^{1,+}(U)[U_1, \dots, U_d]_{\text{pd}}^\wedge$, as argued in the proof of Lemma 2.5, it is easy to see that j uniquely extends to a morphism of $\tilde{\mathbb{C}}^{1,+}(U)$ -algebras

$$j : \mathbb{A}_{\text{inf}}(U)[[V_1, \dots, V_d]][u, \frac{uV_1}{\xi}, \dots, \frac{uV_d}{\xi}]_{\text{pd}}^\wedge \hat{\otimes}_{\mathbb{A}_{\text{inf}}(U)} \tilde{\mathbb{C}}^{1,+}(U) \rightarrow \tilde{\mathbb{C}}^{1,+}(U)[U_1, \dots, U_d]_{\text{pd}}^\wedge$$

sending V_i to $\frac{\xi U_i}{u}$. It is clear that j is exactly the inverse of i .

To see ι_n is an isomorphism, it suffices to note that $\frac{\xi}{u} = \frac{\xi}{p} \cdot \frac{p}{u}$ is nilpotent modulo $(\xi p^{-1})^n$. Using this instead of $|\frac{\xi}{u}|_1 < 1$ above, it is easy to construct the inverse of ι_n . \square

Corollary 2.7. *The morphism of sheaves of $\tilde{\mathbb{C}}_{|X_\infty}^{1,+}$ -algebras (resp. $\mathbb{B}_{\text{dR}|X_\infty}^+$ -algebras)*

$$\iota : \tilde{\mathbb{C}}_{|X_\infty}^{1,+}[U_1, \dots, U_d]_{\text{pd}}^\wedge \rightarrow \mathcal{O}\tilde{\mathbb{C}}_{\text{pd}|X_\infty}^+ \text{ (resp. } \iota : \mathbb{B}_{\text{dR}|X_\infty}^+ \langle U_1, \dots, U_d \rangle_{\text{pd}} \rightarrow \mathcal{O}\mathbb{B}_{\text{dR,pd}|X_\infty}^+)$$

sending U_i to $\frac{u(T_i - [T_i^b])}{\xi}$ is an isomorphism for $p \geq 3$ (resp. for arbitrary p). Here, $\mathbb{B}_{\text{dR}|X_\infty}^+ \langle U_1, \dots, U_d \rangle_{\text{pd}}$ denotes the ξ -adic completion of

$$\tilde{\mathbb{C}}_{|X_\infty}^1[U_1, \dots, U_d]_{\text{pd}}^\wedge = \tilde{\mathbb{C}}_{|X_\infty}^{1,+}[U_1, \dots, U_d]_{\text{pd}}^\wedge[\frac{1}{p}].$$

Proof. The result follows from Corollary 2.6 directly. For the \mathbb{B}_{dR}^+ case, just note that by definition

$$\mathcal{O}\mathbb{B}_{\text{dR,pd}}^+ = \varprojlim_n \mathcal{O}\tilde{\mathbb{C}}^+[\frac{1}{p}]/(\xi p^{-1})^n.$$

\square

Now, we are able to prove Poincaré's Lemma 2.4

Proof of Theorem 2.4. Since the problem is local on both $\mathfrak{X}_{\text{ét}}$ and $X_{\text{proét}}$, we may assume $\mathfrak{X} = \text{Spf}(\mathcal{R})$ with X_∞ as above. Then the result follows from Corollary 2.7 directly. \square

Now we still assume $\mathfrak{X} = \text{Spf}(\mathcal{R})$ is small and keep the notations as above.

Proposition 2.8. *The series $\frac{u^2}{t} \log \frac{[T_i^b]}{T_i} = \frac{u^2}{t} \sum_{n \geq 1} \frac{(1 - \frac{[T_i^b]}{T_i})^n}{n}$ converges in $\mathcal{O}\tilde{\mathbb{C}}_{\text{pd}|X_\infty}^+$. The morphism of the sheaves of $\tilde{\mathbb{C}}_{|X_\infty}^+$ -algebras (resp. $\mathbb{B}_{\text{dR}|X_\infty}^+$ -algebras)*

$$\iota : \tilde{\mathbb{C}}_{|X_\infty}^{1,+}[W_1, \dots, W_d]_{\text{pd}}^\wedge \rightarrow \mathcal{O}\tilde{\mathbb{C}}_{\text{pd}|X_\infty}^+ \text{ (resp. } \iota : \mathbb{B}_{\text{dR}|X_\infty}^+ \langle W_1, \dots, W_d \rangle_{\text{pd}} \rightarrow \mathcal{O}\mathbb{B}_{\text{dR,pd}|X_\infty}^+)$$

sending W_i to $\frac{u^2}{t} \log \frac{[T_i^b]}{T_i}$ is an isomorphism for $p \geq 3$ (resp. for arbitrary p). Via the above morphism, the derivation d on $\mathcal{O}\tilde{\mathbb{C}}_{\text{pd}|X_\infty}^+$ (resp. $\mathcal{O}\mathbb{B}_{\text{dR},\text{pd}|X_\infty}^+$) reads

$$d = \sum_{i=1}^d -u \frac{\partial}{\partial W_i} \otimes \frac{u}{t} d\log T_i,$$

where we trivialize $\Omega_{\tilde{\mathcal{R}}}^1\{-1\}$ by $\oplus_{i=1}^d \tilde{\mathcal{R}} \cdot \frac{u}{t} d\log T_i$, and the Galois group Γ acts on $\mathcal{O}\tilde{\mathbb{C}}_{\text{pd}|X_\infty}^+$ such that

$$\gamma_i(W_j) = W_j + \delta_{ij}u^2, \quad \forall 1 \leq i, j \leq d.$$

Proof. We first deal with the $\mathcal{O}\tilde{\mathbb{C}}^+$ case. Recall t is a unit-multiple of ξu in $\tilde{\mathbb{C}}^{1,+}(U)$ (cf. Lemma 2.2) and T_i is invertible in $\tilde{\mathcal{R}}$, it follows from Corollary 2.7 that the morphism

$$\tilde{\mathbb{C}}_{|X_\infty}^{1,+}[U_1, \dots, U_d]_{\text{pd}}^\wedge \rightarrow \mathcal{O}\tilde{\mathbb{C}}_{\text{pd}|X_\infty}^+$$

sending U_i to $\frac{u^2}{t}(1 - \frac{[T_i^b]}{T_i})$ is an isomorphism. Thus, we need to show the series

$$\frac{u^2}{t} \sum_{n \geq 1} \frac{(\frac{u^2}{t} U_i)^n}{n}$$

converges in $\tilde{\mathbb{C}}_{|X_\infty}^{1,+}[U_1, \dots, U_d]_{\text{pd}}^\wedge$ and the morphism

$$i : \tilde{\mathbb{C}}_{|X_\infty}^{1,+}[W_1, \dots, W_d]_{\text{pd}}^\wedge \rightarrow \tilde{\mathbb{C}}_{|X_\infty}^{1,+}[U_1, \dots, U_d]_{\text{pd}}^\wedge$$

sending W_i to $\frac{u^2}{t} \sum_{n \geq 1} \frac{(\frac{t}{u^2} U_i)^n}{n}$ is an isomorphism. However, we have

$$\frac{u^2}{t} \sum_{n \geq 1} \frac{(\frac{t}{u^2} U_i)^n}{n} = \sum_{n \geq 1} \left(\frac{t}{u^2}\right)^{n-1} (n-1)! U_i^{[n]}.$$

As $(\frac{t}{u^2})^{n-1} (n-1)! \rightarrow 0$ as $n \rightarrow +\infty$ in $\tilde{\mathbb{C}}^{1,+}(U)$, we get the desired convergence. Now, consider the map

$$j : \tilde{\mathbb{C}}_{|X_\infty}^{1,+}[U_1, \dots, U_d]_{\text{pd}}^\wedge \rightarrow \tilde{\mathbb{C}}_{|X_\infty}^{1,+}[W_1, \dots, W_d]_{\text{pd}}^\wedge$$

sending U_i to $\frac{u^2}{t}(1 - \exp(\frac{t}{u^2} W_i)) = -\sum_{n \geq 1} (\frac{t}{u^2})^{n-1} W_i^{[n]}$. We claim it is well-defined: Indeed, as

$$|\frac{t}{u^2}|_1 = |\frac{\xi}{u}|_1 = p^{\frac{1}{p-1}-1},$$

$(\frac{t}{u^2})^{n-1} \rightarrow 0$ as $n \rightarrow +\infty$ in $\tilde{\mathbb{C}}^{1,+}(U)$ (here we use $p \geq 3$), yielding the desired well-definedness. Now one can conclude by checking j is the inverse of i directly.

For the $\mathcal{O}\mathbb{B}_{\text{dR},\text{pd}|X_\infty}^+$ case, note that for arbitrary p , the above j is always defined modulo $(\xi p^{-1})^m$ for an $m \geq 1$ (as $(\frac{t}{u^2})^m$ vanishes in $\tilde{\mathbb{C}}^{1,+}(U)/(\xi p^{-1})^m$). Thus, the above i is an isomorphism modulo $(\xi p^{-1})^m$ for any $m \geq 1$. Then one can conclude by inverting p and taking ξ -adic completion. \square

The above construction is compatible with base-change of $(A, A^+) \in \text{Perfd}$.

Proposition 2.9. *Let $(A, A^+) \rightarrow (B, B^+)$ be a morphism in Perfd . Let \mathfrak{X}_A be a smooth formal scheme over A^+ with a fixed lifting $\tilde{\mathfrak{X}}_A$ over $\mathbb{A}_{\text{inf}}(A, A^+)$. Let $\mathfrak{X}_B := \mathfrak{X}_A \times_{\text{Spf}(A^+)} \text{Spf}(B^+)$ be the induced smooth formal scheme over B^+ with the lifting $\tilde{\mathfrak{X}}_B = \tilde{\mathfrak{X}}_A \times_{\text{Spf}(\mathbb{A}_{\text{inf}}(A, A^+))} \text{Spf}(\mathbb{A}_{\text{inf}}(B, B^+))$. Let $(\mathcal{O}\mathbb{B}_{\text{dR},\text{pd},A}^+, d)$ and $(\mathcal{O}\mathbb{B}_{\text{dR},\text{pd},B}^+, d)$ be the period sheaves with connections corresponding to $\tilde{\mathfrak{X}}_A$ and $\tilde{\mathfrak{X}}_B$, respectively. Then there exists a natural isomorphism*

$$(\mathbb{B}_{\text{dR}}^+(B, B^+) \hat{\otimes}_{\mathbb{B}_{\text{dR}}^+(A, A^+)} \mathcal{O}\mathbb{B}_{\text{dR},\text{pd},A}^+, \text{id} \otimes d) \rightarrow (\mathcal{O}\mathbb{B}_{\text{dR},\text{pd},B}^+, d)$$

compatible with connections.

Proof. The existence of the morphism

$$\mathbb{B}_{\mathrm{dR}}^+(B, B^+) \hat{\otimes}_{\mathbb{B}_{\mathrm{dR}}^+(A, A^+)} \mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}, A}^+ \rightarrow \mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}, B}^+$$

follows directly from the construction of $\mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}}^+$ above. To see it is an isomorphism, we are reduced to the case $\mathfrak{X}_A = \mathrm{Spf}(\mathcal{R})$ is small. Then $\mathfrak{X}_B = \mathrm{Spf}(\mathcal{R} \hat{\otimes}_{A^+} B^+)$. Note that $X_{\infty, B} = X_{\infty, A} \times_{\mathrm{Spa}(A, A^+)} \mathrm{Spa}(B, B^+)$. One can conclude by using Proposition 2.8. \square

Remark 2.10. If we put \mathbb{A}_{dR} and $\mathcal{O}\mathbb{A}_{\mathrm{dR}}$ the (ξp^{-1}) -adic completion of $\tilde{\mathcal{C}}^{1,+}$ and $\mathcal{O}\tilde{\mathcal{C}}^{1,+}$, respectively, then the derivation d on $\mathcal{O}_{\tilde{\mathfrak{X}}}$ induces a derivation

$$d : \mathcal{O}\mathbb{A}_{\mathrm{dR}} \rightarrow \mathcal{O}\mathbb{A}_{\mathrm{dR}} \otimes_{\mathcal{O}_{\tilde{\mathfrak{X}}}} u \cdot \Omega_{\tilde{\mathfrak{X}}}^1 \{-1\}.$$

The last paragraph in the proof of Proposition 2.8 together with Proposition 2.6 tells us when $\mathfrak{X} = \mathrm{Spf}(\mathcal{R})$ is small, then the morphism

$$\iota : \mathbb{A}_{\mathrm{dR}|X_{\infty}} [W_1, \dots, W_d]_{\mathrm{pd}}^{\wedge} \rightarrow \mathcal{O}\mathbb{A}_{\mathrm{dR}|X_{\infty}}$$

sending W_i to $\frac{u^2}{t} \log \frac{[T_i^p]}{T_i}$ is a well-defined isomorphism and gives the identification $d = -\sum_{i=1}^d u \frac{\partial}{\partial W_i} \otimes \frac{u}{t} d \log T_i$. One can check directly the sequence

$$0 \rightarrow \mathbb{A}_{\mathrm{dR}} \rightarrow \mathcal{O}\mathbb{A}_{\mathrm{dR}} \otimes_{\mathcal{O}_{\tilde{\mathfrak{X}}}} u \cdot \Omega_{\tilde{\mathfrak{X}}}^1 \{-1\} \rightarrow \dots \rightarrow \mathcal{O}\mathbb{A}_{\mathrm{dR}} \otimes_{\mathcal{O}_{\tilde{\mathfrak{X}}}} u^d \cdot \Omega_{\tilde{\mathfrak{X}}}^d \{-d\}$$

is exact; that is, the Poincaré's Lemma holds true in this case. This leads to an integral version of p -adic Riemann–Hilbert correspondence lifting the integral p -adic Simpson correspondence in [MW23].

Remark 2.11. In [CLWZ24, §3], given a lifting $\tilde{\mathfrak{X}}$ of \mathfrak{X} , one can construct an overconvergent de Rham period sheaf with connection $(\mathcal{O}\mathbb{B}_{\mathrm{dR}}^{\dagger,+}, d : \mathcal{O}\mathbb{B}_{\mathrm{dR}}^{\dagger,+} \rightarrow \mathcal{O}\mathbb{B}_{\mathrm{dR}}^{\dagger,+} \otimes_{\mathcal{O}_{\tilde{\mathfrak{X}}}} \Omega_{\tilde{\mathfrak{X}}}^1 \{-1\})$. One can check $(\mathcal{O}\mathbb{B}_{\mathrm{dR}}^{\dagger,+}, d)$ can be viewed as a sub- $\mathbb{B}_{\mathrm{dR}}^+$ -algebra of $(\mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}}^+, d)$ compatible with connections. Indeed, let $(\widehat{\mathcal{O}\mathbb{B}_{\mathrm{dR}}^+}, d)$ be the ξ -adic completion of the period sheaf with connection $(\mathcal{O}\tilde{\mathcal{C}}^{1,1}, d)$ in [CLWZ24, Def. 2.29] with the d -preserving lattice $\mathcal{O}\tilde{\mathcal{C}}^{1,1,+}$, and then it clearly contains $(\mathcal{O}\mathbb{B}_{\mathrm{dR}}^{\dagger,+}, d)$ as a sub- $\mathbb{B}_{\mathrm{dR}}^+$ -algebra compatible with connections. On the other hand, as u admits pd -powers in $\mathcal{O}\tilde{\mathcal{C}}^{1,1,+}$, one can construct a natural map $\mathcal{O}\tilde{\mathcal{C}}_{\mathrm{pd}}^+ \rightarrow \mathcal{O}\tilde{\mathcal{C}}^{1,1,+}$ compatible with connections, yielding a well-defined injection $\mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}}^+ \rightarrow \widehat{\mathcal{O}\mathbb{B}_{\mathrm{dR}}^+}$ compatible with connections by inverting p and taking ξ -adic completion. One can show the natural inclusion $\mathcal{O}\mathbb{B}_{\mathrm{dR}}^{\dagger,+} \hookrightarrow \widehat{\mathcal{O}\mathbb{B}_{\mathrm{dR}}^+}$ factors through $\mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}}^+$ by working locally; that is, working with \mathfrak{X} small affine.

Remark 2.12. The $(\mathcal{O}\tilde{\mathcal{C}}_{\mathrm{pd}}^+, d)$ and $(\mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}}^+, d)$ reduce to the $(\mathcal{O}\widehat{\mathcal{C}}_{\mathrm{pd}}^+, \Theta)$ and $(\mathcal{O}\widehat{\mathcal{C}}_{\mathrm{pd}}, \Theta)$ modulo (ξp^{-1}) . This follows by combining arguments in Remark 2.11 and [CLWZ24, Rem. 3.24], together with the identification in [MW23, Prop. 2.10]. We now check this on $X_{\infty, \mathrm{pro\acute{e}t}}$ for \mathfrak{X} small affine: Put $Y_i = \frac{u}{t} \log \frac{[T_i^p]}{T_i}$, and then the reduction of $(\mathcal{O}\tilde{\mathcal{C}}_{\mathrm{pd}}^+, d)$ is given by

$$(\widehat{\mathcal{O}}_X^+[(\zeta_p - 1)\underline{Y}]_{\mathrm{pd}}^{\wedge}, -\sum_{i=1}^d \frac{\partial}{\partial Y_i} \otimes (\zeta_p - 1) \cdot \frac{1}{t} d \log T_i)$$

with the Γ -action such that

$$\gamma_i(Y_j) = Y_j + \delta_{ij}(\zeta_p - 1), \quad \forall 1 \leq i, j \leq d$$

by noting that the reduction of u is $\zeta_p - 1$. Now, one can conclude by applying [MW23, Cor. 2.6]. One can also check the reduction of $\mathcal{O}\tilde{\mathcal{C}}_{\mathrm{pd}}^+$ modulo ξp^{-1} behaves the same as $\mathcal{B}_{\tilde{\mathfrak{X}}}$ appearing in [AHLB23b, Lem. 3.8] (see also [AHLB23a, Prop. 4.8]).

3. LOCAL SMALL RIEMANN–HILBERT CORRESPONDENCE

In this section, we fix a $\mathrm{Spa}(A, A^+) \in \mathrm{Perfd}$ and let \mathfrak{X} be a liftable smooth formal scheme over A^+ with a fixed lifting $\tilde{\mathfrak{X}}$ over $\mathbf{A}_{\mathrm{inf}}$. Let $(\mathcal{O}\mathbb{B}_{\mathrm{dR}, \mathrm{pd}}^+, d)$ be the period sheaf with connection defined in the previous section and denote by $(\mathcal{O}\widehat{\mathcal{C}}_{\mathrm{pd}}, \Theta)$ its reduction modulo t .

3.1. A criterion for being Hitchin-small. Denote by $v\text{Bun}_r(X)(A)^{\text{H-small}}$ and by $\text{HIG}_r(X)(A)^{\text{H-small}}$ the category of Hitchin-small v -vector bundles on X_v of rank r and the category of Hitchin-small Higgs bundles on $X_{\text{ét}}$ of rank r . Then

$$v\text{Bun}(X)(A)^{\text{H-small}} := \bigcup_{r \geq 1} v\text{Bun}_r(X)(A)^{\text{H-small}} \text{ and } \text{HIG}(X)(A)^{\text{H-small}} := \bigcup_{r \geq 1} \text{HIG}_r(X)(A)^{\text{H-small}}$$

are the whole categories of v -vector bundles and Higgs bundles. Let $\mathcal{A}_r(A)$ and $\mathcal{A}_r^{\text{H-small}}(A)$ be the Hitchin base and its Hitchin-small locus associated to X with respect to the fixed rank r . Recall a v -vector bundle \mathcal{L} (resp. a Higgs bundle (\mathcal{H}, θ)) of rank r is Hitchin-small if and only if its image $\tilde{h}(\mathcal{L})$ (resp. $h(\mathcal{H}, \theta)$) in $\mathcal{A}_r(A)$ belongs to $\mathcal{A}_r^{\text{H-small}}(A)$.

We first give a more convenient description of Hitchin-small Higgs bundles in [AHLB23a].

Proposition 3.1. *A Higgs bundle (\mathcal{H}, θ) on $X_{\text{ét}}$ is Hitchin-small if and only if étale locally on \mathfrak{X} , there is a Higgs bundle on \mathfrak{X} , i.e. a locally finite free $\mathcal{O}_{\mathfrak{X}}$ -module \mathcal{H}^+ with a Higgs field $\theta^+ : \mathcal{H}^+ \rightarrow \mathcal{H}^+ \otimes_{\mathcal{O}_{\mathfrak{X}}} \Omega_{\mathfrak{X}}^1\{-1\}$, such that*

- (1) $(\mathcal{H}, \theta_{\mathcal{H}}) = (\mathcal{H}^+, \theta^+)[\frac{1}{p}]$, and that
- (2) the $(\zeta_p - 1)$ -twist of θ^+ is topologically nilpotent; that is, θ^+ takes values in $(\zeta_p - 1)\mathcal{H} \otimes_{\mathcal{O}_{\mathfrak{X}}} \Omega_{\mathfrak{X}}^1\{-1\}$ such that

$$\frac{1}{\zeta_p - 1} \theta^+ : \mathcal{H} \rightarrow \mathcal{H} \otimes_{\mathcal{O}_{\mathfrak{X}}} \Omega_{\mathfrak{X}}^1\{-1\}$$

is (p -adically) topologically nilpotent.

Indeed, one can always find such an $(\mathcal{H}^+, \theta^+)$ on any $\mathfrak{Y} = \text{Spf}(\mathcal{R}) \in \mathfrak{X}_{\text{ét}}$ which is small affine.

Proof. Recall one can check Hitchin-smallness of Higgs bundle by passing to geometric points in the sense of [AHLB23b, Def. 3.2]. Compared with [AHLB23a, Def. 6.24(2)], the result follows from [AHLB23a, Th. 7.13(3)]. Indeed, one can find the desired lattice $(\mathcal{H}^+, \theta^+)$ on any affine $\mathfrak{Y} = \text{Spf}(\mathcal{R}) \in \mathfrak{X}_{\text{ét}}$ splitting the Hodge–Tate gerbe \mathfrak{Y}^{HT} (cf. [AHLB23a, Th. 7.11(3)]). Note that the Hodge–Tate gerbe \mathfrak{Y} always splits when \mathfrak{Y} is affine small by [BL22, Prop. 5.12]. \square

We also want to give an explicit description of Hitchin-small v -vector bundles. Let $(\mathcal{B}, \Theta_{\mathcal{B}})$ be the sheaf with Higgs field in [AHLB23a, Def. 7.17]. Recall the following constructions in *loc.cit.*: For any $(\mathcal{H}, \theta) \in \text{HIG}(X)(A)^{\text{H-small}}$, define

$$(3.1) \quad \Theta'_{\mathcal{H}} := \theta \otimes \text{id} + \text{id} \otimes \Theta_{\mathcal{B}} : \mathcal{H} \otimes_{\mathcal{O}_X} \mathcal{B} \rightarrow \mathcal{H} \otimes_{\mathcal{O}_X} \mathcal{B} \otimes_{\mathcal{O}_{\mathfrak{X}}} \Omega_{\mathfrak{X}}^1\{-1\}$$

and define

$$\mathcal{L}(\mathcal{H}, \theta_{\mathcal{H}}) := (\mathcal{H} \otimes_{\mathcal{O}_X} \mathcal{B})^{\Theta'_{\mathcal{H}}=0}.$$

Then we have the following result:

Theorem 3.2 ([AHLB23b, Th. 3.20], [AHLB23a, Th. 7.13(1)]). *The functor $(\mathcal{H}, \theta_{\mathcal{H}}) \mapsto \mathcal{L}(\mathcal{H}, \theta_{\mathcal{H}})$ above induces an equivalence between the categories*

$$v\text{Bun}(X)(A)^{\text{H-small}} \simeq \text{HIG}(X)(A)^{\text{H-small}}$$

which is compatible with cohomology in the sense that we have a quasi-isomorphism

$$\text{R}\nu_*(\mathcal{L}(\mathcal{H}, \theta_{\mathcal{H}})) \simeq \text{HIG}(\mathcal{H}, \theta_{\mathcal{H}}).$$

Corollary 3.3. *Assume \mathfrak{X} is small affine. For any $(\mathcal{H}, \theta) \in \text{HIG}(X)^{\text{H-small}}(A)$, define*

$$\Theta_{\mathcal{H}} := \theta \otimes \text{id} + \text{id} \otimes \Theta : \theta \otimes \text{id} + \text{id} \otimes \Theta : \mathcal{H} \otimes_{\mathcal{O}_X} \mathcal{O}\widehat{\mathcal{C}}_{\text{pd}} \rightarrow \mathcal{H} \otimes_{\mathcal{O}_{\mathfrak{X}}} \mathcal{O}\widehat{\mathcal{C}}_{\text{pd}} \otimes_{\mathcal{O}_{\mathfrak{X}}} \Omega_{\mathfrak{X}}^1\{-1\}.$$

Then $\mathcal{L}(\mathcal{H}, \theta_{\mathcal{H}}) = (\mathcal{H} \otimes_{\mathcal{O}_X} \mathcal{O}\widehat{\mathcal{C}}_{\text{pd}})^{\Theta_{\mathcal{H}}=0}$. In particular, the functor

$$\text{HIG}(X)(A)^{\text{H-small}} \rightarrow v\text{Bun}(X)(A)^{\text{H-small}}, \quad (\mathcal{H}, \theta) \mapsto (\mathcal{H} \otimes_{\mathcal{O}_X} \mathcal{O}\widehat{\mathcal{C}}_{\text{pd}})^{\Theta_{\mathcal{H}}=0}$$

is an equivalence of categories which is compatible with cohomology.

Proof. Recall we have an identification $(\mathcal{O}\widehat{\mathcal{C}}_{\text{pd}}, \Theta)|_{X_{\infty}} = (\mathcal{B}, \Theta_{\mathcal{B}})|_{X_{\infty}}$ by Remark 2.12. The result follows from Theorem 3.2 directly. \square

Now, let $\mathfrak{X} = \mathrm{Spf}(\mathcal{R})$ be small affine with the generic fiber $\mathrm{Spa}(R, R^+)$ and write $\Omega_{\mathfrak{X}}^1\{-1\} = \bigoplus_{i=1}^d \mathcal{R} \cdot (\zeta_p - 1) \frac{d\log T_i}{t}$. Define

$$(B_{\infty}^+, \Theta) := (\mathcal{O}\widehat{\mathcal{C}}_{\mathrm{pd}}^+, \Theta)(X_{\infty})$$

and then we have

$$(3.2) \quad (B_{\infty}^+, \Theta) = (\widehat{R}_{\infty}^+[W]_{\mathrm{pd}}^{\wedge}, - \sum_{i=1}^d (\zeta_p - 1) \frac{\partial}{\partial W_i} \otimes (\zeta_p - 1) \cdot \frac{1}{t} d\log T_i)$$

by Proposition 2.8 such that $\Gamma = \bigoplus_{i=1}^d \mathbb{Z}_p \gamma_i$ acts on B_{∞}^+ via

$$\gamma_i(W_j) = W_j + \delta_{ij}(\zeta_p - 1)^2, \quad \forall 1 \leq i, j \leq d.$$

Set $B^+ = R^+[W]_{\mathrm{pd}}^{\wedge}$, and then it is Θ -preserving and stable under the action of Γ .

Now let (H, θ) be the global section of a Hitchin-small Higgs bundle on $X = \mathrm{Spa}(R, R^+)$, and

$$(H^+, \theta^+ : H^+ \rightarrow H^+ \cdot (\zeta_p - 1) \frac{d\log T_i}{t})$$

be the lattice of (H, θ) satisfying the condition in Proposition 3.1. Then one can write

$$\theta^+ = \sum_{i=1}^d (\zeta_p - 1) \theta_i \otimes (\zeta_p - 1) \frac{d\log T_i}{t}$$

with $\theta_i \in \mathrm{End}_{R^+}(H^+)$ topologically nilpotent. Put

$$\Theta_H^+ := \theta^+ \otimes \mathrm{id} + \mathrm{id} \otimes \Theta : H^+ \otimes_{\mathcal{R}} B_{\infty}^+ \rightarrow H^+ \otimes_{\mathcal{R}} B_{\infty}^+ \otimes_{\mathcal{R}} \Omega_{\mathcal{R}}^1\{-1\}$$

and then $H^+ \otimes_{\mathcal{R}} B_{\infty}^+$ is Θ_H^+ -preserving. Put

$$L^+(H^+, \theta^+) := (H^+ \otimes_{\mathcal{R}} B^+)_{\Theta_H^+ = 0} \text{ and } L_{\infty}^+(H^+, \theta^+) := (H^+ \otimes_{\mathcal{R}} B_{\infty}^+)_{\Theta_H^+ = 0}.$$

Then $L^+(H^+, \theta^+)$ and $L_{\infty}^+(H^+, \theta^+)$ inherit the Γ -actions from B_{∞}^+ satisfying

$$L_{\infty}^+(H^+, \theta^+) = L^+(H^+, \theta^+) \otimes_{\mathcal{R}} \widehat{R}_{\infty}^+.$$

Lemma 3.4. *The $L^+ := L^+(H^+, \theta^+)$ is a finite projective R^+ -module which is isomorphic to H^+ such that for any $x \in L^+$ and any $1 \leq i \leq d$, we have*

$$(3.3) \quad L^+ = \exp\left(\sum_{i=1}^d \theta_i W_i\right)(H^+) = \{x \in H^+ \otimes_{\mathcal{R}} B^+ \mid x = \sum_{J \in \mathbb{N}^d} \underline{\theta}^J(y) \underline{W}^{[J]}, y \in H^+\}$$

$$\gamma_i(x) = \exp(-\theta_i(\zeta_p - 1)^2)(x).$$

Proof. Let $x = \sum_{J \in \mathbb{N}^d} x_J \underline{W}^{[J]}$ be an element in $H^+ \otimes_{\mathcal{R}} B^+ \cong H^+[\underline{W}]_{\mathrm{pd}}^{\wedge}$. Then we have

$$\begin{aligned} \Theta_H^+(x) &= \sum_{i=1}^d \sum_{J \in \mathbb{N}^d} ((\zeta_p - 1) \theta_i(x_J) \underline{W}^{[J]} - (\zeta_p - 1) x_J \underline{W}^{[J - E_i]}) \otimes (\zeta_p - 1) \frac{d\log T_i}{t} \\ &= (\zeta_p - 1) \sum_{i=1}^d \sum_{J \in \mathbb{N}^d} (\theta_i(x_J) - x_{J+E_i}) \underline{W}^{[J]} \otimes (\zeta_p - 1) \frac{d\log T_i}{t}. \end{aligned}$$

Thus, $x \in L^+$ if and only if for any $J \in \mathbb{N}^d$ and any $1 \leq i \leq d$, $x_{J+E_i} = \theta_i(x_J)$. Using this, by iteration, we see that $x \in L^+$ if and only if

$$x = \sum_{J \in \mathbb{N}^d} \underline{\theta}^J(x_0) \underline{W}^{[J]} = \exp\left(\sum_{i=1}^d \theta_i W_i\right)(x).$$

So we have

$$L^+ = \exp\left(\sum_{i=1}^d \theta_i W_i\right)(H^+) \cong H^+$$

and the description of Γ on L^+ follows as $\gamma_i(W_j) = W_j + \delta_{ij}(\zeta_p - 1)^2$ for any $1 \leq i, j \leq d$. \square

Note that $L_\infty(H, \theta) := L_\infty^+(H^+, \theta^+)$ is exactly $\mathcal{L}(\mathcal{H}, \theta)(X_\infty)$. Thus, if $\mathcal{L} \in v\text{Bun}(X)^{\text{H-small}}(A)$, then there must be a finite projective \mathcal{R} -module H^+ endowed with an action of Γ such that

$$\mathcal{L}(X_\infty) \cong H^+ \otimes_{\mathcal{R}} \widehat{R}_\infty$$

and that for any $1 \leq i \leq d$, there exists a topologically nilpotent $\theta_i \in \text{End}_{\mathcal{R}}(H^+)$ such that for any $x \in H^+$, we have $\gamma_i(x) = \exp(-\theta_i(\zeta_p - 1)^2)(x)$. We will show this condition is also sufficient. The arguments in what follows are closely related with [AHLB23b, §3] and [MW23, §3 and Th. 4.1].

Definition 3.5. Let $C \in \{\mathcal{R} = R^+, R, \widehat{R}_\infty^+, \widehat{R}_\infty\}$

- (1) By an *Hitchin-small representation* of Γ over \mathcal{R} of rank r , we mean a finite projective \mathcal{R} -module L^+ endowed with an action of Γ such that for any $1 \leq i \leq d$, there exists a topologically nilpotent $\theta_i \in \text{End}_{\mathcal{R}}(L^+)$ such that for any $x \in L^+$, we have $\gamma_i(x) = \exp(-\theta_i(\zeta_p - 1)^2)(x)$. Denote by $\text{Rep}_{\Gamma}^{\text{H-small}}(\mathcal{R})$ the category of Hitchin-small representations of Γ over \mathcal{R} .
- (2) In general, by an *Hitchin-small representation* of Γ over B of rank r , we mean a finite projective C -module L endowed with an action of Γ such that there exists some $L^+ \in \text{Rep}_{\Gamma}^{\text{H-small}}(\mathcal{R})$ such that

$$L \cong L^+ \otimes_{\mathcal{R}} C.$$

Denote by $\text{Rep}_{\Gamma}^{\text{H-small}}(C)$ the category of Hitchin-small representations of Γ over C .

- (3) By a *Hitchin-small Higgs module* over \mathcal{R} of rank r , we mean a pair

$$(H^+, \theta^+ : H^+ \rightarrow H^+ \otimes_{\mathcal{R}} \Omega_{\mathcal{R}}^1\{-1\} = \bigoplus_{i=1}^d H^+ \cdot (\zeta_p - 1) \frac{d\log T_i}{t})$$

satisfying $\theta^+ \wedge \theta^+ = 0$ such that H^+ is a finite projective \mathcal{R} -module of rank r and $\theta = \sum_{i=1}^d (\zeta_p - 1) \theta_i \otimes (\zeta_p - 1) \frac{d\log T_i}{t}$ with $\theta_i \in \text{End}_{\mathcal{R}}(H^+)$ topologically nilpotent. Denote by $\text{HIG}^{\text{H-small}}(\mathcal{R})$ the category of Hitchin-small Higgs module over \mathcal{R} . A Higgs module (H, θ) over R of rank r is called *Hitchin-small* if it is of the form $(H, \theta) = (H^+, \theta^+)[\frac{1}{p}]$ for some $(H^+, \theta^+) \in \text{HIG}^{\text{H-small}}(\mathcal{R})$ of rank r . Denote by $\text{HIG}^{\text{H-small}}(R)$ the category of Hitchin-small Higgs module over R .

The following lemma is obvious.

Lemma 3.6. The base-change $L^+ \mapsto L_\infty^+ := L^+ \otimes_{\mathcal{R}} \widehat{R}_\infty^+$ induces an equivalence of categories

$$\text{Rep}_{\Gamma}^{\text{H-small}}(\mathcal{R}) \simeq \text{Rep}_{\Gamma}^{\text{H-small}}(\widehat{R}_\infty^+)$$

such that the natural map

$$\text{R}\Gamma(\Gamma, L^+) \rightarrow \text{R}\Gamma(\Gamma, L_\infty^+)$$

identifies the former with a direct summand of the latter whose complement is concentrated in degree ≥ 1 and killed by $\zeta_p - 1$.

Proof. To get the desired equivalence, we need to show the base-change functor is fully faithful. Note that L_∞ admits a Γ -equivariant decomposition:

$$(3.4) \quad L_\infty = \widehat{\bigoplus_{\underline{\alpha}=(\alpha_1, \dots, \alpha_d) \in (\mathbb{N} \cap [0,1))^d}} L^+ \cdot \underline{T}^\alpha,$$

where $T^\alpha = T_1^{\alpha_1} \cdots T_d^{\alpha_d}$. Then the desired full faithfulness and the whole lemma will follow from that for any $\underline{\alpha} \neq 0$,

$$\text{R}\Gamma(\Gamma, L^+ \cdot \underline{T}^\alpha)$$

is concentrated in degree ≥ 1 and killed by $\zeta_p - 1$. To do so, without loss of generality, we may assume $\alpha_1 \neq 0$. Then $\gamma_1 - 1$ acts on $L^+ \cdot \underline{T}^\alpha$ via

$$\zeta^\alpha \exp(-\theta_1(\zeta_p - 1)^2) - 1 = (\zeta^\alpha - 1) \left(1 + \zeta^\alpha \sum_{n \geq 1} \frac{(1 - \zeta_p)^n}{n!} \frac{(\zeta_p - 1)^n}{\zeta^\alpha - 1} \theta_1^n \right)$$

with θ_1 topologically nilpotent. As $1 + \zeta^\alpha \sum_{n \geq 1} \frac{(1 - \zeta_p)^n}{n!} \frac{(\zeta_p - 1)^n}{\zeta^\alpha - 1} \theta_1^n$ is invertible, we see that $H^i(\Gamma, L^+ \cdot \underline{T}^\alpha)$ is killed by $\zeta_p - 1$ for any i and vanishes for $i = 0$. Now one can conclude by using Hochschild–Serre spectral sequence. \square

We also need the following lemma.

Lemma 3.7. *For any $L^+ \in \text{Rep}_\Gamma^{\text{H-small}}(\mathcal{R})$ with $L_\infty^+ = L^+ \otimes_{\mathcal{R}} \widehat{R}_\infty^+$, the natural inclusion*

$$\text{R}\Gamma(\Gamma, L^+ \otimes_{\mathcal{R}} B^+) \rightarrow \text{R}\Gamma(\Gamma, L_\infty^+ \otimes_{\widehat{R}_\infty^+} B_\infty^+)$$

identifies the former with a direct summand of the latter whose complement is concentrated in degree ≥ 1 and killed by $\zeta_p - 1$. Moreover, let $\theta_i \in \text{End}_{\mathcal{R}}(L^+)$ be as in Definition 3.5(1), we have

$$(3.5) \quad \begin{aligned} H^+(L^+) &:= H^0(\Gamma, L \otimes_{\mathcal{R}} B^+) = \exp\left(\sum_{i=1}^d \theta_i W_i\right)(L^+) \\ &:= \{x \in L^+ \otimes_{\mathcal{R}} B^+ \mid x = \sum_{J \in \mathbb{N}^d} \underline{\theta}^J(y) \underline{W}^{[J]}, y \in L^+\}. \end{aligned}$$

with a natural isomorphism

$$(3.6) \quad H^+(L^+) \otimes_{\mathcal{R}} B^+ \xrightarrow{\cong} L^+ \otimes_{\mathcal{R}} B^+,$$

and that $H^n(\Gamma, L^+ \otimes_{\mathcal{R}} B^+)$ is killed by $(\zeta_p - 1)^2$ for any $n \geq 1$.

Proof. The decomposition (3.4) induces a Γ -equivariant decomposition

$$L_\infty^+ \otimes_{\widehat{R}_\infty^+} B_\infty^+ = \widehat{\bigoplus}_{\underline{\alpha}=(\alpha_1, \dots, \alpha_d) \in (\mathbb{N} \cap [0,1))^d} (L^+ \otimes_{\mathcal{R}} B^+) \cdot \underline{T}^\alpha.$$

For the first claim, it suffices to show that for any $\underline{\alpha} \neq 0$,

$$\text{R}\Gamma(\Gamma, (L^+ \otimes_{\mathcal{R}} B^+) \cdot \underline{T}^\alpha)$$

is concentrated in degree ≥ 1 and killed by $(\zeta_p - 1)^2$. But this follows from [MW23, Prop. 3.17(1)] by using Hochschild–Serre spectral sequence as in the proof of Lemma 3.6. For the “moreover” part follows from [MW23, Prop. 3.17(2)] directly. \square

Now, we are going to prove the following Simpson correspondence, which can be viewed as an analogue of Theorem 3.2 on the integral level.

Proposition 3.8. (1) *For any $L_\infty^+ \in \text{Rep}_\Gamma^{\text{H-small}}(\widehat{R}_\infty^+)$ of rank r , we have*

$$H^i(\Gamma, L_\infty^+ \otimes_{\widehat{R}_\infty^+} B_\infty^+) = \begin{cases} H^+(L_\infty^+), & i = 0 \\ (\zeta_p - 1)^2\text{-torsion}, & i \geq 1, \end{cases}$$

where $H^+(L_\infty^+)$ is a finite projective \mathcal{R} -module of rank r . The restriction of

$$\Theta_{L_\infty}^+ : L_\infty^+ \otimes_{\widehat{R}_\infty^+} B_\infty^+ \rightarrow L_\infty^+ \otimes_{\widehat{R}_\infty^+} B_\infty^+ \otimes_{\mathcal{R}} \Omega_{\mathcal{R}}^1\{-1\}$$

to $H^+(L_\infty^+)$ defines a Higgs field θ^+ on $H^+(L_\infty^+)$ making $(H^+(L_\infty^+), \theta^+) \in \text{HIG}^{\text{H-small}}(\mathcal{R})$.

(2) *For any $(H^+, \theta^+) \in \text{HIG}^{\text{H-small}}(\mathcal{R})$ of rank r , put*

$$\Theta_H^+ := \theta^+ \otimes \text{id} + \text{id} \otimes \Theta : H^+ \otimes_{\mathcal{R}} B_\infty^+ \rightarrow H^+ \otimes_{\mathcal{R}} B_\infty^+ \otimes_{\mathcal{R}} \Omega_{\mathcal{R}}^1\{-1\}.$$

Then $L_\infty^+(H^+, \theta^+) := (H^+ \otimes_{\mathcal{R}} B_\infty^+)^{\Theta_H^+ = 0}$ equipped with the induced Γ -action is a well-defined object in $\text{Rep}_\Gamma^{\text{H-small}}(\widehat{R}_\infty^+)$.

(3) *The functors $L_\infty^+ \mapsto (H^+(L_\infty^+), \theta^+)$ and $(H^+, \theta^+) \mapsto L_\infty^+(H^+, \theta^+)$ induces an equivalence of categories*

$$\text{Rep}_\Gamma^{\text{H-small}}(\widehat{R}_\infty^+) \simeq \text{HIG}^{\text{H-small}}(\mathcal{R})$$

which preserves ranks, tensor products and dualities. Moreover, for any $L_\infty^+ \in \text{Rep}_\Gamma^{\text{H-small}}(\widehat{R}_\infty^+)$ with corresponding $(H^+, \theta^+) \in \text{HIG}^{\text{H-small}}(\mathcal{R})$, we have a natural isomorphism

$$(3.7) \quad (L_\infty^+ \otimes_{\widehat{R}_\infty^+} B_\infty^+, \Theta_{L_\infty}^+) \cong (H^+ \otimes_{\mathcal{R}} B_\infty^+, \Theta_H^+)$$

compatible with Higgs fields, and a quasi-isomorphism

$$\text{R}\Gamma(\Gamma, L_\infty) \simeq \text{HIG}(H, \theta)$$

where $L_\infty = L_\infty^+[\frac{1}{p}]$ and $(H, \theta) = (H^+, \theta^+)[\frac{1}{p}]$.

Proof. For Item (1): Let $L^+ \in \text{Rep}_\Gamma^{\text{H-small}}(\mathcal{R})$ be the representation corresponding to L_∞^+ in the sense of Lemma 3.6. Let $\theta_i \in \text{End}_{\mathcal{R}}(L^+)$ be topologically nilpotent as in Definition 3.5(1). By Lemma 3.7, we have

$$H^+(L_\infty^+) = H^+(L^+) = \exp\left(\sum_{i=1}^d \theta_i W_i\right)(L^+)$$

is finite projective of rank r and θ^+ is the restriction of Θ_L^+ to $H^+(L^+)$. It then follows from (3.2) that

$$\theta^+ = \sum_{i=1}^d -(\zeta_p - 1)\theta_i \otimes (\zeta_p - 1) \frac{d\log T_i}{t}.$$

So $(H^+(L^+), \theta^+)$ is a well-defined object in $\text{HIG}^{\text{H-small}}(\mathcal{R})$.

Item (2) follows from Lemma 3.4.

For Item (3): Granting we have already obtained the desired equivalence, then (3.7) follows from (3.6). Note that Item (1) implies $H^i(\Gamma, L_\infty \otimes_{\widehat{R}_\infty} B_\infty) = 0$ for any $i \geq 1$. As the Higgs complex $\text{HIG}(B_\infty, \Theta)$ is a resolution of \widehat{R}_∞ by Theorem 2.4, we then have quasi-isomorphisms

$$\text{R}\Gamma(\Gamma, L_\infty) \simeq \text{R}\Gamma(\Gamma, \text{HIG}(L_\infty \otimes_{\widehat{R}_\infty} B_\infty, \Theta_{L_\infty})) \simeq \text{HIG}(H, \theta_H)$$

as desired. It remains to establish the desired equivalence.

For any $L_\infty^+ \in \text{Rep}_\Gamma^{\text{H-small}}(\widehat{R}_\infty^+)$, put $(H^+, \theta^+) = (H^+(L_\infty), \theta^+)$. Then we have a canonical morphism

$$\iota_{L_\infty^+} : L_\infty(H^+, \theta^+) \rightarrow L_\infty$$

compatible with Γ -actions defined by the composites

$$\begin{aligned} L_\infty(H^+, \theta^+) &= (H^+ \otimes_{\mathcal{R}} B_\infty^+)^{\Theta_H^+ := \theta^+ \otimes \text{id} + \text{id} \otimes \Theta = 0} \\ &\xrightarrow{=} ((L_\infty^+ \otimes_{\widehat{R}_\infty^+} B_\infty^+)^{\Gamma} \otimes_{\mathcal{R}} B_\infty^+)^{\Theta_L^+ \otimes \text{id} + (\text{id} \otimes \text{id}) \otimes \Theta = 0} \\ &\hookrightarrow (L_\infty^+ \otimes_{\widehat{R}_\infty^+} B_\infty^+ \otimes_{\widehat{R}_\infty^+} B_\infty^+)^{\text{id} \otimes \Theta \otimes \text{id} + \text{id} \otimes \text{id} \otimes \Theta = 0} \\ &\rightarrow (L_\infty^+ \otimes_{\widehat{R}_\infty^+} B_\infty^+)^{\text{id} \otimes \Theta = 0} \\ &= L_\infty. \end{aligned}$$

Here, the last but second arrow is induced by the multiplication on B_∞ .

For any $(H^+, \theta^+) \in \text{HIG}^{\text{H-small}}(\mathcal{R})$, put $L_\infty^+ = L_\infty^+(H^+, \theta^+)$. Then we have a canonical morphism

$$\iota_{(H^+, \theta^+)} : H^+(L_\infty^+) \rightarrow H^+$$

compatible with Higgs fields defined by the composites

$$\begin{aligned} H^+(L_\infty^+) &= (L_\infty^+ \otimes_{\widehat{R}_\infty^+} B_\infty^+)^{\Gamma} \\ &\xrightarrow{=} ((H^+ \otimes_{\mathcal{R}} B_\infty^+)^{\Theta_H^+ = 0} \otimes_{\widehat{R}_\infty^+} B_\infty^+)^{\Gamma} \\ &\hookrightarrow (H^+ \otimes_{\mathcal{R}} B_\infty^+ \otimes_{\widehat{R}_\infty^+} B_\infty^+)^{\Gamma} \\ &\rightarrow (H \otimes_{\mathcal{R}} B_\infty^+)^{\Gamma} \\ &= H^+. \end{aligned}$$

Here the last but second arrow is again induced by the multiplication on B_∞ while the last follows as $(B_\infty^+)^{\Gamma} = (B^+)^{\Gamma} = \mathcal{R}$ by Lemma 3.7. To conclude, it is enough to show both $\iota_{L_\infty^+}$ and $\iota_{(H^+, \theta^+)}$ are isomorphism. But this follows from the explicit description (3.3) and (3.5) immediately. \square

Now, we are able to give a criterion for being Hitchin-small v -vector bundles.

Proposition 3.9. *Let \mathfrak{X} be a liftable smooth formal scheme (not necessarily small affine) over A^+ with the fixed lifting $\widetilde{\mathfrak{X}}$ as before. Then an v -vector bundle \mathcal{L} on X_v is Hitchin-small if and only if there exists an étale covering $\{\mathfrak{X}_i \rightarrow \mathfrak{X}\}_{i \in I}$ by affine small $\mathfrak{X}_i = \text{Spf}(\mathcal{R}_i)$ with the corresponding $X_{i,\infty} = \text{Spa}(\widehat{R}_{i,\infty}^+, \widehat{R}_{i,\infty}^+)$ such that $\mathcal{L}(X_{i,\infty}) \in \text{Rep}_\Gamma^{\text{H-small}}(\widehat{R}_{i,\infty}^+)$ for all i .*

Proof. Recall for each affine small $\mathfrak{X}_i = \mathrm{Spf}(\mathcal{R}_i)$ with the generic fiber $X_i = \mathrm{Spa}(R_i, R_i^+ = \mathcal{R}_i)$, taking global sections induces an equivalence of categories

$$\mathrm{HIG}^{\mathrm{H-small}}(X_i)(A) \simeq \mathrm{HIG}^{\mathrm{H-small}}(R_i).$$

The result then follows from Corollary 3.3 and Proposition 3.8 immediately. \square

Remark 3.10. Thanks to existence of geometric Sen operator of Rodriguez-Carmago [RC22] (and of Heuer and Xu [HX24, Th. 3.2.1] in our setting), one can give another criterion for being a Hitchin-small v -vector bundle \mathcal{L} in terms of its geometric Sen operator $\phi_{\mathcal{L}}$, just like Proposition 3.1. However, this definition is not good enough for in our setting as up to now, there is no “geometric Sen operator” for general $\mathbb{B}_{\mathrm{dR},n}^+$ -local systems.

3.2. Local small Riemann–Hilbert correspondence. In this part, we always assume $\mathfrak{X} = \mathrm{Spf}(\mathcal{R})$ is small affine with the chart ψ over A^+ with the given lifting $\mathrm{Spf}(\tilde{\mathcal{R}})$ over $\mathbb{A}_{\mathrm{inf}}(A, A^+)$. Let $X = \mathrm{Spa}(R, R^+)$ be the generic fiber of \mathfrak{X} and \tilde{R} be the induced lifting of R over $\mathbb{B}_{\mathrm{dR}}^+(A, A^+)$. Then the lifting (cf. the paragraph above Lemma 2.5)

$$\tilde{\psi} : \mathbb{A}_{\mathrm{inf}}(A, A^+) \langle \underline{T}^{\pm 1} \rangle \rightarrow \tilde{\mathcal{R}}$$

induces a unique morphism

$$\tilde{\psi}_{\mathrm{dR}} : \tilde{R} \rightarrow \mathbb{B}_{\mathrm{dR}}^+(X_{\infty}).$$

Put $\tilde{\mathbf{B}}_{\psi,\infty} = \mathbb{B}_{\mathrm{dR}}^+(X_{\infty})$ and then it is equipped with a Γ -action such that for any $1 \leq i, j \leq d$,

$$\gamma_i([T_j^{\flat}]) = [\epsilon]^{\delta_{ij}} [T_j^{\flat}].$$

The $\tilde{\psi}_{\mathrm{dR}}$ identifies \tilde{R} with a Γ -stable sub- $\mathbb{B}_{\mathrm{dR}}^+(A, A^+)$ -algebra, denoted by $\tilde{\mathbf{B}}_{\psi}$, such that $\tilde{\mathbf{B}}_{\psi,\infty}$ admits a Γ -equivariant decomposition

$$\tilde{\mathbf{B}}_{\psi,\infty} = \bigoplus_{\underline{\alpha}=(\alpha_1,\dots,\alpha_d) \in (\mathbb{N}[1/p] \cap [0,1))^d} \tilde{\mathbf{B}}_{\psi} \cdot [T_1^{\flat\alpha_1}] \cdots [T_d^{\flat\alpha_d}].$$

Clearly, $\tilde{\mathbf{B}}_{\psi,\infty}/t = \hat{R}_{\infty}$ and $\tilde{\mathbf{B}}_{\psi}/t = R$. For simplicity, in what follows, we put

$$\tilde{\mathbf{B}}_{\psi,\infty}/t^{\infty} := \tilde{\mathbf{B}}_{\psi,\infty}, \tilde{\mathbf{B}}_{\psi}/t^{\infty} = \tilde{\mathbf{B}}_{\psi} \text{ and } \tilde{R}/t^{\infty} = \tilde{R}.$$

Definition 3.11. Fix an $1 \leq n \leq \infty$.

- (1) Let $C \in \{\tilde{\mathbf{B}}_{\psi,\infty}/t^n, \tilde{\mathbf{B}}_{\psi}/t^n\}$ By an *Hitchin-small representation* of Γ over C of rank r , we mean a finite projective C -module L endowed with an action of Γ such that its reduction $L/t \in \mathrm{Rep}_{\Gamma}^{\mathrm{H-small}}(C/t)$. Denote by $\mathrm{Rep}_{\Gamma}^{\mathrm{H-small}}(C)$ the category of Hitchin-small representation of Γ over C .
- (2) By a *integrable connection* over \tilde{R}/t^n of rank r , we mean a pair

$$(D, \nabla : D \rightarrow D \otimes_{\tilde{\mathcal{R}}} \Omega_{\tilde{\mathcal{R}}}^1 \{-1\})$$

satisfying $\nabla \wedge \nabla = 0$ such that D is a finite projective \tilde{R}/t^n -module of rank r and ∇ satisfies the Leibniz rule with respect to the derivation $d : \tilde{\mathcal{R}} \rightarrow \Omega_{\tilde{\mathcal{R}}}^1 \{-1\}$. An integrable connection (D, ∇) is called *Hitchin-small* if its reduction $(D, \nabla)/t \in \mathrm{HIG}^{\mathrm{H-small}}(R)$ modulo t . Denote by $\mathrm{MIC}^{\mathrm{H-small}}(\tilde{R}/t^n)$ the category of Hitchin-small integrable connection over \tilde{R}/t^n . For an integrable connection (D, ∇) , denoted by

$$\mathrm{DR}(D, \nabla) := [D \xrightarrow{\nabla} D \otimes_{\tilde{\mathcal{R}}} \Omega_{\tilde{\mathcal{R}}}^1 \{-1\} \rightarrow \cdots \rightarrow D \otimes_{\tilde{\mathcal{R}}} \Omega_{\tilde{\mathcal{R}}}^d \{-d\}]$$

the induced de Rham complex and put

$$\mathrm{H}_{\mathrm{dR}}^n(D, \nabla) := \mathrm{H}^n(\mathrm{DR}(D, \nabla)).$$

Recall we have deduce the following isomorphism from Proposition 2.8:

$$(\mathcal{O}_{\mathbb{B}_{\mathrm{dR},\mathrm{pd}}^+}, d)(X_{\infty}) = (\tilde{\mathbf{B}}_{\psi,\infty} \langle \underline{W} \rangle)_{\mathrm{pd}} = \tilde{\mathbf{B}}_{\psi,\infty} \langle W_1, \dots, W_d \rangle_{\mathrm{pd}}, \sum_{i=1}^d -u \frac{\partial}{\partial W_i} \otimes \frac{u}{t} d \log T_i$$

such that

$$\gamma_i(W_j) = W_j + \delta_{ij} u^2, \forall 1 \leq i, j \leq d.$$

Now, we can give the local Riemann–Hilbert correspondence for Hitchin-small representations of Γ over $\tilde{\mathbf{B}}_{\psi,\infty}/t^n$ and Hitchin-small integrable connections over \tilde{R}/t^n .

Proposition 3.12. *Fix $1 \leq n \leq \infty$.*

- (1) *For any $L_\infty \in \text{Rep}_\Gamma^{\text{H-small}}(\tilde{\mathbf{B}}_{\psi,\infty}/t^n)$ of rank r , we have*

$$H^n(\Gamma, L_\infty \otimes_{\tilde{\mathbf{B}}_{\psi,\infty}} \tilde{\mathbf{B}}_{\psi,\infty} \langle \underline{W} \rangle_{\text{pd}}) = \begin{cases} D(M_\infty), & n = 0 \\ 0, & n \geq 1, \end{cases}$$

where $D(L_\infty)$ is a finite free \tilde{R}/t^n -module of rank r . Moreover, the restriction of d to $D(L_\infty)$ induces a flat connection ∇ on $D(L_\infty)$ making it an object in $\text{MIC}^{\text{H-small}}(\tilde{R}/t^n)$.

- (2) *For any $(D, \nabla) \in \text{MIC}^{\text{H-small}}(\tilde{R}/t^n)$ of rank r , define*

$$\nabla_D = \nabla \otimes \text{id} + \text{id} \otimes \nabla : D \otimes_{\tilde{R}} \tilde{\mathbf{B}}_{\psi,\infty} \langle \underline{W} \rangle_{\text{pd}} \rightarrow D \otimes_{\tilde{R}} \tilde{\mathbf{B}}_{\psi,\infty} \langle \underline{W} \rangle_{\text{pd}} \otimes_{\tilde{R}} \Omega_{\tilde{R}}^1\{-1\}.$$

Then it satisfies $\nabla_D \wedge \nabla_D = 0$ and we have

$$H_{\text{dR}}^n(D \otimes_{\tilde{R}} \tilde{\mathbf{B}}_{\psi,\infty} \langle \underline{W} \rangle_{\text{pd}}, \nabla_D) = \begin{cases} L_\infty(D, \nabla), & n = 0 \\ 0, & n \geq 1, \end{cases}$$

where $L_\infty(D, \nabla)$ is a finite free $\tilde{\mathbf{B}}_{\psi,\infty}/t^n$ -module of rank r . Moreover, the Γ -action on $\tilde{\mathbf{B}}_{\psi,\infty} \langle \underline{W} \rangle_{\text{pd}}$ induces a Γ -action on $L_\infty(D, \nabla)$ making it an object in $\text{Rep}_\Gamma^{\text{H-small}}(\tilde{\mathbf{B}}_{\psi,\infty}/t^n)$.

- (3) *The functors $L_\infty \mapsto (D(L_\infty), \nabla)$ and $(D, \nabla) \mapsto L_\infty(D, \nabla)$ induce an equivalence of categories*

$$\text{Rep}_\Gamma^{\text{H-small}}(\tilde{\mathbf{B}}_{\psi,\infty}/t^n) \simeq \text{MIC}^{\text{H-small}}(\tilde{R}/t^n)$$

preserving ranks, tensor products and dualities. Moreover for any $L_\infty \in \text{Rep}_\Gamma^{\text{H-small}}(\tilde{\mathbf{B}}_{\psi,\infty}/t^n)$ corresponding to (D, ∇) , we have a quasi-isomorphism

$$\text{R}\Gamma(\Gamma, L_\infty) \simeq \text{DR}(D, \nabla).$$

Before proving Proposition 3.12, we first give some lemmas.

Lemma 3.13 ([CLWZ24, Lem. 6.10]). *Let B be a ring with a non-zero divisor t and $B_m := B/t^m$ for any $m \geq 1$.*

- (1) *Let M be a B_m -module. If M/tM is a finite free B_1 -module of rank r and for any $0 \leq n \leq m-1$, the multiplication by t^n induces an isomorphism of B_1 -modules*

$$M/tM \xrightarrow{\cong} t^n M/t^{n+1} M,$$

then M is a finite free B_m -module of rank r .

- (2) *Assum moreover B is t -adically complete. Let M be a B -module such that M/tM is a finite free B_1 -module of rank r and for any $n \geq 0$, the multiplication by t^n induces an isomorphism of B_1 -modules*

$$M/tM \xrightarrow{\cong} t^n M/t^{n+1} M,$$

then M is a finite free B -module of rank r .

Proof. The Item (2) is a consequence of Item (1) together with [MT20, Lem. 1.9]. So it is enough to prove Item (1).

Let e_1, \dots, e_r be elements in M whose reductions modulo t induces an B_1 -basis of M/tM . We claim that e_1, \dots, e_r form a B_m -basis of M . We now prove this by induction on m . We may assume $m > 1$ and then by inductive hypothesis, the reduction of e_1, \dots, e_r modulo t^{m-1} form a B_{m-1} -basis of $M/t^{m-1}M$. Thus, for any $x \in M$, there exist $b_1, \dots, b_r \in B_m$ such that

$$x \equiv b_1 e_1 + \dots + b_r e_r \pmod{t^{m-1}M}.$$

As $t^{m-1}M \cong M/tM$, one can find $c_1, \dots, c_r \in B_m$ such that

$$x - (b_1 e_1 + \dots + b_r e_r) = t^{m-1} c_1 e_1 + \dots + t^{m-1} c_r e_r.$$

Put $a_i = b_i + t^{m-1} c_i$ for any $1 \leq i \leq r$ and then we have

$$x = a_1 e_1 + \dots + a_r e_r.$$

So e_1, \dots, e_r generate M over B_m . To conclude, it remains to show for any $d_1, \dots, d_r \in B_m$ such that

$$d_1 e_1 + \dots + d_r e_r = 0,$$

we must have

$$d_1 = d_2 = \dots = d_r = 0.$$

By inductive hypothesis, we have

$$d_1 \equiv d_2 \equiv \dots \equiv d_r \pmod{t^{m-1}}.$$

That is, there are $f_1, \dots, f_r \in B_m$ such that $d_i = t^{m-1} f_i$ for any $1 \leq i \leq r$. Thus, we have

$$f_1 t^{m-1} e_1 + \dots + f_r t^{m-1} e_r = 0.$$

Using $t^{m-1} M \cong M/tM$ again, we conclude that

$$f_1 \equiv f_2 \equiv \dots \equiv f_r \pmod{t},$$

which forces that for any $1 \leq i \leq r$,

$$d_i = t^{m-1} f_i = 0$$

as desired. This completes the proof. \square

Lemma 3.14. (1) Fix $B \in \{\tilde{\mathbf{B}}_{\psi, \infty}, \tilde{\mathbf{B}}_{\psi}\}$. Then the functor $M \mapsto \{M/t^n\}_{n \geq 1}$ induces an equivalence of categories

$$\mathrm{Rep}_{\Gamma}^{\mathrm{H-small}}(B) \simeq \varprojlim_n \mathrm{Rep}_{\Gamma}^{\mathrm{H-small}}(B/t^n)$$

which is compatible with cohomologies; that is, we have a canonical quasi-isomorphism

$$\mathrm{R}\Gamma(\Gamma, M) \xrightarrow{\sim} \varprojlim_n \mathrm{R}\Gamma(\Gamma, M/t^n).$$

(2) Then the functor $(D, \nabla) \mapsto (D/t^n, \nabla)$ induces an equivalence of categories

$$\mathrm{MIC}^{\mathrm{H-small}}(\tilde{R}) \simeq \varprojlim_n \mathrm{MIC}^{\mathrm{H-small}}(\tilde{R}/t^n),$$

which is compatible with cohomologies; that is, we have a canonical quasi-isomorphism

$$\mathrm{DR}(D, \nabla) = \varprojlim_n \mathrm{DR}(D/t^n \nabla)$$

Proof. Note that $\tilde{\mathbf{B}}_{\psi}$, $\tilde{\mathbf{B}}_{\psi, \infty}$ and \tilde{R} are all t -adically complete and t -torsion free. The desired equivalences follows immediately from Lemma 3.13(2) while the cohomological comparison follows from the (derived) Nakayama's Lemma. \square

Proof of Proposition 3.12: By Lemma 3.14, it suffices to deal with the case where $n < \infty$. Note that the $n = 1$ case has been established in Proposition 3.8. We will finish the proof by induction on n . From now on, assume there exists some $n \geq 1$ such that the result holds true for any $1 \leq m \leq n$.

For Item (1): Fix an $M_{\infty} \in \mathrm{Rep}_{\Gamma}^{\mathrm{H-small}}(\tilde{\mathbf{B}}_{\psi, \infty}/t^{n+1})$ of rank r . Then we have

$$tM_{\infty} \in \mathrm{Rep}_{\Gamma}^{\mathrm{H-small}}(\tilde{\mathbf{B}}_{\psi, \infty}/t^n) \text{ and } M_{\infty}/t \in \mathrm{Rep}_{\Gamma}^{\mathrm{H-small}}(\tilde{\mathbf{B}}_{\psi, \infty}/t).$$

The short exact sequence

$$0 \rightarrow tM_{\infty} \rightarrow M_{\infty} \rightarrow M_{\infty}/t \rightarrow 0$$

gives rise to an exact triangle

$$\mathrm{R}\Gamma(\Gamma, tM_{\infty} \otimes_{\tilde{\mathbf{B}}_{\psi, \infty}} \tilde{\mathbf{B}}_{\psi, \infty} \langle \underline{W} \rangle_{\mathrm{pd}}) \rightarrow \mathrm{R}\Gamma(\Gamma, M_{\infty} \otimes_{\tilde{\mathbf{B}}_{\psi, \infty}} \tilde{\mathbf{B}}_{\psi, \infty} \langle \underline{W} \rangle_{\mathrm{pd}}) \rightarrow \mathrm{R}\Gamma(\Gamma, M_{\infty}/t \otimes_{\tilde{\mathbf{B}}_{\psi, \infty}} \tilde{\mathbf{B}}_{\psi, \infty} \langle \underline{W} \rangle_{\mathrm{pd}}).$$

Considering the induced long exact sequence, by inductive hypothesis, we have

$$\mathrm{H}^n(\Gamma, M_{\infty} \otimes_{\tilde{\mathbf{B}}_{\psi, \infty}} \tilde{\mathbf{B}}_{\psi, \infty} \langle \underline{W} \rangle_{\mathrm{pd}}) = 0$$

for any $n \geq 1$ and there is a short exact sequence

$$0 \rightarrow D_{\mathrm{RH}}(tM_{\infty}) \rightarrow D_{\mathrm{RH}}(M_{\infty}) \rightarrow D_{\mathrm{RH}}(M_{\infty}/t) \rightarrow 0.$$

We claim that

$$D_{\mathrm{RH}}(tM_{\infty}) = tD_{\mathrm{RH}}(M_{\infty}) \text{ and } D_{\mathrm{RH}}(M_{\infty}/t) = D_{\mathrm{RH}}(M_{\infty})/t.$$

Indeed, consider the exact sequence

$$\cdots \rightarrow M_\infty \xrightarrow{\times t} M_\infty \xrightarrow{\times t^n} M_\infty \xrightarrow{\times t} M_\infty \rightarrow M_\infty/t \rightarrow 0.$$

Applying $\mathrm{R}\Gamma(\Gamma, - \otimes_{\tilde{\mathbf{B}}_{\psi, \infty}} \tilde{\mathbf{B}}_{\psi, \infty} \langle W \rangle_{\mathrm{pd}})$ to the above sequence, we obtain an exact sequence

$$\cdots \rightarrow D_{\mathrm{RH}}(M_\infty) \xrightarrow{\times t} D_{\mathrm{RH}}(M_\infty) \xrightarrow{\times t^n} D_{\mathrm{RH}}(M_\infty) \xrightarrow{\times t} D_{\mathrm{RH}}(M_\infty) \rightarrow D_{\mathrm{RH}}(M_\infty/t) \rightarrow 0.$$

So we have the short exact sequence

$$0 \rightarrow tD_{\mathrm{RH}}(M_\infty) \rightarrow D_{\mathrm{RH}}(M_\infty) \rightarrow D_{\mathrm{RH}}(M_\infty/t) \rightarrow 0.$$

Thus the claim follows.

A similar argument implies that for any $0 \leq m \leq n$, we have $D_{\mathrm{RH}}(t^m M_\infty) \cong t^m D_{\mathrm{RH}}(M_\infty)$, yielding that

$$t^m D_{\mathrm{RH}}(M_\infty)/t^{m+1} D_{\mathrm{RH}}(M_\infty) \cong D_{\mathrm{RH}}(t^m M_\infty/t^{m+1} M_\infty) \cong D_{\mathrm{RH}}(M_\infty/t).$$

By Lemma 3.13, we see that $D_{\mathrm{RH}}(M_\infty)$ is finite free of rank r over \tilde{R}/t^{n+1} as desired. Finally, as

$$(D_{\mathrm{RH}}(M_\infty/t), \nabla) = (D_{\mathrm{RH}}(M_\infty)/t, \nabla) \in \mathrm{MIC}^{\mathrm{H-small}}(\tilde{R}/t)$$

we have $(D_{\mathrm{RH}}(M_\infty/t), \nabla) \in \mathrm{MIC}^{\mathrm{H-small}}(\tilde{R}/t^{n+1})$ as desired. This completes the proof of Item (1).

Item (2) can be deduced from the similar argument above.

For Item (3): Similar to the proof of Proposition 3.8 (3), for any $M_\infty \in \mathrm{Rep}_\Gamma^{\mathrm{H-small}}(\tilde{\mathbf{B}}_{\psi, \infty}/t^n)$ and any $(D, \nabla) \in \mathrm{MIC}^{\mathrm{H-small}}(\tilde{R}/t^{n+1})$, one can construct natural morphisms

$$\iota_{M_\infty} : M_\infty(D_{\mathrm{RH}}(M_\infty), \nabla) \rightarrow M_\infty$$

and

$$\iota_{(D, \nabla)} : (D_{\mathrm{RH}}(M_\infty(D, \nabla)), \nabla) \rightarrow (D, \nabla).$$

To see they are both isomorphism, by Nakayama's Lemma, one can check this modulo t , and thus reduces to Theorem Proposition 3.8 (3). This establishes the desired equivalence of categories. The cohomological comparison follows easily. \square

To complete the local theory, we state the following result, which will not be used in this paper.

Corollary 3.15. *For any $1 \leq n \leq \infty$, the base-change $L \mapsto L_\infty := L \otimes_{\tilde{\mathbf{B}}_\psi} \tilde{\mathbf{B}}_{\psi, \infty}$ induces an equivalence of categories*

$$\mathrm{Rep}_\Gamma^{\mathrm{H-small}}(\tilde{\mathbf{B}}_\psi/t^n) \xrightarrow{\sim} \mathrm{Rep}_\Gamma^{\mathrm{H-small}}(\tilde{\mathbf{B}}_{\psi, \infty}/t^n)$$

which is compatible with cohomologies; that is, we have a quasi-isomorphism

$$\mathrm{R}\Gamma(\Gamma, L) \cong \mathrm{R}\Gamma(\Gamma, L_\infty).$$

Proof. The full faithfulness together with the cohomological comparison follows from the $n = 1$ case (cf. Lemma 3.6) together with the derived Nakayama's Lemma. It remains to show the essential surjectivity. Note that

$$\tilde{\mathbf{B}}_\psi \langle W \rangle_{\mathrm{pd}} \subset \tilde{\mathbf{B}}_{\psi, \infty} \langle W \rangle_{\mathrm{pd}}$$

is a d -preserving sub-ring stable under the action of Γ . For any $L_\infty \in \mathrm{Rep}_\Gamma^{\mathrm{H-small}}(\tilde{\mathbf{B}}_{\psi, \infty}/t^n)$ with the induced $(D, \nabla) \in \mathrm{MIC}^{\mathrm{H-small}}(\tilde{R}/t^n)$, one can achieve an $L \in \mathrm{Rep}_\Gamma^{\mathrm{H-small}}(\tilde{\mathbf{B}}_\psi/t^n)$ by using $\tilde{\mathbf{B}}_\psi \langle W \rangle_{\mathrm{pd}}$ instead of $\tilde{\mathbf{B}}_{\psi, \infty} \langle W \rangle_{\mathrm{pd}}$ in Proposition 3.12. Then one can conclude by checking $L \otimes_{\tilde{\mathbf{B}}_\psi} \tilde{\mathbf{B}}_{\psi, \infty} \cong L_\infty$ directly. \square

4. THE STACKY RIEMANN–HILBERT CORRESPONDENCE: PROOF OF THEOREM 1.1

This section is devoted to proving Theorem 1.1. To do so, we need the following result:

Theorem 4.1. *Fix a $\mathrm{Spa}(A, A^+) \in \mathrm{Perfd}$. Let \mathfrak{X} be a liftable smooth formal scheme over A^+ with a fixed lifting $\tilde{\mathfrak{X}}$ over $\mathbb{A}_{\mathrm{inf}}(A, A^+)$. Let X be the generic fiber of \mathfrak{X} and \tilde{X} be the lifting of X over $\mathbb{B}_{\mathrm{dR}}^+(A, A^+)$ induced by $\tilde{\mathfrak{X}}$. Let $(\mathcal{O}_{\mathrm{dR}, \mathrm{pd}}^+, d)$ be the period sheaf with connection constructed in §2. Let $\nu : X_v \rightarrow X_{\mathrm{\acute{e}t}}$ be the natural morphism of sites. Let $1 \leq n \leq \infty$.*

(1) For any $\mathbb{L} \in \mathrm{LS}^{\mathrm{H-small}}(X, \mathbb{B}_{\mathrm{dR},n}^+)(A)$ of rank r , we have

$$R^n \nu_* (\mathbb{L} \otimes_{\mathbb{B}_{\mathrm{dR}}^+} \mathcal{O}\mathbb{B}_{\mathrm{dR,pd}}^+) = \begin{cases} \mathcal{D}(\mathbb{L}), & n = 0 \\ 0, & n \geq 1 \end{cases}$$

where $\mathcal{D}(\mathbb{L})$ is a locally finite free $(\mathcal{O}_{\tilde{X}_n})$ -module of rank r on $X_{\mathrm{ét}}$ such that the

$$\mathrm{id}_{\mathbb{L}} \otimes d : \mathbb{L} \otimes_{\mathbb{B}_{\mathrm{dR}}^+} \mathcal{O}\mathbb{B}_{\mathrm{dR,pd}}^+ \rightarrow \mathbb{L} \otimes_{\mathbb{B}_{\mathrm{dR}}^+} \mathcal{O}\mathbb{B}_{\mathrm{dR,pd}}^+ \otimes_{\mathcal{O}_{\tilde{X}}} \Omega_{\tilde{X}}^1 \{-1\}$$

induces a flat connection $\nabla_{\mathbb{L}}$ on $\mathcal{D}(\mathbb{L})$ making $(\mathcal{D}(\mathbb{L}), \nabla_{\mathbb{L}})$ an object in $\mathrm{MIC}^{\mathrm{H-small}}(\tilde{X}_n)(A)$.

(2) For any $(\mathcal{D}, \nabla) \in \mathrm{MIC}^{\mathrm{H-small}}(\tilde{X}_n)$ of rank r , define

$$\nabla_{\mathcal{D}} := \nabla \otimes \mathrm{id} + \mathrm{id} \otimes d : \mathcal{D} \otimes_{\mathcal{O}_{\tilde{X}}} \mathcal{O}\mathbb{B}_{\mathrm{dR,pd}}^+ \rightarrow \mathcal{D} \otimes_{\mathcal{O}_{\tilde{X}}} \mathcal{O}\mathbb{B}_{\mathrm{dR,pd}}^+ \otimes_{\mathcal{O}_{\tilde{X}}} \Omega_{\tilde{X}}^1 \{-1\}.$$

Then

$$\mathbb{L}(\mathcal{D}, \nabla) := (\mathcal{D} \otimes_{\mathcal{O}_{\tilde{X}}} \mathcal{O}\mathbb{B}_{\mathrm{dR,pd}}^+)^{\nabla_{\mathcal{D}}=0}$$

is an object of rank r in $\mathrm{LS}^{\mathrm{H-small}}(X, \mathbb{B}_{\mathrm{dR},n}^+)(A)$.

(3) The functors $\mathbb{L} \mapsto (\mathcal{D}(\mathbb{L}), \nabla_{\mathbb{L}})$ and $(\mathcal{D}, \nabla) \mapsto \mathbb{L}(\mathcal{D}, \nabla)$ in Items (1) and (2) respectively define an equivalence of categories

$$\rho_{\tilde{X}} : \mathrm{LS}^{\mathrm{H-small}}(\mathfrak{X}, \mathbb{B}_{\mathrm{dR},n}^+)(A) \xrightarrow{\sim} \mathrm{MIC}^{\mathrm{H-small}}(\tilde{X}_n)(A)$$

which preserves ranks, tensor products and dualities. Moreover, for any $\mathbb{L} \in \mathrm{LS}^{\mathrm{H-small}}(X, \mathbb{B}_{\mathrm{dR},n}^+)$ with corresponding $(\mathcal{D}, \nabla) \in \mathrm{MIC}^{\mathrm{H-small}}(\tilde{X}_n)(A)$, there exists a quasi-isomorphism

$$R\nu_* \mathbb{L} \simeq \mathrm{DR}(\mathcal{D}, \nabla).$$

In particular, we have a quasi-isomorphism

$$\mathrm{R}\Gamma(X_v, \mathbb{L}) \simeq \mathrm{R}\Gamma(X_{\mathrm{ét}}, \mathrm{DR}(\mathcal{D}, \nabla)).$$

Before proving this theorem, we explain how to use it to obtain Theorem 1.1.

Proof of Theorem 1.1: Let \mathfrak{X} be a liftable smooth formal scheme over \mathcal{O}_C with the fixed lifting $\tilde{\mathfrak{X}}$ over $\mathbf{A}_{\mathrm{inf}}$, the generic fiber X and the lifting \tilde{X} of X over $\mathbf{B}_{\mathrm{dR}}^+$. For any $\mathrm{Spa}(A, A^+) \in \mathrm{Perfd}$, we denote by \mathfrak{X}_A , $\tilde{\mathfrak{X}}_A$, X_A and \tilde{X}_A their corresponding base-changes to A^+ , $\mathbf{A}_{\mathrm{inf}}(A, A^+)$, A and $\mathbb{B}_{\mathrm{dR}}^+(A, A^+)$. Let $(\mathcal{O}\mathbb{B}_{\mathrm{dR,pd},A}^+, d)$ be the period sheaf with connection on $X_{A,v}$ corresponding to the lifting $\tilde{\mathfrak{X}}_A$. Then it follows from Proposition 2.9 that for any $f : \mathrm{Spa}(B, B^+) \rightarrow \mathrm{Spa}(A, A^+)$, we have an isomorphism

$$\mathcal{O}\mathbb{B}_{\mathrm{dR,pd},B}^+ = \mathcal{O}\mathbb{B}_{\mathrm{dR,pd},A}^+ \hat{\otimes}_{\mathbb{B}_{\mathrm{dR}}^+(A, A^+)} \mathbb{B}_{\mathrm{dR}}^+(B, B^+)$$

compatible with connection. Note that f induces the obvious base-change functors

$$f_{\mathrm{LS}} : \mathrm{LS}(X, \mathbb{B}_{\mathrm{dR},n}^+)^{\mathrm{H-small}}(A) \rightarrow \mathrm{LS}(X, \mathbb{B}_{\mathrm{dR},n}^+)^{\mathrm{H-small}}, \quad \mathbb{L} \mapsto \mathbb{L} \hat{\otimes}_{\mathbb{B}_{\mathrm{dR}}^+(A, A^+)} \mathbb{B}_{\mathrm{dR}}^+(B, B^+)$$

and

$$f_{\mathrm{MIC}} : \mathrm{MIC}(\tilde{X}_n)^{\mathrm{H-small}}(A) \rightarrow \mathrm{MIC}^{\mathrm{H-small}}(\tilde{X}_n)(B), \quad (\mathcal{D}, \nabla) \mapsto (\mathcal{D} \hat{\otimes}_{\mathbb{B}_{\mathrm{dR}}^+(A, A^+)} \mathbb{B}_{\mathrm{dR}}^+(B, B^+), \nabla \otimes \mathrm{id}).$$

As the construction in Theorem 4.1 is clearly functorial in $\mathrm{Spa}(A, A^+)$, we have

$$f_{\mathrm{MIC}} \circ \rho_{\tilde{\mathfrak{X}}_A} = \rho_{\tilde{\mathfrak{X}}_B} \circ f_{\mathrm{LS}}.$$

As $\rho_{\tilde{\mathfrak{X}}_A}$ preserves ranks, the equivalence criterion of $\rho_{\tilde{\mathfrak{X}}_A}$ yields the desired equivalence of stacks

$$\rho_{\tilde{X}} : \mathrm{LS}_r(X, \mathbb{B}_{\mathrm{dR},n}^+)^{\mathrm{H-small}} \xrightarrow{\sim} \mathrm{MIC}_r(\tilde{X}_n)^{\mathrm{H-small}}.$$

□

Now, we focus on the proof of Theorem 4.1 by proceeding as in the proof of [CLWZ24, Th. 7.11].

Lemma 4.2. Suppose that \mathfrak{X} is small affine and let $\mathbb{L} \in \mathrm{LS}(X, \mathbb{B}_{\mathrm{dR},n}^+)(A)$. Then for any affinoid perfectoid $U = \mathrm{Spa}(S, S^+) \in X_{\infty,v}$ and for any $i \geq 1$, we have

$$H^i(U, \mathbb{L} \otimes_{\mathbb{B}_{\mathrm{dR}}^+} \mathcal{O}\mathbb{B}_{\mathrm{dR,pd}}^+) = 0.$$

Proof. We need to show that $R\Gamma(U, \mathbb{L} \otimes_{\mathbb{B}_{\text{dR}}^+} \mathcal{O}\mathbb{B}_{\text{dR}, \text{pd}}^+)$ is concentrated in degree 0. By derived Nakayama's Lemma, we are reduced to showing the case for $n = 1$. That is, we have to show for any v -vector bundle \mathbb{L} ,

$$R\Gamma(U, \mathbb{L} \otimes_{\widehat{\mathcal{O}}_X} \widehat{\mathcal{O}}_{\widehat{\mathbb{C}}_{\text{pd}}})$$

is concentrated in degree 0. It follows from [KL16, Th. 3.5.8] that there is a finite projective S -module L such that $\mathbb{L}|_U \cong L \otimes_{S^+} \widehat{\mathcal{O}}_U$. So without loss of generality, we may assume $\mathbb{L}|_U = \widehat{\mathcal{O}}_U$, and are reduced to showing that

$$R\Gamma(U, \mathcal{O}\widehat{\mathbb{C}}_{\text{pd}}) = R\Gamma(U, \widehat{\mathcal{O}}_U[W]_{\text{pd}}^\wedge)$$

is concentrated in degree 0, by using Proposition 2.8. It suffices to show that

$$R\Gamma(U, \widehat{\mathcal{O}}_U^+[W]_{\text{pd}}^\wedge)$$

has cohomologies killed by \mathfrak{m}_C in degree ≥ 1 . However, as $\widehat{\mathcal{O}}_U^+[W]_{\text{pd}}^\wedge$ is the p -adic completion of a directly limit of finite free $\widehat{\mathcal{O}}_U^+$ -modules, by the quasi-compactness of U , we are reduced to show that

$$R\Gamma(U, \widehat{\mathcal{O}}_U^+)$$

has cohomologies killed by \mathfrak{m}_C in degree ≥ 1 . This is well-known (cf. [Sch17, Prop. 8.8]). \square

Lemma 4.3. *Suppose that \mathfrak{X} is small affine and let $\mathbb{L} \in \text{LS}(X, \mathbb{B}_{\text{dR}, n}^+)(A)$. Then there exists a natural quasi-isomorphism*

$$R\Gamma(\Gamma, \mathbb{L} \otimes_{\mathbb{B}_{\text{dR}}^+} \mathcal{O}\mathbb{B}_{\text{dR}, \text{pd}}^+(X_\infty)) \rightarrow R\Gamma(X_v, \mathbb{L} \otimes_{\mathbb{B}_{\text{dR}}^+} \mathcal{O}\mathbb{B}_{\text{dR}, \text{pd}}^+).$$

Proof. This follows from the same argument for the proof of [Wan23, Lem. 5.11] by using Lemma 4.2 instead of [Wan23, Lem. 5.7]. \square

Now, we are prepared to show Theorem 4.1.

Proof of Theorem 4.1: For Item (1): Fix an $\mathbb{L} \in \text{LS}^{\text{H-small}}(X, \mathbb{B}_{\text{dR}, n}^+)(A)$ of rank r . Since the problem is local on $\mathfrak{X}_{\text{ét}}$, we may assume $\mathfrak{X} = \text{Spf}(\mathcal{R})$ is small affine. According to Remark 1.4 and Proposition 3.9, comparing Definition 3.5(2) with Definition 3.11(1), we have $\mathbb{L}(X_\infty) \in \text{Rep}_\Gamma^{\text{H-small}}(\widetilde{\mathcal{B}}_{\psi, \infty}/t^n)$. Then one can deduce Item (1) from Lemma 4.3 together with Proposition 3.12(1).

For Item (2): Fix an $(\mathcal{D}, \nabla) \in \text{MIC}^{\text{H-small}}(\widetilde{X}_n)(A)$. Since the problem is again local on $\mathfrak{X}_{\text{ét}}$, we may assume $\mathfrak{X} = \text{Spf}(\mathcal{R})$ is small affine. According to Remark 1.4 and Proposition 3.1, comparing Definition 3.5(3) with Definition 3.11(2), we see that the global section

$$(D, \nabla) := (\mathcal{D}, \nabla)(X) \in \text{MIC}^{\text{H-small}}(\widetilde{R}/t^n).$$

Then one can deduce Item (2) from Proposition 3.12(2).

For Item (3): For any $\mathbb{L} \in \text{LS}^{\text{H-small}}(X, \mathbb{B}_{\text{dR}, n}^+)(A)$ and any $(\mathcal{D}, \nabla) \in \text{MIC}^{\text{H-small}}(\widetilde{X}_n)(A)$, similar to the proof of Proposition 3.12(3), one can construct two canonical morphisms

$$\iota_{\mathbb{L}} : \mathbb{L}(\mathcal{D}(\mathbb{L}), \nabla_{\mathbb{L}}) \rightarrow \mathbb{L}$$

and

$$\iota_{(\mathcal{D}, \nabla)} : (\mathcal{D}(\mathbb{L}(\mathcal{D}, \nabla)), \nabla_{\mathbb{L}(\mathcal{D}, \nabla)}) \rightarrow (\mathcal{D}, \nabla).$$

To get desired equivalence of categories, we have to show both $\iota_{\mathbb{L}}$ and $\iota_{(\mathcal{D}, \nabla)}$ are isomorphisms. But this is still a local problem and thus reduces to the proof of Proposition 3.12(3). Finally, we have to show that for any $\mathbb{L} \in \text{LS}^{\text{H-small}}(X, \mathbb{B}_{\text{dR}, n}^+)(A)$ with corresponding $(\mathcal{D}, \nabla) \in \text{MIC}^{\text{H-small}}(\widetilde{X}_n)(A)$, we have

$$R\nu_* \mathbb{L} \simeq \text{DR}(\mathcal{D}, \nabla).$$

By Poincaré's Lemma (cf. Theorem 2.4), we have

$$R\nu_* \mathbb{L} \simeq R\nu_*(\text{DR}(\mathbb{L} \otimes_{\mathbb{B}_{\text{dR}}^+} \mathcal{O}\mathbb{B}_{\text{dR}, \text{pd}}^+, \text{id}_{\mathbb{L}} \otimes d)).$$

It follows from Item (1) that

$$R\nu_*(\text{DR}(\mathbb{L} \otimes_{\mathbb{B}_{\text{dR}}^+} \mathcal{O}\mathbb{B}_{\text{dR}, \text{pd}}^+, \text{id}_{\mathbb{L}} \otimes d)) \simeq \nu_*(\text{DR}(\mathbb{L} \otimes_{\mathbb{B}_{\text{dR}}^+} \mathcal{O}\mathbb{B}_{\text{dR}, \text{pd}}^+, \text{id}_{\mathbb{L}} \otimes d)) \simeq \text{DR}(\mathcal{D}, \nabla_{\mathcal{D}}).$$

This completes the proof. \square

REFERENCES

- [AHLB23a] Johannes Anschütz, Ben Heuer, and Arthur-César Le Bras. Hodge-Tate stacks and non-abelian p -adic Hodge theory of v -perfect complexes on smooth rigid spaces. *arXiv preprint arXiv:2302.12747*, 2023.
- [AHLB23b] Johannes Anschütz, Ben Heuer, and Arthur-César Le Bras. The small p -adic Simpson correspondence in terms of moduli spaces. *arXiv preprint arXiv:2312.07554*, 2023.
- [Bha21] Bhargav Bhatt. Algebraic geometry in mixed characteristic. *arXiv preprint arXiv:2112.12010*, 2021.
- [BL22] Bhargav Bhatt and Jacob Lurie. The prismatization of p -adic formal schemes. *arXiv:2201.06124*, 2022.
- [CLWZ24] Zekun Chen, Ruochuan Liu, Yupeng Wang, and Xinwen Zhu. A p -adic riemann–hilbert correspondence over robba rings. *In preparation*, 2024.
- [DLLZ23] Hansheng Diao, Kai-Wen Lan, Ruochuan Liu, and Xinwen Zhu. Logarithmic Riemann–Hilbert correspondences for rigid varieties. *Journal of the American Mathematical Society*, 36(2):483–562, 2023.
- [Fal05] Gerd Faltings. A p -adic Simpson correspondence. *Advances in Mathematics*, 198(2):847–862, 2005.
- [GMW] Hui Gao, Yu Min, and Yupeng Wang. On de Rham crystals in the relative case, tentative title. *in preparation*.
- [Heu22] Ben Heuer. Moduli spaces in p -adic non-abelian Hodge theory. *arXiv preprint arXiv:2207.13819*, 2022.
- [Heu23] Ben Heuer. A p -adic Simpson correspondence for smooth proper rigid varieties. *arXiv preprint arXiv:2307.01303*, 2023.
- [HX24] Ben Heuer and Daxin Xu. p -adic non-abelian hodge theory for curves via moduli stacks. *arXiv preprint arxiv:2402.01365*, 2024.
- [KL16] Kiran S Kedlaya and Ruochuan Liu. Relative p -adic hodge theory, II: Imperfect period rings. *arXiv preprint arXiv:1602.06899*, 2016.
- [Li] Yixiao Li. p -adic riemann–hilbert correspondence for zariski \mathbb{Q}_p -constructible sheaves. *In preparation*.
- [LQN24] Yudong Liu, Xiaoyu Qu, and Xiecheng Nie. A stacky nilpotent Riemann–Hilbert correspondence. *In preparation*, 2024.
- [LZ17] Ruochuan Liu and Xinwen Zhu. Rigidity and a Riemann–Hilbert correspondence for p -adic local systems. *Invent. Math.*, 207(1):291–343, 2017.
- [MT20] Matthew Morrow and Takeshi Tsuji. Generalised representations as q -connections in integral p -adic Hodge theory. *arXiv:2010.04059v2*, 2020.
- [MW22] Yu Min and Yupeng Wang. p -adic Simpson correspondence via prismatic crystals. *arXiv preprint arXiv:2201.08030v4*. *Appear to J. Eur. Math. Soc.*, 2022.
- [MW23] Yu Min and Yupeng Wang. Integral p -adic non-abelian hodge theory for small representations. *arXiv preprint arXiv:2304.07078v2*, 2023.
- [RC22] Juan Esteban Rodríguez Camargo. Geometric sen theory over rigid analytic spaces. *arXiv preprint arxiv:2205.02016v3*, 2022.
- [Sch13] Peter Scholze. p -adic Hodge theory for rigid-analytic varieties. *Forum Math. Pi*, 1:e1, 77, 2013.
- [Sch17] Peter Scholze. Étale cohomology of diamonds. *arXiv preprint arXiv:1709.07343*, 2017.
- [Wan23] Yupeng Wang. A p -adic Simpson correspondence for rigid analytic varieties. *Algebra & Number Theory*, 17(8):1453–1499, 2023.
- [Yu24] Jiahong Yu. On smooth adic spaces over \mathbf{B}_{dR}^+ and sheafified p -adic riemann–hilbert correspondence. *arXiv preprint arXiv:2403.01363*, 2024.

Y.L., SCHOOL OF MATHEMATICAL SCIENCE, PEKING UNIVERSITY, YIHEYUAN ROAD 5, BEIJING, 100190, CHINA.

Email address: 2100010764@stu.pku.edu.cn

X.N., ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, UNIVERSITY OF CHINESE ACADEMY OF SCIENCE, ZHONGGUANCUN EAST ROAD 80, BEIJING, 100190, CHINA

Email address: niexiecheng23@mails.ucas.ac.cn

X.Q., SCHOOL OF MATHEMATICAL SCIENCE, PEKING UNIVERSITY, YIHEYUAN ROAD 5, BEIJING, 100190, CHINA.

Email address: 2200010912@stu.pku.edu.cn

Y.W., BEIJING INTERNATIONAL CENTER FOR MATHEMATICAL RESEARCH, PEKING UNIVERSITY, YIHEYUAN ROAD 5, BEIJING, 100190, CHINA.

Email address: 2306393435@pku.edu.cn