

# Chapter 0 Exercises

Hui

## 1 Exercise 0.1

- a) The set of all odd natural numbers.
- b) The set of all integers that are multiples of 2.
- c) The set of all natural numbers divisible by 2.
- d) The set of all natural numbers divisible by 2 and 3.
- e) The set of all strings of 0s and 1s that are palindromic.
- f) An empty set.

## 2 Exercise 0.2

- a)  $\{n | n = 10^m \text{ for some } m \in \mathbb{N}\}$
- b)  $\{n \in \mathbb{Z} | n > 5\}$
- c)  $\{n \in \mathbb{N} | n < 5\}$
- d)  $\{\text{aba}\}$
- e)  $\{\epsilon\}$
- f)  $\emptyset$

## 3 Exercise 0.3

- a)  $A$  is not a subset of  $B$ .
- b)  $B$  is a subset of  $A$ .
- c)  $A \cup B = \{x, y, z\}$
- d)  $A \cap B = \{x, y\}$
- e)  $A \times B = \{(x, x), (x, y), (y, x), (y, y), (z, x), (z, y)\}$
- f)  $\mathcal{P}(B) = \{\emptyset, \{x\}, \{y\}, B\}$

## 4 Exercise 0.4

The Cartesian product of  $n$  sets is the set of all  $n$ -tuples where the first element is an element of the first operand, the second element is an element of the second operand, and so on. In the case of  $A \times B$ , the first element of each 2-tuple is an element of  $A$ , and the second element of each 2-tuple is an element of  $B$ . So, if  $A$  has  $a$  elements and  $B$  has  $b$  elements, to make every possible pair, you would

need  $a$  elements  $b$  times, or  $a * b$  pairs. Similarly, if  $A$  has  $a$  elements,  $B$  has  $b$  elements, and  $C$  has  $c$  elements,  $A \times B \times C$  has  $a * b * c$  elements.

## 5 Exercise 0.5

Each element of  $\mathcal{P}(C)$  is a subset of  $C$ , meaning each element of  $\mathcal{P}(C)$  is a permutation where each element of  $C$  is either present or not present (2 possibilities per element). Thus, if  $C$  is a set with  $c$  elements, there are  $2^c$  elements in  $\mathcal{P}(C)$ .

## 6 Exercise 0.6

- a)  $f(2) = 7$
- b)  $f : [1, 5] \rightarrow \{6, 7\}$
- c)  $g(2, 10) = 6$
- d)  $g : [1, 5] \times [6, 10] \rightarrow [6, 10]$
- e)  $g(4, f(4)) = g(4, 7) = 8$

## 7 Exercise 0.7

Let  $A = \{1, 2, 3\}$ .

- a)  $R_a = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 1), (2, 3), (3, 2)\}$   
 $R_a$  is reflexive because  $\forall x \in A(xR_ax)$ ,  $R_a$  is symmetric because  $\forall x \in A(xR_ay \rightarrow yR_ax)$ , and  $R_a$  is not transitive because  $\exists x \in A((xR_ay \wedge yR_az) \wedge \neg(xR_az))$
- b)  $R_b = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 3), (1, 3)\}$   
 $R_b$  is reflexive because  $\forall x \in A(xR_bx)$ ,  $R_b$  is transitive because  $\forall x \in A((xR_by \wedge yR_bz) \rightarrow xR_bz)$ , and  $R_b$  is not symmetric because  $\exists x \in A(xR_by \wedge \neg(yR_bx))$
- c)  $R_c = \{(1, 2), (2, 1), (1, 3), (3, 1), (2, 3), (3, 2)\}$   
 $R_c$  is symmetric because  $\forall x \in A(xR_cy \rightarrow yR_cx)$ ,  $R_c$  is transitive because  $\forall x \in A((xR_cy \wedge yR_cz) \rightarrow xR_cz)$ , and  $R_c$  is not reflexive because  $\exists x \in A \neg(xR_cx)$

## 8 Exercise 0.8

Hey Alexa how to use tikz-automata

## 9 Exercise 0.9

$(\{1, 2, 3, 4, 5, 6\}, \{(1, 4), (1, 5), (1, 6), (2, 4), (2, 5), (2, 6), (3, 4), (3, 5), (3, 6)\})$