


Intramammary tube for mastitis treatment in lactating animals

Antibiotic combination of Ampicillin and Cloxacillin indicated for the treatment of clinical mastitis in milking cows.



Udder of a cow infected with mastitis causing organisms

Solution;











Osho Chemical Industries Limited



Intramammary tube for mastitis treatment in lactating animals



COMPOSITION:

Ampicillin 75 mg
Cloxacillin 200mg
Base q.s

INDICATIONS:

TILOX is indicated for the treatment of mastitis in domestic animals. It is particularly recommended where the causative organisms have not been identified and where penicillin resistant *staphylococci* and E. Coli are known or are suspected to be involved.



DOSAGE AND ADMINISTRATION:

One Tilox syringe per quarter to be repeated every twelve hours for 3 days or as directed by the veterinarian.



After milking, clean and disinfect the teat, insert the nozzle of the tube into the teat and apply gentle continuous pressure on the plunger until all the suspension is expressed. The treated guarters may be milked out at the next normal milking time.



WITHDRAWAL PERIOD:

Milk from the treated animal must not be consumed until 60 hours after the last treatment.

PRESENTATION:

5g disposable syringes.





Cow's udder





Healthy herd





Industrial Area Office: Osho Complex, Sasio Rd. off Lunga lunga, Industrial Area
Postal Address: P. O. Box 49916 - GPO 00100 Nbi-Kenya Pilot Line: (+254) 020 391 2000, (+254) 0711-045 000,
Customer Service: (+254) 020 391 2500, (+254) 0711-045 500 Fax: (+254) 020 3912600
E-mail: oshochem@oshochem.com Website: www.oshochem.com

