

MancoFlo 455 SC ni dawa ya maji yenye nguvu wa mchanganyiko wa kemikali ya mancozeb ya kudhibiti Botrytis (kwenye maua na shina) kwenye Waridi, Anthracnose kwenye Mishiri, Early & Late Blight kwenye Nyanya na Anthracnose na Ascochyta Blights kwenye Mbaazi. Ni dawa isiyo na vumbi na haihitaji kutangulia kuchanganywa. Ikitumiwa kwa kiasi kilichopendekezwa, MancoFlo 455 SC haina madhara kwa maua, lakini yafaa kujaribiwa kwa mimae chache.

MAPENDEKEZO YA MATUMIZI

Nvunvizia MancoFlo 455 SC kwa mpango wa kuzuja ugoniwa.

• Tumia kiwango cha maji cha Lita 1000 kwa hekta na nyunyizia kulowesha mimea.

| | Mmea | Ugonjwa | Kiwango |
|---|---------|---------------------------------|------------------|
| | Mishiri | Anthracnose | Turrango |
| | Nyanya | Early & Late Blight | Lita 2 / Hektari |
| , | Mbaazi | Anthracnose & Ascochyta Blights | |
| | Waridi | Botrytis (ya shina na maua) | Lita 2 / Hektari |

Muda Salama wa kuvuna: siku 14 kwa mboga.

Muda salama wa kuingia pahali palipo nyunyiziwa: Masaa 6 TAHADHARI

Usivute sigara, usinywe wala kula unapotumia dawa hii. Vaa mavazi ya kinga kama vile overall, mifuko ya mikono, masks na kadhalika. Jihadhari dawa isiguze ngozi wala kuingia machoni.

UHIFADHI NA UTUPAJI

Uweke dawa hii katika chombo chake asili kilichofunikwa kamili na ihifadhiwe mbali na vyakula na watoto. Isafishe mikoba tupu ya dawa mara tatu kisha itoboe na kuitupa au kuizika mahili palipoidhinishwa kulingana na sheria za utupaji, mbali na visima au mito. Vyombo hivi visivitumiwe kamwe kwa shudhuli nyincine zozote.

MADHARA KWA MAZINGIRA

MancoFlo 455 SC haidhuru nyuki, lakini ni sumu kwa samaki. Hivyo basi mabaki ya dawa au mikebe funu haifai kutuowa kwenve visima au mito.

MAELEZO YA HUDUMA YA KWANZA

Ikigusana na ngozi: yafaa ngozi isafishwe vizuri na maji na sabuni. Mvuke ukivutwa yafaa mtu apelekwe kwenye hewa safi. Ikimezwa: Mhusika afaa kupelekwa kwa daktari mara moja. DALILI ZA SUMU

Madawa aina ya Carbamates yaweza kusababisha mikaso ya misuli ya tumbo, kutapika, n.k. Mgonjwa afaa kupelekwa kwa daktari mara moja.

IWAPO DHARURA: Wasiliana na nambari hii isiyo malipo-0800720021/0800730030 (inanatikana masaa 24)

(inapatikana masaa 24)

Dawa hii sharti itumiwe kulingana na maagizo yaliyo kwenye kibandiko. Ni hatia chini ya sheria za madawa ya kuangamiza wadudu waharibifu kutumia au kuhifadhi dawa hii katika hali isiyo

THIBITISHO

Dhamana ya muuzaji ni maelezo yaliyo kwenye kibandiko, ambayo ni lazima yafuatwe kikamilifu. Mnunuzi lazima akubali hatari zozote zinazoweza kutokana na matumizi ya dawa hii kwa binadamu au mali na lazima aikubali dawa hii katika hali hiyo.





MANCOFLO® FUNGICIDE 455 SC

Fungicide for control of Botrytis on Roses and leafspot in vegetables.

(Kiua dudu cha kudhibiti Botrytis kwenye waridi na leafspot kwenye mboga)

Distributed by (Husambazwa na): DSHO CHEMICAL INDUSTRIES LTD

P. O. Box 49916 00100 Nbi, Kenya Tel: 3912000 Mobile: 0726 880480 / 0733 880480 0733 524968 / 0711 045 000 Email: oshochem@oshochem.com Web: www.oshochem.com

Manufacturer (Imetengenezwa na): SULPHUR MILLS LTD. 303/304, T.V. ESTATE S.K. AHIRE MARG, WORLI, MUMBAI 400 025 INDIA

MEMBER OF AGROCHEMICAL ASSOCIATION OF KENYA AAK REG. NO. 275





COMMERCIAL AND AGRICULTURAL CLASS
KUNDI LA BIASHARA NA KILIMO.
GUARANTE (PHAMNA):
455g/L Mancozeb
REGISTRATION NO.
(NAMBARI YA USAJILI):(CR) 0431

READ THE LABEL BEFORE USING (SOMA KIBANDIKO KABLA YA KUTUMIA) KEEP LOCKED & OUT OF REACH OF CHILDREN (WEKA MBALI NA WATOTO) SHAKE WELL BEFORE USE



Batch No. (Nambari ya fungu):

Mfg Date(Tarehe ya kutengenezwa):

Expiry Date (Tumia kabla ya):

Shelf Life (Maisha Rafuni):

2 years in original unopened container stored in a cool, dry & dark place (Miaka miwili toka tarehe ya kutengenezwa ikiwa kwenye vyombo vyake asili visivyofunguliwa kwenye sehemu tulivu na kavu.)



DESCRIPTION. MancoFlo 455 SC is a liquid suspension concentrate of mancozeb for control of Botrytis (both flowers and stem) in Roses, Anthracnose on French beans, Early & Late Blight in Tomatoes and Anthracnose & Ascochyta Blights in Peas. It has no need for pre-mixes hence saves time and labour. Easy to pour and measure, no dust inhalation and minimum exposure to hazard. It is less abrasive to expensive spray equipment.

N.B: Roses: Mancozeb when used at the recommended rates does not cause phytotoxicity. Always check for varietal tolerance on a few plants.

RECOMMENDATIONS FOR USE

- Spray MancoFlo 455 SC in a preventative programme.
- MancoFlo may leave some residues on some plants and it is advisable to check out on some few plants before application.
- Use spray volumes of 1000 litres per Hectare and spray to wet the plant.

| | Crop | Disease | Rate |
|---|--------------|---------------------------------|----------|
| | French beans | Anthracnose | |
| | Tomatoes | Early & Late Blight | 2L / 1Ha |
| | Peas | Anthracnose & Ascochyta Blights | |
| - | Roses | Botrytis (Stem & Flower) | 2L / 1Ha |

Preharvest Interval (PHI): 14 days for vegetables

Re-entry period: 6 hours

HANDLING PRECAUTIONS: Do not smoke, eat or drink when handling this product. Wear protective clothing; overalls, face shields, gloves and boots when handling or spraying this product. Avoid skin or eye contact with product or spray mixture. Wash body and clothes after application.

STORAGE AND DISPOSAL: Store in original tightly closed container, away from children and livestock. Store in a cool, dry and dark place. Triple rinse empty container and add washings to spray tank. Puncture and bury or burn the container in accordance with National Regulation and Legislation, away from water sources and human habitation. Do not reuse containers for any other purpose.

ENVIRONMENTAL HAZARDS: MancoFlo 455 SC is hazardous to fish. Water sources or ponds should not be contaminated with concentrate or its solution or its empty containers.

FIRST AID INSTRUCTIONS: Skin or eye contact: Wash with plenty of water for at least 15 minutes. If ingested: Induce vomiting. Give patient copious quantities of water and obtain medical attention for gastric lavage. Never give anything by mouth to an unconscious person. Seek medical attention immediately.

TOXICOLOGICAL INFORMATION: Symptoms of poisoning: Carbamates can cause abdominal cramps, vomiting, nausea and mucous irritation. Antidote: No known antidote

Note to Physician: Treatment is essentially symptomatic.

In case of Emergency Call Toll free No. 0800720021 / 0800730030 (24 hrs)

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

SELLER'S GUARANTEE (WARRANTY): Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.