

TOTAL PROTECTION AGAINST INSECT PESTS



BETACYDE[®] 25 EC

Beta-cyfluthrin 25g/L

Broad spectrum, contact & stomach action insecticide with Rapid knockdown

for the control of wide range of insect pests like Thrips, Aphids, Whiteflies, Caterpillars & stink bugs etc.

Available packs
50ml, 100ml, 250ml, 1 litre




Pilot lines: 0711 045000 | 0732 167000
20560 (SMS) @
oshochem@oshochem.com
www.oshochem.com

osho[®]
CHEMICAL INDUSTRIES LTD.

BETACYDE[®] 25 EC
Beta-cyfluthrin 25g/L

DESCRIPTION: BETACYDE 25EC is a broad spectrum insecticide containing beta-cyfluthrin. It has rapid knock-down effect.

MODE OF ACTION: Non-systemic insecticide with contact, stomach action and long residual activity. Acts on the nervous system, with rapid knockdown and long residual activity. Uses Insecticide effective against Lepidoptera, Coleoptera, Hemiptera and Homoptera on cotton, fruit, vegetables, cereals and other crops, at 7.5-20 g/ha. Also against migratory locusts and grasshoppers



CROP	PEST	RATES/20L	RATES/Ha	RATES/Ha
Roses	Thrips, aphids, whiteflies, caterpillars	10ml	500ml	500
Wheat, barley, oats, rye and triticale	Grain aphid, rose grain aphid, wheat blossom midge	10ml	500ml	Apply when pests are seen or in response to specialist advice. Do not apply after flowering
Oilseed rape	Cabbage seed weevil, Cabbage stem weevil	10ml	500ml	Apply during the flowering period when pests have reached threshold levels. Repeat at a 10-14 day interval. Apply as soon as pest is seen or as advised by specialist advice.
	Brassica pod midge	10ml	500ml	Apply as soon as pest is seen or as advised by specialist advice.
Sugar beet	<i>Aphis fabae</i>	10ml	500ml	Apply as soon as pest is seen or as advised by specialist advice.
Cauliflower and cabbage	Cabbage moth, small cabbage moth and peach potato aphid	10ml	500ml	Apply as soon as the pest or damage is first seen. Allow 10-14 days between applications.

DIRECTIONS FOR USE: Pour the required quantity of BETACYDE 25EC into the spray tank, which is half filled with water and stir. Fill the tank with water to the required level as you agitate/mix. Spray at first sign of infestation and repeat every 7-14 days depending on pest infestation. Use 400 - 600 litres of water per hectare.

Available packs: 8g, 40g, 1kg