

hrs)

**Fungicide for the** control of downy in Emergency call Toll Free No. (24 ti wa Dharura piga Simu bila malipo (masaa 24) mildew and Botrytis in Roses. (Kiua kuvu cha kuzuia na kudhibiti downy mildew na Botrytis kwenye waridi).

## **OSHO CHEMICAL** INDUSTRIES LTD.

0726 880 480 | 0733 880 480 SMS: 20560 | Fax: [+254] 0711 045600

**INDOFIL INDUSTRIES LTD.** 

4th Floor, Kalpataru Sqaure, Kondivita Road, Off Andheri-Kurla Road, Andheri (E), Mumbai- 400 059, India.

MEMBER OF **AGROCHEMICAL** ASSOCIATION AAK REG. NO. 275






## **COMMERCIAL AND AGRICULTURAL CLASS** (KUNDI LA BIASHARA NA KILIMO)

GUARANTEE (DHAMANA): Carbendazim 12% w/w Mancozeb 63% w/w


REGISTRATION NO. (NAMBARI YA USAJILI): PCPB(CR)

**READ THE LABEL BEFORE USING** (SOMA KIBANDIKO KABLA YA KUTUMIA) KEEP LOCKED AND OUT OF REACH OF CHILDREN (FUNGIA MBALI NA WATOTO)









Expiry Date (Tumia kabla ya):

Shelf Life: 2 years in original unopened container stored in a cool, dry & dark place (Maisha Rafuni: Miaka miwili toka tarehe ya kutengenezwa ikiwa kwenye chombo chake asili kisichofunguliwa kwenye sehemu tulivu na kavu.)



ΥK

case of an (*Wakati* w

2





















**DESCRIPTION:** COMPANION®75WP is a curative and preventive fungicide with broad spectrum, systemic and contact activity in the control of Downy mildew and Botrytis on Roses. It has long disease control duration.

**DIRECTIONS FOR USE:** It is recommended that you avoid weather conditions that favour drift. Use a water volume of 1000L per her Hectare.

## **RATES OF APPLICATION**

CROP	DISEASE	RATE
Roses	Downy mildew, Botrytis	1.5kg/ha (30g/20L)

PREPARATION OF SPRAY MIXTURE: Fill the tank to half its capacity. In separate container add a small amount of water to the required quantity of COMPANION®75WP and mix well until a uniform mixture is obtained. Agitate and filter mixture into the tank, while adding water to the required amount. Ensure good mixing before starting to spray. Do not store diluted solution for later use. Ensure uniformity while spraying. Swirl-type nozzles are preferred. **REI (Re- entry Interval):** 12hrs.

**HAZARDOUS / SAFETY PRECAUTIONS:** Wear safety goggles or face shield. Eye contact will cause very slight irritation. Mist or dust concentration may be irritating to respiratory tract and eyes. Protective clothing (overall) and water proof gloves. Repeated or prolonged Skin contact may cause irritation. Wash hands before eating, drinking, chewing gum, using tobacco or using toilet. Ingestion of this product may cause liver abnormalities on prolonged or excess exposure. Remove clothing immediately if pesticide gets inside &wash

thoroughly. Wash the outside of gloves before removing .As soon as possible,

wash thoroughly and change into clean clothing.

STORAGE AND DISPOSAL: Keep out of reach of children under lock and key. Keep in original container. Store in a cool, dry place away from heat or open flame. Protect from sunlight. Keep away from feed , foodstuff and drinking water. Do not store above 37°C. Triple rinse empty pouch and add washings to spray mixture, then dispose empty pouch in a safe and authorised place away from human habitation and water sources, in accordance with national

WILDLIFE AND ENVIRONMENTAL PROTECTION: Do not mix, load or clean equipment within 30 metres of wellheads or aquatic systems. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes. Is safe to bees when recommended rates are used. TOXIC to wild mammals. Toxic to birds. Do not apply this product to blooming trees. This product is highly toxic to aquatic

regulations and legislation. Do not reuse pouch for any other purpose

FIRST AID INSTRUCTIONS: Remove the affected person from the danger zone to a well-ventilated room or to fresh air, and protect from under cooling. IN CASE OF SUSPECTED POISONING: Immediately call a physician. If in eye: Wash eyes with fresh water for 15 minutes. If irritation continues, see a doctor. If on skin: Remove contaminated clothing and wash skin with plenty of soap and water. Call doctor if irritation persists. If inhaled: Remove person from exposure area to fresh air, Keep patient at most comfort level and keep warm. Get medical attention. If swallowed: Do not induce vomiting. Give water to drink. Do not give anything by mouth if patient is unconscious. Seek medical

TOXICOLOGICAL INFORMATION: Symptoms of poisoning: sweating, headache, allergy, coughing, sneezing, itching, nausea maybe observed. Antidote: There is no specific antidote. SEEK MEDICAL ATTENTION.

Antidote: No specific antidote.

Note to Physician: Symptomatic treatment is recommended.

In case of Emergency Call Toll free No. 0800720021 / 0800730030 (24 hrs) **NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions. WARRANTY: Seller's guarantee is limited to the terms set out on the label and

subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

MAELEZO: COMPANION®75WP ni dawa yenye uwezo mkubwa, inayoponya na kuzuia magonjwa mengi ya kuvu. Hupenya ndani ya mmea na hutenda kazi inapoguza vijimea vya kuvu. Huzuia magonjwa *Downy mildew* na *Botrytis* kwenye Waridi. Inanguvu za kuzuia magonjwa kwa muda mrefu.

JINSI YA KUTUMIA: Nyunyizia dawa pasipo na upepo mkali. Tumia Lita 1000 za maji kwa Hektari . VIDIMO VVA MATUMIZI

VIFIMO VIA MATOMIZI				
MMEA	UGONJWA	KIWANGO		
Waridi	Downy mildew, Botrytis	kilo 1.5/hekta (gramu 30/Lita 20)		

JINSI YA KUTAYARISHA DAWA: Weka maji hadi nusu ya bomba la kunyunyizia. Katika mtungi mwingine ongeza maji kidogo kwa kiwango cha COMPANION®75WP kinachostahili halafu koroga sawa sawa. Tikisa nakuchunga mchanganyiko kwenye bomba, ukiongeza maji hadi kiwango kinachihitajika. Hakikisha umepata mchanganyiko maalum kabla yakuanza kunyunyizia. Usitumie mchanganyiko wa dawa uliopita siku moja.

Muda salama wa kuingia mahali palipo nyunyiziwa dawa: Masaa 12

HADHARI: Valia miwani ya kazi (goggles) au kizuizi cha uso. Ikiingia machoni huleta mwasho kidogo. Mvuke wa dawa hii pia huleta mwasho katika sehemu za kupumua na machoni, Nguo aina ya overall na vizuizi vya mikono (gloves). Ikiguza ngozi kwa muda mrefu huleta mwasho. Nawa mikono kabla ya kula, kunywa kumumunya peremende, kuvuta sigara au kuenda choo. Ikimezwa kwa kiasi kikubwa, huenda ikadhuru ini. Vua nguo dawa inapopenya halafu oga mara moja. Osha vizuizi vya mikono kabla kuvivua. Punde, nawa auoga nakubadilisha mayazi.

UHIFADHI NA UTUPAJI: Fungia mbali na watoto. Hifadhi katika kiwekeo chake halisi. Weka katika sehemu iliyokauka, isiyo najoto, yenye hewa mbali na mwangaza wajua. Mbali nalishe ya mifugo, vyakula, mbole, dawa za kilimo na maji ya kunywa. Funga kiwekeo kabisa wakati hautumii dawa. Usihifadhi katika stoo yenye kiwango cha joto juu ya 37°C. Tupa pochi tupu mahali palipo salama na paliporuhusiwa, mbali na makao ya watu na vianzo vya maji, kulingana na sheria za utupaji.Usitumie pochi tupu kwa matumizi mengine

MAZINGIRA: Usichanganye, usipakie au kusafisha dawa mita 30 kutoka kwa visima au sehemu zenye maji ardhini. Usisumishe maji yanayotumiwa kunyunyizia mimea, yanayotumiwa kunywa au yanayofugwa samaki unapoosha vyombo vya dawa au masalio ya dawa. Ni salama kwa nyuki ukizingatia viwango vilivvopendekezwa. Ni sumu kwa wanyama waporina ndege waangani. Usinyunyizie kwa mimea inayootesha maua. Ni sumu dhidi va wanyama wamajini.

**HUDUMA YA KWANZA:** Toa mhadhiriwa kwenye eneo lahatari na kumweka mahali penye hewa safi wala sio kwenye baridi kali. Iwapo amesumika, mwarifu daktari mara moja. Ikiingia machoni: Osha macho kwa maji mengi kwa muda wa dakika15. Mwasho ukizidi, muone daktarii. **Ikiguzangozi:** Osha kwa sabuni na maji mengi. Mwasho ukizidi, muone daktari. Ikipumuliwa: Muondoe mwasiriwa kwenye hatari hadi mahali penye hewa safi. Ikiwa anashida ya kupumua, mpe pumzi kupitia kinywa chake. Pata huduma ya daktari haraka iwezekanavyo. **Ikimezwa:** Usimsaidie mwasiriwa kutapika. Kunywa glasi. Usimpe kitu chochote kupitia kinywa ikiwa amepoteza fahamu. Peleka mjeruhi kwa daktari.

MAELEZO YA SUMU: Dalili za kusumishwa: kutokwa na jasno, kuumwa na kichwa, allergy, kukohoa, kupiga chafya, mwashowakujikuna,nakuhisi kutapika. PELEKA MJERUHI KWA DAKTARI.

Tiba ya sumu: Haina tiba maalum

Arifa kwa daktari: Tibu kulingana na dalili za sumu

Wakati wa dharura piga simu bila malipo-0800720021/0800730030

ILANI KWA MTUMIAJI: ILANI KWA MTUMIAJI: Dawa hii sharti itumiwe kulingana na maagizo yaliyo kwenye kibandiko. Ni hatia chini ya sheria za madawa ya kuangamiza wadudu waharibifu kutumia au kuhifadhi dawa hii

THIBITISHO: Dhamana ya muuzaji ni maelezo yaliyo kwenye kibandiko, ambayo ni lazima yafuatwe kikamilifu. Mnunuzi lazima akubali hatari zozote zinazoweza kutokana na matumizi ya dawa hii kwa binadamu au mali na lazima aikubali dawa hii katika hali hiyo.