

1. 解: (Ch 2.2.16)

$$P_X(x) = \begin{cases} x^2/a, & x = -3, -2, -1, 0, 1, 2, 3 \\ 0 & \text{otherwise} \end{cases}$$

$$(a) \sum P_X(x) = 1 \Rightarrow \frac{1}{a}(9+4+1) \times 2 = 1 \Rightarrow a = 28$$

$$E(X) = \sum x P_X(x) = 0 \quad (\text{由对称性})$$

$$(b) Z = (X - E(X))^2. \text{ PMF.}$$

$$\text{由 } EX=0, \quad Z=X^2. \quad x=\sqrt{z} \Rightarrow P_Z(z) = P_X(\sqrt{z}) + P_X(-\sqrt{z})$$

$$\text{当 } (z=1, 4, 9) \Rightarrow P_Z(z) = \begin{cases} z/14, & z=1, 4, 9 \\ 0 & \text{otherwise} \end{cases}$$

$$(c) \text{var}(X) = E(X - EX)^2 = EZ = \sum P_Z(z) \cdot z = \frac{1}{14} + \frac{16}{14} + \frac{81}{14} = 7$$

$$(d) \text{var}(X) = \sum_x (x - EX)^2 P_X(x)$$

$$= \left(9 \times \frac{9}{28} + 4 \times \frac{4}{28} + 1 \times \frac{1}{28} \right) \times 2 = 7$$

2. 解: (Ch 2.2.2)

(a). 同时抛, 结果可为 $H_1H_2, H_1T_2, T_1H_2, T_1T_2$ 由独立性

记事件 $A = \{H_1T_2, T_1H_2\}$. 则在一次抛掷中,

$$P(A) = p(1-p) + p(1-p)$$

记事件 A_k 为在第 k 次抛掷中 A 发生. 记 X 为题中要求的次数.

$$\text{所求 } P_X(k) = P(\bar{A}_1 \bar{A}_2 \cdots \bar{A}_{k-1} A_k) = (1 - p(1-p) - p(1-p))^{k-1} (p(1-p) + p(1-p))$$
$$(k=1, 2, \dots), \quad X \sim G(p(1-p) + p(1-p)), \text{ 记 } m = \text{参数}.$$

因此有 (2 课结论), $EX = \frac{1}{p(1-p) + p(1-p)} = \frac{1}{p(1-p)}$

$$EX^2 = E(X(X-1)) + EX = \sum_{j=1}^{\infty} j(j-1)(1-m)^{j-1} \cdot m + \frac{1}{m}$$

$$= \sum_{j=0}^{\infty} (1-m) \frac{\partial^2}{\partial m^2} (1-m)^j \cdot m + \frac{1}{m} = (1-m) \cdot \frac{2}{m^3} \cdot m + \frac{1}{m}$$

$$\Rightarrow \text{Var} X = EX^2 - (EX)^2 = \frac{2(1-m)}{m^2} + \frac{1}{m} - \frac{1}{m^2} = \frac{2-2m+m-1}{m^2} = \frac{1-m}{m^2}$$

$$\Rightarrow \text{本题 } \text{Var}(X) = \frac{pq + (1-p)(1-q)}{[p(1-q) + q(1-p)]^2}$$

$$(b) \text{ 所求概率 } P(H_1 T_2 | \{H_1 T_2, H_2 T_1\}) = \frac{P(H_1 T_2)}{P(H_1 T_2 \cup H_2 T_1)} = \frac{P(H_1 T_2)}{P(H_1 T_2) + P(H_2 T_1)}$$

$$= \frac{p(1-q)}{p(1-q) + q(1-p)}$$

3. 解 (ch 3.13)

$$X \sim N(\mu, \sigma^2) = N(10, 10^2) \quad \text{令 } Z = \frac{X - \mu}{\sigma} \quad (\text{Standardization})$$

X : in Celsius. Y : in Fahrenheit.

$$\text{relation between } X \text{ and } Y: \quad X = \frac{5}{9}(Y - 32)$$

$$\Rightarrow P(Y \leq 59) = P(X \leq 15) = P(X \leq \frac{1}{2}) = \Phi(0.5) = 0.692$$

$$\therefore P(Y \leq 59) = 0.692$$

4. 解:

① $X \sim \chi_n^2$. proof of $EX = n$, $\text{Var}(X) = 2n$.

由 χ_n^2 分布的定义 $\Rightarrow X = Z_1^2 + Z_2^2 + \dots + Z_n^2$. Z_1, \dots, Z_n 为 i.i.d., $\sim N(0, 1)$

$$\Rightarrow \text{则 } EX = E(Z_1^2 + Z_2^2 + \dots + Z_n^2) = n E Z_1^2$$

$$Z_1 \sim N(0,1) \text{ 即有 } f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-z^2/2}, \int_{-\infty}^{\infty} z^2 f_Z(z) dz = 1$$

$$\Rightarrow EX = n \times 1 = n.$$

$$\text{Var}(X) = \text{Var}(Z_1^2 + Z_2^2 + \dots + Z_n^2) \text{ 由 } Z_1, \dots, Z_n \text{ 独立, } Z_1^2, \dots, Z_n^2 \text{ 也 independent}$$

$$\therefore \text{Var}(X) = \sum_{i=1}^n \text{Var}(Z_i^2) = n \cdot \text{Var}(Z_1^2)$$

$$\text{Var}(Z_1^2) = E(Z_1^4) - (EZ_1^2)^2 = \int_{-\infty}^{\infty} z^4 f_Z(z) dz - 1 = 2, \text{ Var}(X) = 2n$$

$$(\text{利用了 } \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} z^4 e^{-z^2/2} dz = 3. \text{ 由 Gaussian 积分得到})$$

$$\text{thus, } EX = n, \text{ Var}(X) = 2n.$$

$$\textcircled{2} T \sim t_n.$$

$$Z \sim N(0,1), X \sim \chi_n^2, X \perp Z. T = \frac{Z}{\sqrt{\frac{X}{n}}}, T \sim t_n.$$

$$\text{由 } X \perp Z \Rightarrow g(X) \perp Z \therefore \text{有如下的式子成立}$$

$$ET = E\left(\frac{Z}{\sqrt{\frac{X}{n}}}\right) = EZ \cdot E\left(\frac{1}{\sqrt{\frac{X}{n}}}\right)$$

$$\text{由 } EZ = 0. \text{ 且 } n \geq 2 \text{ 时, } E\left(\frac{1}{\sqrt{\frac{X}{n}}}\right) \text{ 存在. } \therefore ET = 0. (n \geq 2).$$

$$\text{Var}(T) = ET^2 - (ET)^2 \quad (EZ^2 = 1)$$

$$= E\left(\frac{Z^2}{\frac{X}{n}}\right) = n E\left(\frac{Z^2}{X}\right) = n EZ^2 E\left(\frac{1}{X}\right) = n E\left(\frac{1}{X}\right)$$

$$\text{只要证明 } E\left(\frac{1}{X}\right) = \frac{1}{n-2} \text{ 即可. 由 } X \sim \chi_n^2. \text{ 有 } f_X = \frac{(\frac{1}{2})^{\frac{n}{2}} \cdot x^{\frac{n}{2}-1}}{\Gamma(\frac{n}{2})} e^{-\frac{x}{2}}$$

$$\Rightarrow E\left(\frac{1}{X}\right) = \int_0^{\infty} x f(x) dx$$

$$= \int_0^{\infty} \frac{1}{x} \frac{(\frac{1}{2})^{\frac{n}{2}} \cdot x^{\frac{n}{2}-1}}{\Gamma(\frac{n}{2})} e^{-\frac{x}{2}} dx$$

$$= \frac{(\frac{1}{2})^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} \int_0^{\infty} x^{\frac{n}{2}-2} e^{-\frac{x}{2}} dx$$

$$(n \geq 2)$$

$$= \frac{(\frac{1}{2})^n}{P(\frac{n}{2})} \frac{P(\frac{n}{2}-1)}{(\frac{1}{2})^{\frac{n}{2}-1}} = \frac{1}{n-2} \quad \therefore \text{Var}(T) = \frac{n}{n-2} \text{ 证毕!}$$

(n>3)

$$\textcircled{3} F = \frac{X_1/m}{X_2/n} \sim F(m, n). \quad X_1 \sim \chi_m^2, \quad X_2 \sim \chi_n^2$$

$$EF = E\left(\frac{X_1/m}{X_2/n}\right) = \frac{n}{m} E\left(\frac{X_1}{X_2}\right) = \frac{n}{m} E(X_1) E\left(\frac{1}{X_2}\right)$$

利用 ① 问结论有 $E(X_1) = m$, ② 问结论有 $E\left(\frac{1}{X_2}\right) = \frac{1}{n-2}$

$$\therefore EF = \frac{n}{n-2} \text{ 证毕!} \quad (n>3)$$

补充: 求 F 分布的方差.

$$\text{Var } F = EF^2 - (EF)^2 = EF^2 - \frac{n^2}{(n-2)^2}.$$

$$\text{而由 } EF^2 = \frac{n^2}{m^2} EX_1^2 E\left(\frac{1}{X_2^2}\right)$$

$$EX_1^2: \quad X_1^2 \sim \chi_m^2,$$

$$EX_1^2 = \int_0^\infty x^2 \cdot \frac{(\frac{1}{2})^{m/2}}{P(\frac{m}{2})} x^{\frac{m}{2}-1} e^{-\frac{x}{2}} dx = \frac{(\frac{1}{2})^{m/2}}{P(\frac{m}{2})} \int_0^\infty x^{\frac{m}{2}+1} e^{-\frac{x}{2}} dx$$

$$= \frac{(\frac{1}{2})^{m/2}}{P(\frac{m}{2})} \times \frac{P(\frac{m}{2}+1)}{(\frac{1}{2})^{\frac{m}{2}+2}} = m(m+1)$$

$$E\frac{1}{X_2^2}: \quad X_2^2 \sim \chi_n^2$$

$$E\frac{1}{X_2^2} = \int_0^\infty \frac{1}{x^2} \cdot \frac{(\frac{1}{2})^{n/2}}{P(\frac{n}{2})} x^{\frac{n}{2}-1} e^{-\frac{x}{2}} dx = \frac{(\frac{1}{2})^{n/2}}{P(\frac{n}{2})} \times \frac{P(\frac{n}{2}-2)}{(\frac{1}{2})^{\frac{n}{2}-2}} = \frac{1}{(n-2)(n-4)}$$

$$\therefore EF^2 = EF^2 - (EF)^2 = \frac{2n^2(m+n-2)}{m(n-2)(n-4)}$$

5. 解:

Chebyshev's inequality: $P(|X - EX| \geq a) \leq \frac{\text{Var}(X)}{a^2}$

设 X 表示正面向上的次数. $X \sim B(N, p)$. 有 $\mu = EX = Np = \frac{1}{2}N$

$$\sigma^2 = EX^2 = Np(1-p) = \frac{1}{4}N$$


$$\therefore P(|X - \frac{1}{2}N| \geq \frac{1}{4}N) \leq \frac{\frac{1}{4}N}{(\frac{1}{4}N)^2} = \frac{4}{N}$$

我得到了这个事件发生概率的一个上界, $\frac{4}{N}$.

6. 解. (造数)

7. 解: (Ch 3.34)

$$(a). f_p(p) = \begin{cases} pe^p & p \in [0, 1] \\ 0 & \text{otherwise} \end{cases}$$



$$E(p) = \int_0^1 p f_p(p) dp = \int_0^1 p^2 e^p dp = \int_0^1 p^2 de^p = e - 2 \int_0^1 p e^p dp$$

$$\text{而 } \int_0^1 p e^p dp = 1 \therefore E(p) = e - 2$$

\therefore probability of $\{H_1\} = e - 2$

$$(b). \text{求 } f(p|H) = \frac{f(p, H)}{P(H)} = \frac{P(H|p) \cdot f_p(p)}{E(p)} = \frac{p \cdot f_p(p)}{e - 2}$$

$$\therefore f_{p|H} = \begin{cases} \frac{p^2 e^p}{e - 2} & p \in [0, 1] \\ 0 & \text{otherwise} \end{cases}$$

(c) 求 $P(H_2|H_1)$

$$P(H_2|H_1) = \frac{P(H_1 \cap H_2)}{P(H_1)} = \frac{E_p(P(H_1 \cap H_2))}{E(p)} = \frac{\int_0^1 p^2 f_p(p) dp}{e-2}$$

$$\int_0^1 p^2 f_p(p) dp = \int_0^1 p^3 e^p dp = 6-2e$$

$$\therefore P(H_2|H_1) = \frac{6-2e}{e-2} \approx 0.784 > P(H_1)$$

大于说明我们在得知 H_1 的信息后, 更新了对 p 的认知.

8. 解: $(a) \xrightarrow{\frac{1}{2}} (a) \xrightarrow{\frac{1}{2}} (a+1)$ 状态转移

停止时一共只输了 X_0 局, 本金为 a , 求 EX_a .

由全期望定理:

$$EX_a = \frac{1}{2} \times (EX_{a+1} + 1) + \frac{1}{2} (EX_{a-1} + 1)$$

$$\therefore \text{得到递推关系 } EX_{a+1} - 2EX_a + EX_{a-1} + 2 = 0 \quad (*)$$

$$\text{边界条件: } EX_0 = 0, \quad EX_{a+b} = 0,$$

$$\text{由 } (*) \text{ 式有 } \underbrace{EX_{a+1}}_{S_{a+1}''} - EX_a = \underbrace{EX_a - EX_{a-1}}_{S_a''} - 2$$

$$\Rightarrow S_{a+1}'' - S_a'' = -2 \Rightarrow S_a'' = -2 \cdot a + \underset{\substack{\uparrow \\ \text{常数项}}}{m}$$

$$\therefore EX_a - EX_{a-1} = -2a + n \quad ①$$

$$\left\{ \begin{array}{l} EX_{a-1} - EX_{a-2} = -2(a-1) + n \quad ② \\ \vdots \\ EX_1 - EX_0 = -2 \cdot 1 + n \quad ④ \end{array} \right.$$

$$\Sigma: EX_a = -2 \cdot \frac{a+1}{2}a + an \Rightarrow$$

$$\text{由 } EX_{a+b} = -2 \cdot \frac{a+b+1}{2}(a+b) + n(a+b) = 0$$

$$\therefore n = a+b+1$$

$$\therefore EX_a = -(a+1)a + a(a+b+1) = ab !$$