

MRN:	00000690	Department:	Endocrinology and Diabetes
Patient Name :	Shahid Ehsan	Date/Time of visit:	10/2/2026 3:11:00 PM
Patient Contact No:	0300-0068443	Gender	[Male / 44 Y]
Date Of Birth:	01-Jan-1982		

Diagnosis

Active Problems :

[Primary] - [Acute] R7303 - Prediabetes

[Primary] - [Acute] E669 - Obesity, unspecified

CURRENT MEDICATION :

Medicine :

1. Tirzee 2.5 mg
2. Dolmet Tab
3. Vonoprazan 20mg Cap
4. TIRZEE 10MG/0.5ML PRE-FILLED PEN 1'S
5. Dolmet Tab
 - Zyloric 300mg

Visit Summary

- Follow Up
- Taking Tirzee 10 mg

Vitals :

Weight: 84.0 kgs

Height: 1.74 m

BMI: 27.74

Temp:

BP: 110 / 91 mmHg

SpO2:

RR:

Pulse: 96 BPM

BG: 127 mg/dl

INVESTIGATION RESULTS

Lab Results :

- HbA1c 5.5%

17/08/26

- TSH 3.53 (T4 1.09)
- Serum Uric acid 5.5

Printed By
Ali Raza Naqvi

*This is an system generated report with no signature required

Generated By
Ali Raza Naqvi

consultant
Diabetologist &
Endocrinologist

10/2/2026 3:41:37 AM

Prime Health Hub Dew Plaza No. 154, CCA 1, Sector C DHA Phase 6, Lahore

10/2/2026 3:41:37 AM

MRN:	00000690	Department:	Endocrinology and Diabetes
Patient Name :	Shahid Ehsan	Date/Time of visit:	10/2/2026 3:11:00 PM
Patient Contact No:	0300-0068443	Gender	[Male / 44 Y]
Date Of Birth:	01-Jan-1982		

- HbA1c 5.8%
- CRT 1.20

PLAN

Comments :

- Follow a high-protein, low-carb diet (under 30 g carbs/day).
- Complete at least 150 minutes of cardio weekly and strength train 2–3 times per week.
- Drink at least 3 L of water daily; avoid juices and sugary drinks.
- We discussed GLP-1 receptor agonists (semaglutide vs. tirzepatide), including their mechanism of action, expected benefits, and potential side effects.
- Referred to Rheumatologist (Dr. Shakaib Oureshi) for his heel pain**
- Continue with Tirzee 10 mg for now; increase the dose to 12.5 mg after the rheumatologist appointment.

Plan : wean off dose of tirzepatide on next consultation

Treatment Medications :

1. TIRZEE 12.5 MG/0.5ML PRE-FILLED PEN 1'S - -

Directions :

1 Injection Once a week for 4 Weeks

بدایات :

ایک انجکشن () ہفتے میں ایک بار چار ہفتے کے لئے

2. ZANOV 20 mg - -

Directions :

1 Capsule(s) Before Breakfast

بدایات :

ایک کیپسول () ناشتے سے پہلے

3. METHIX TAB 20'S - -

Directions :

1 Tablet(s) After Breakfast

بدایات :

ایک گولی () ناشتے کے بعد

Follow Up : 1. Follow-Up Visit after 1 Month (01-Nov-2026)

Printed By
Ali Raza Naqvi

*This is an system generated report with no signature required

Generated By
Ali Raza Naqvi

consultant
Diabetologist &
Endocrinologist

10/2/2026 3:41:37 AM

Prime Health Hub Dew Plaza No. 154, CCA 1, Sector C DHA Phase 6, Lahore

10/2/2026 3:41:37 AM