Print Client Name:		Client Identification Number: Date:		Adolescent 65D-30 Intensive		Outpatient		
						ASAM	Level	II.1
DIMENSIONS	[1] ADMISSION (		Place a check	in the appropri	Pension that apply to the ate box that indicates accement into this level	validation	YES	ИО
ASAM Requirements	may be transferred intensive level and dimension. A clien to address the clie	to this level of ca require this inten t may transfer from nt's needs or when	re when they h sity of servic ASAM Level I ASAM Level I h	ave met essenti e provided at t when services a as consisted of	, or 6. Transfer crite: al treatment objectives his level of care in at t that level have been motivational intervent are for which admission	in a more least one insufficient ions to		
Dimension 1: Acute Intoxication and/or Withdrawal Potential	$\square$ c. The client has	ee from intoxication intoxication or wit	n or withdrawal thdrawal sympto t to sustain to	l symptoms/risks oms/risks can be reatment and to	s; or e managed at this level follow treatment recomm			
Dimension 2: Biomedical Conditions and Complications	dimension is charac  a. The client's will not inte  b. The client's sufficient to medical manage	terized by one of t biomedical condition erfere with treatment biomedical condition of distract from reco	he following: ons, if any, and the this level ons are not seven efforts. e provided by the	re stable or are el of care; or vere enough to volume Such conditions the intensive or	el II. The client's state being concurrently add warrant inpatient treatments require medical monitoutpatient program or threatments.	dressed and ment but are pring and/or		
Dimension 3: Emotional, Behavioral or Cognitive Conditions and Complications	property and safety between daily superviols.  b. Interference affected by and requires  c. Social Functioning, daily living  d. Ability for some manage the addinterventions illness. Poor transitioning e. Course of II.	thality: The client requires frequent ren IOP sessions. He ision; with Addiction Recommendation and emotional, behave increased intensity ioning: The clients but not to such a cort of fulfill response of the client structures of daily is a problems may involve self-care or lack to adulthood, or increase: The client's	is at mild rismonitoring to a power, his or powery Efforts: ioral or cognity to support to symptoms are called the possibilities at it is experient living, and the policy poor hygical of independent in a younger achistory and proposed to the policy of the policy of independent in a younger achistory and proposed to the policy of the pol	sk of behaviors assure that then her condition if the client's retive problem, who reatment participates and it is to mooth the client of the	endangering self, other re is a reasonable likeled is not so severe as to recovery efforts are negation causes mild interfeit pation and/or compliance moderate difficulty in sele to manage the activit	cihood of require  citively rence with re; rocial ries of reatment ronic mental ris roports; roal,		

Revised 7/10/06 1 F-PR-1035

Dimension 4: Readiness to Change	□ a. □ b.	the stages of change, as evidenced a verbally compliant, but does not der passively involved in treatment; or attendance at outpatient sessions of interventions are not feasible or as The client's perspective inhibits his change. The client thus requires stinterventions are not feasible or as	apy and a programmatic milieu to promote proposed by behaviors such as the following: [1] the monstrate consistent behaviors; [2] the clie [3] the client demonstrates variable completed in self- or mutual-help meetings or support of the not likely to succeed in a level I service or her ability to make progress through the tructured therapy and a programmatic milieus or not likely to succeed in a Level II.1 settlements.	adolescent is ent is only iance with groups. Such ces; or the stages of . Such	
Dimension 5: Relapse/Continu ed Use Potential	□ a. □ b. □ c.	structured support, as indicated by difficulty in postponing immediate of or The client is assessed as being unable behaviors, which threaten abstinence Despite active participation in treaddiction symptoms (such as craving relapse.	ol or drug use without close outpatient mon- the client's lack of awareness of relapse of gratification, and/or ambivalence/resistance ole to interrupt his or her impulsive and see in the absence of ongoing clinical support atment, the client is experiencing an inten- and drug seeking behavior) with associated	triggers, e to treatment; elf-defeating t; or sification of	
Dimension 6: Recovery Environment	□ a. □ b.	Continued exposure to current job, and the client has insufficient or adequate level of functioning without. The client lacks social contacts, or recovery, or has few friends or peer has insufficient resources or skills without the services of a level II. of functioning between sessions; or The client family is supportive of a	r has inappropriate social contacts that jed rs who do not use alcohol or other drugs. I s necessary to maintain an adequate level of l program, but is capable of maintaining an	to maintain an opardize He or she also f functioning adequate level amily dysfunction	
Recommendations,		Counselor	Signature/Credential:	Date:	