

Search for Particles with Anomalous Charge in the IceCube Detector

Ward Van Driessche



Search for Particles with Anomalous Charge in the IceCube Detector

Ward Van Driessche

Author DATUM

Certified by prof. dr. D. Ryckbosch

Thesis Supervisor

Accepted by ???

Chair Examination Committee

Supervisor: prof. dr. D. Ryckbosch
Dissertation submitted in fulfillment of the requirements for the degree of
Doctor (Ph.D.) in Science: Physics

Department of Physics and Astronomy
Faculty of Sciences
Ghent University
Academic year ???



- To bla bla bla

Abstract

Bla bla bla Ik zou graag ook mezelf willen bedanken Of dat je in het middelbaar leert dat je eigenlijk een bundel bent van elektronen en protonen en bla bla bla, maar van zodra je fysica studeert besef je dat je eigenlijk een bundel trillingen in velden bent.




Table of Content

I	Theory and Experiment	
1	The Standard Model of Particle Physics	7
1.1	What we call matter: fermions	7
1.1.1	Leptons	8
1.1.2	Quarks	8
1.2	How particles communicate: interactions	8
1.3	The Standard Model in theory	11
1.3.1	SU(3): quantum chromodynamics	12
1.3.2	SU(2) \times U(1): electroweak	12
1.3.3	Brout-Englert-Higgs mechanism	13
1.3.4	Particle mixing	17
1.4	A success story	19
1.4.1	Wave-particle duality	19
1.4.2	Predictions	20
1.4.3	Precision tests	20
1.5	The need for physics beyond the Standard Model	21
1.5.1	Running of the coupling constants	24
1.5.2	Unifying theories	26
1.5.3	How to look for new physics	26
2	Theoretical Motivation of the Analysis	29
2.1	Introduction	29
2.2	Theory	29
2.3	Properties of the signal	30
2.4	Previous searches	30
2.4.1	Searches with accelerators and fixed targets	30

2.4.2	Colliders	31
2.4.3	Searches for particles with telescopes	32
3	Cosmic Rays and Neutrinos	35
3.1	Discovery of cosmic rays	35
3.2	What are cosmic rays?	35
3.2.1	Solar modulation	38
3.2.2	Galactic component	38
3.2.3	Extragalactic component	38
3.3	Acceleration mechanism	40
3.3.1	Supernova (remnants)	41
3.3.2	Active Galactic Nuclei	45
3.3.3	Blazars	45
3.3.4	Gamma Ray Bursts	45
3.4	Neutrinos	45
3.5	Air shower	45
3.5.1	Cosmological	45
3.5.2	Solar	45
3.5.3	Atmospheric	45
3.5.4	Astrophysical	45
3.6	Cosmic ray and neutrino detectors	45
4	Relativistic Particles in Matter	47
5	The IceCube Detector	49

II

Simulation, Data Processing and Analysis

6	Event Generation, Simulation and Reconstruction	53
6.1	Line-Fit	53
6.2	SPE and MPE	53
6.3	Millipede	53
6.4	Boosted Decision Tree Classifiers	53
7	The SPACE Analysis	55
7.1	Motivation	55
7.2	General strategy	55
7.3	Event cleaning	55
7.4	Variables	55
7.5	BDT results	55
7.6	Pull validation	55
7.7	Systematic Uncertainties	55
7.8	Results	55
8	Summary and Discussion	57
	Appendices	59
A	Gauge symmetries	61

B	Planck's law	63
B.1	Electromagnetic waves in a cubical cavity	63
B.1.1	Classical approach	64
B.1.2	Quantum approach	64
9	Some useful things for LaTeX	65
9.1	Definitions	65
9.2	Remarks	65
9.3	Corollaries	65
9.4	Propositions	65
9.4.1	Several equations	66
9.4.2	Single Line	66
9.5	Examples	66
9.5.1	Equation and Text	66
9.5.2	Paragraph of Text	66
9.6	Exercises	66
9.7	Problems	66
9.8	Vocabulary	66
	Bibliography	67
	Articles	67
	Books	70
	Index	71



Paragraphs of Text

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetuer id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetuer adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo


velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea dictumst. Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed, volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend faucibus, vehicula eu, lacus.

Theory and Experiment

1	The Standard Model of Particle Physics .	7
1.1	What we call matter: fermions	
1.2	How particles communicate: interactions	
1.3	The Standard Model in theory	
1.4	A success story	
1.5	The need for physics beyond the Standard Model	
2	Theoretical Motivation of the Analysis .	29
2.1	Introduction	
2.2	Theory	
2.3	Properties of the signal	
2.4	Previous searches	
3	Cosmic Rays and Neutrinos	35
3.1	Discovery of cosmic rays	
3.2	What are cosmic rays?	
3.3	Acceleration mechanism	
3.4	Neutrinos	
3.5	Air shower	
3.6	Cosmic ray and neutrino detectors	
4	Relativistic Particles in Matter	47
5	The IceCube Detector	49



The aim of this chapter is to give a summary of the theoretical framework that is used in particle physics. This framework was developed in stages throughout the latter half of the 20th century and is known as the Standard Model of particle physics. The Standard Model is a quantum field theory which is able to describe most of what is seen in particle physics experiments and proved to be successful in predicting later experimental discoveries. In this chapter a brief historical overview of the development of this theory will be given together with a limited description in order to familiarize the reader with concepts that will be used throughout this work. For a more in-depth and exhaustive discussion I refer to Refs. [14, 25, 60, 62]. [Sections bla bla bla](#)

1.1 What we call matter: fermions

Physics (from Ancient Greek: φυσική - *physikē*, “knowledge of nature”) is the natural science that studies matter. Matter is made up by *atoms* (from Greek: *atomos*, “indivisible”)* which can bind together into molecules which results into what is around us in our everyday life. Atoms are made up of a positively charged *nucleus* and surrounded by one or more *electrons* which are bound to the nucleus. The nucleus is made up of one or more *protons* and typically an approximately equal amount of *neutrons*. Because of their similar characteristics protons and neutrons are often referred to as *nucleons* and together they make up of more than 99.94% of an atom’s mass. Nucleons are made up of smaller particles called *quarks*[†] which are, as far as we know, *fundamental particles*. This means that we believe there are no smaller substructures making up these objects and are in essence mathematically best described as infinitely small. Because of this they are often referred to as pointlike. In the Standard Model (SM) these particles are characterized in our three-dimensional space as *fermions* which have odd half-integer spins obeying the laws of quantum mechanics. The spin of a particle is often illustrated with its classical counterpart in which an object is spinning and thus carries an intrinsic angular momentum. This analogy cannot be extrapolated to pointlike particles but the property happens to hold the same units as the classical orbital momentum. The spin of a particle seems to be

*Coined by ancient Greek philosophers Leucippus and his pupil Democritus who believed matter was made up of discrete units.

[†]The word “quark” originally appeared in the novel *Finnegans Wake* written by the Irish author James Joyce (1882–1941). The protagonist of the book dreams that he is serving beer to a drunken seagull. Instead of asking for “three quarts for Mister Mark” the inebriated bird says “three quarks for Muster Mark”. Murray Gell-Man had the habit of using names like “squeak” and “squark” for peculiar objects and after encountering the sentence in the book the name struck him as appropriate since the hypothetical came in threes.

just another property particles have, like charge or mass. Fermions follow Fermi-Dirac statistics and therefore obey the Pauli exclusion principle. As a consequence fermions cannot occupy the same place at the same time. (More formally, no two fermions may be described by the same quantum numbers.) This agrees with our macroscopic observations of matter in everyday life: people cannot walk through walls!

In total the SM distinguished 24 different fermions which can be subdivided into two distinct classes: *quarks* and *leptons*. There are six quarks (up, down, charm, strange, top and bottom), and six leptons (electron, electron neutrino, muon, muon neutrino, tau and tau neutrino), along with the corresponding antiparticle of each of these. A summary of the particles in the Standard Model is given in Figure 1.1.

1.1.1 Leptons

Leptons* can be subdivided into two classes: electromagnetically charged particles (e , μ and τ) and the neutral neutrinos (ν_e , ν_μ and ν_τ). Because of their charge, electrons are the well known particles combining into atoms together with nucleons. Being the lightest of the three, the electron is said to be part of the first *generation* together with the electron neutrino. Muons differ only from electrons in mass[†] and make up the second generation together with muon neutrinos. Similarly tau particles and tau neutrinos define the third generation. All leptons have a corresponding antiparticle indicated by a positive charge (i.e. e^+) or a bar (i.e. $\bar{\nu}_e$). Neutrinos[‡] are proven to have a very small mass **REFERENCE** and interact only using the weak force (Section **HE?**) making them inherently very hard to detect.

1.1.2 Quarks

The six quarks are called up, down, charm, strange, top and bottom quarks ((u, d), (c, s), (t, b)). One generation is made up of a particle with charge $1/3$ and one with $2/3$. This is again visualized in Figure 1.1. The difference between generations is essentially again the bare mass of the particles. Because quarks also interact through the strong force (see Section **????**) they combine into *hadrons*[§] of which the nucleons are best known. Because of their color charge and the intrinsic behaviour of the strong force quarks cannot be observed freely: they always combine into color neutral particles, a property called *confinement*. When a hadron, with it's constituent quarks, is pulled apart the attractive force between the quarks does not fall down rapidly because gluons carry color charge. When these particles are pulled apart far enough it becomes energetically more favorable to produce new quark-antiquark pairs which again combine into color neutral particles[¶]. The energy requirement for the production of new particles is far below that to separate the quarks far enough from each other to observe them separately.

Antiparticles are again denominated with a bar (i.e. \bar{u}). Because of their ability to interact with the strong force; particle accelerators in the 19???'s led to the discovery of a plethora of possible combinations. Something which is often referred to as the "particle zoo".

1.2 How particles communicate: interactions

There are four fundamental interactions known to exist: gravity and electromagnetism, which produce significant long-range forces, and the strong and weak force which only express themselves

*λεπτός (leptos) meaning thin, delicate, lightweight, or small. These particles don't need to bind to each other, which keeps them "thin" in a certain sense. Originally leptons were considered the "light" particles and hadrons the "heavy" particles, but the discovery of the tau lepton in 1975 broke that rule

[†]This characteristic of often referred to as *lepton universality* [??]

[‡]The name is a wordplay. The Italian word for neutron (neutrono) sounds like the word neutral (neutro) with an augmentative suffix (-one) tacked on the end. That is, it sounds something like "big neutral" to Italian ears. Replace the augmentative suffix -one with the diminutive suffix -ino and you have a "little neutral", which is a good description of what a neutrino is — a diminutive neutral particle.

[§]αδρός (adros) meaning thick, robust, massive, or large. This name alludes to the ability of the point-like quarks to bind together and form particles that are "thick" in a certain sense.

[¶]This process is called *hadronization* and results into the production of "jets" in particle accelerators [34]




Figure 1.1: info from: <https://home.cern/about/physics/standard-model>

Interaction	Gravitation	Weak	Electromagnetic	Strong	
		Electroweak		Fundamental	Residual
Acts on:	Mass/Energy	Flavor	Electric charge	Color charge	Atomic nuclei
Experiencing	All particles	All fermions	Electrically charged	Quarks, Gluons	Hadrons
Mediation:	Not yet observed	W^+, W^-, Z	γ	Gluons	π, ρ, ω mesons
Strength	10^{-35}	10^{-11}	1	10^2	/
Long-distance	$\frac{1}{r^2}$	$\frac{1}{r} e^{-m_{W,Z} \cdot r}$	$\frac{1}{r^2}$		r^*

Table 1.1: Bla bla bla. Ref: <https://web.archive.org/web/20160304133522/> <https://www.pha.jhu.edu/> dfehling
A nice webpage explaining... :<https://profmattstrassler.com/articles-and-posts/particle-physics-basics/the-known-forces-of-nature/the-strength-of-the-known-forces/>
<http://web.mit.edu/sahughes/www/8.022/lec01.pdf>

at (sub)atomic distances and govern nuclear interactions. These are explained in more detail below and an overview is given in Table 1.2.

Particles interact with each other through the exchange of *gauge bosons* or *force carriers*^{||}. Gauge bosons are bundles of energy (*quanta*) and can be seen as excitations of one of the force fields. Fields are a mathematical approach used by physicists to describe what we observe in experiments.

Although the use of fields is very natural, the concept might feel a bit unfamiliar. In the following the known forces are described in more detail. Although it plays less of a role in subatomic physics, the theory of gravity is added for completeness but is mainly used to explain the concept of fields more in depth.

Gravity

Gravity (from Latin/old French: *gravitas*/grave, “weighty, heavy”) is the phenomenon in which objects with mass are attracted to each other. Gravitation is famously described by the general theory[†] of general relativity proposed by Albert Einstein in 1915. Compared to the other forces

^{||}A classic way of looking at force carriers is imagining two people standing in a boat. The force carrier is a heavy ball which can be thrown from one person to the other. Doing so both persons will move in opposite direction.

[†]A scientific theory is an explanation of an aspect of the natural world that can be repeatedly tested, in accordance with the scientific method, using a predefined protocol of observation and experiment. Established scientific theories have withstood rigorous scrutiny and embody scientific knowledge. [REF WIKI?](#)

gravity is intrinsically very weak* and is not described in the SM (see Section???). Because of this, gravity is often left out in discussions of particle physics experiments. It can however be used to explain the concept of a field in a very natural way.

First published on July 5th 1686 in Newton's work *Philosophiae Naturalis Principia Mathematica* ("the Principia"), a very good description of gravitation was already well known centuries ago. The equation of the force exerted by two massive bodies takes the following form:

$$F = G \frac{m_1 m_2}{r^2} \quad (1.1)$$

where F is the gravitational force acting between two objects, m_1 and m_2 are the masses of the objects, r is the distance between the centers of their masses, and G is the gravitational constant.

Newton's law states that two massive bodies will exert a force onto one another proportional to their masses but inversely proportional to the square of the distance between them. Newton realized this would mean that at any given instant in time this would mean that all massive objects in the universe would know of every other object in the universe where it is located†. Because of this Newton himself believed his explanation could not be the final answer. The answer is fully described in Einstein's work **RELATIVITY** but was already explained by Pierre-Simon Laplace in 1783. Gravitation is the slope of a field that pervades space and because of this one only needs to know the value of the field in a local region to calculate the attractional force. Massive objects do not "feel" each other but distort space and time in such a way that objects are attracted "fall" towards each other. Field theory makes it able to treat the laws of physics as local instead of action at a distance.

To date it is not possible for gravity to be described in the framework of quantum field theory like the other fundamental forces in a compatible way with the theory of general relativity. The gauge bosons from such a quantum field theory for gravity are referred to as *gravitons*.

Electromagnetism

The electromagnetic field (from Ancient Greek: ἥλεκτρον *ēlektron*, "amber", and μαγνῆτις λίθος, *magnetislithos*, which means "Magnesian stone")‡ presents itself in the electrical and magnetical forces. In the late 1870's the publication of James Clerk Maxwell's *A Treatise on Electricity and Magnetism* showed that the interactions of negative and positive charges are mediated by one force. Particles carrying a quantity (charge) of one of these forces can attract each other or repel.

Similar to the theory of gravity, the electromagnetic field pervades all around us and the interaction of nuclei, which have a positive electric charge, and electrons makes up most of what is described in chemistry. The force carrier of electromagnetism is called a *photon*, or in other words: light.

Weak force

The weak force is one aspect of the overarching electroweak theory which combines electromagnetism and the weak force. As opposed to gravity and electromagnetism it only takes place at very small subatomic distances§. One well known phenomenon that is described by the weak force is *beta decay* in which free neutrons decay into protons and produce an extra electron and anti-electron neutrino. Another beautiful example of the weak force is the driving mechanism in the Sun's thermonuclear process which makes it shine. This process cannot be explained by chemical processes but with the fusion of hydrogen into deuterium. Two protons are squeezed together into a He atom which consists of a proton and a neutron. The conversion of the proton into a neutron can be explained by the weak force.

*Two magnets that fit in the palm of your hand can deliver a force which is of similar strength than what the whole of Earth exerts on a human body.

†Imagine the attraction of the Moon towards the Earth: how are both "communicating" to each other?

‡In 1641 Athanasius Kircher titled one of the chapters in his book Magnus [BLABLABLA]: "Elektromagnetismos i.e. On the Magnetism of amber, or electrical attractions and their causes" in which the ???

§The reason being that the force carriers are massive, see more info in section???

The force carriers of the weak force are the W^+ , W^- and Z bosons. Because of the mass of these particles the coupling of particles with the weak field is inversely proportional to the square of their mass the force seems to be very weak, hence it's name*.

The weak force also carries some peculiar properties which are unique in a number of respects:

- It is the only theory that violates parity symmetry and even does so maximally.
- It's force carriers are massive as opposed to all other force carriers.
- It is the only force capable of changing quarks from one family into a quark of another family.

Strong force

As indicated in Section 1.1 nuclei are made up of protons and neutrons. However, the forces described in this section up to now cannot explain how they can make up a stable combination. The positive/neutral electromagnetic charge of the protons/neutrons would even suggest the opposite. Protons and neutrons are made up by quarks which carry a quantity which is called *color charge*. Particles carrying a color charge participate in interactions of the strong force. Due to the principle of *self interaction* the strong force only manifests itself on very small scales[†]. When nucleons are squeezed together (either due to high temperatures or pressure) and come close enough the quarks that make up the nucleons interact and make up the binding energy between nucleons. The force carriers of the strong force are called *gluons* which carry a color charge themselves and are massless.

Aside from holding nucleons together the strong force is also responsible for around 99% of the mass of the nucleon mass. The binding energy (which includes the kinetic energy of the quarks and the energy of the gluon fields that bind the quarks together).

A note about bosons

As opposed to fermions, which obey Fermi-Dirac REF statistics and cannot occupy the same quantum state, bosons follow Bose-Einstein statistics[‡]. Bosons carry integer spins ($s = 0, 1, 2$, etc.), fermions carry half-integer spins ($s = 1/2, 3/2$, etc.). As a result, bosons have no problem occupying the same place at the same time. (More formally, two or more bosons may be described by the same quantum numbers.) As the particles that make up light and other forms of electromagnetic radiation, photons are the bosons we have the most direct experience with. In our everyday experience, we never see beams of light crash into one another. Photons can go through each other with no effect.

1.3 The Standard Model in theory

A lot from Mandel and Saw: QFT

The Standard Model is a *quantum field theory*, meaning its fundamental objects are fields of a quantum nature which are defined at all points in spacetime. These fields are

- fermion fields, ψ , which account for “matter particles”;
- electroweak boson fields, W^1, W^2, W^3 and B ;
- gluon field, G^a ; and
- Higgs field, ϕ

Quantum field theory treats particles as excited states of one of these underlying fields, so called *field quanta*. The difference between classical and quantum fields is that they are operator-valued. Classical fields can in principle take on distinct values at each point in space whereas a quantum field accommodates observations of quantum mechanics as:

- energies are quantized, meaning that only discrete energy values are possible,
- these discrete energy levels are equally spaced,

*Iets van Fermi? Dat hij dacht dacht het een eenpuntsinteractie was?

[†]As opposed to the weak force where the short distance is explained due to the mass of the force carriers.

[‡]The name boson originates from Paul Dirac REFCOMMENTED OUT who wanted to commemorate the contributions of Indian physicist Satyendra Nath Bose who, together with Albert Einstein, theorized the characteristics of elementary particles that follow the Bose-Einstein statistics.

- the lowest achievable energy is not equal to absolute zero, but has a zero-point energy* (REFERENCE)?.

REFERENCE: <https://arxiv.org/pdf/hep-ph/0609174.pdf>, chapter 2 The dynamics of the quantum state and the fundamental fields are determined by the Lagrangian density \mathcal{L} . Writing the time and space coordinates in the form $(t, \mathbf{x}) = (x^0, x^1, x^2, x^3) = x^\mu$ the equations of motion of these fields can be written as:

$$\frac{\partial}{\partial x_\mu} \left[\frac{\partial \mathcal{L}}{\partial (\partial \phi / \partial x^\mu)} \right] - \frac{\partial \mathcal{L}}{\partial \phi} = 0, \quad (1.2)$$

which follow from the principle of least action REFERENCE. The lagrangian function depends on the fields and how these fields change in spacetime: $\mathcal{L}(\phi, \nabla\phi)$. Quantization of these fields can be obtained by interpreting the coordinates and momenta as Heisenberg operators, and subjecting these to canonical commutation relations REFRENENCE MANDEL AND SHAW.

Furthermore the Standard Model is a gauge theory in which the Lagrangian is invariant under certain Lie groups (referred to as the symmetry group or the gauge group of the theory) of local transformations. For quantized gauge groups the quanta of the gauge fields are referred to as *gauge bosons*. A gauge theory is a mathematical model that has a gauge freedom of some of the mathematical degrees of freedom are redundant. In other words: different mathematical expressions describe the exact same physical system and in that sense unphysical. An experiment could never uniquely determine their values, even in principle. If the phase of the wavefunction is changed by a different amount at each point in spacetime and the physics remains unchanged, the Lagrangian is said to follow a *local phase symmetry*.[†].

The Standard Model is defined by the local $SU(3) \times SU(2) \times U(1)$ gauge symmetry. Each factor gives rise to three fundamental forces:

1.3.1 **SU(3): quantum chromodynamics**

The quantum chromodynamics (QCD) sector defines the interactions between quarks and gluons. Since leptons do not carry colour charge they do not participate in this interaction. The Dirac Lagrangian of the quarks coupled to the gluon fields is given by

$$\mathcal{L}_{QCD} = \sum_{\psi} \bar{\psi}_i \left(i\gamma^\mu \left(\partial_\mu \delta_{ij} - ig_s G_\mu^a T_{ij}^a \right) - m_\psi \delta_{ij} \right) \psi_j - \frac{1}{4} G_{\mu\nu}^a G_a^{\mu\nu}, \quad (1.3)$$

where

ψ_i is the Dirac spinor of the quark field, where $i = r, g, b$ represents the color charges,

γ^μ are the Dirac matrices REFERENCE,

G_μ^a is the 8-component ($a = 1, 2, \dots, 8$) $SU(3)$ gauge field,

T_{ij}^a are the 3×3 Gell-Mann matricesREF, generators of the $SU(3)$ color group,

$G_{\mu\nu}^a$ are the field strength tensors for the gluons,

g_s is the strong coupling constant.

1.3.2 **SU(2) × U(1): electroweak**

Definitie van electroweak charge, Y?

The electroweak sector is a Yang-Mills gauge theoryREF with the symmetry group $SU(2)_L \times U(1)$. The Lagrangian is given by

*This is in accordance with the well known Heisenberg uncertainty principle which states that because of the zero-point energy, the position and momentum of a particle are not fixed but have a small range of variance: $\sigma_x \sigma_p \geq \frac{\hbar}{2}$.

[†]See appendix A

$$\begin{aligned}
\mathcal{L}_{EW} &= \sum_{\psi} \bar{\psi} \gamma^{\mu} \left(i\partial_{\mu} - g' \frac{1}{2} Y_W B_{\mu} - g \frac{1}{2} \vec{\tau}_L \vec{W}_{\mu} \right) \psi - \frac{1}{4} W_a^{\mu\nu} W_a^{\mu\nu} - \frac{1}{4} B^{\mu\nu} B_{\mu\nu} \\
&= \bar{Q}_i i D_{\mu} \gamma^{\mu} Q_i + \bar{u}_i i D_{\mu} \gamma^{\mu} u_i + \bar{d}_i i D_{\mu} \gamma^{\mu} d_i + \bar{L}_i (i D_{\mu} \gamma^{\mu}) L_i + \bar{e}_{R,i} (i D_{\mu} \gamma^{\mu}) e_{R,i} \\
&\quad - \frac{1}{4} W_a^{\mu\nu} W_a^{\mu\nu} - \frac{1}{4} B^{\mu\nu} B_{\mu\nu},
\end{aligned} \tag{1.4}$$

where

B_{μ} is the U(1) gauge field,

Y_W is the weak hypercharge - the generator of the U(1) group,

\vec{W}_{μ} is the 3-component SU(2) gauge field,

$\vec{\tau}_L$ are the Pauli matrices - infinitesimal generators of the SU(2) group - with subscript L to indicate that they only act on left-chiral fermions,

g' (weak isospin) and g (weak hypercharge) are the U(1) and SU(2) coupling constants respectively,

$W^{a\mu\nu}$ ($a = 1, 2, 3$) and $B^{\mu\nu}$ are the field strength tensors for the weak isospin and weak hypercharge fields

Q, u and d are the left-handed doublet, right-handed singlet up right-handed singlet down quark fields,

L and e are the left-handed doublet and right-handed singlet electron fields.

The field strengths are given by

$$\begin{aligned}
W_{\mu\nu}^a &= \partial_{\mu} W_{\nu}^a - \partial_{\nu} W_{\mu}^a + g \epsilon^{abc} W_{\mu}^b W_{\nu}^c, \\
B_{\mu\nu} &= \partial_{\mu} B_{\nu} - \partial_{\nu} B_{\mu},
\end{aligned}$$

and the covariant derivative for the left- and right-handed leptons by,

$$\begin{aligned}
D_{\mu} L_L &= \left(\partial_{\mu} - i \frac{g}{2} \tau_a W_{\mu}^a - ig' Y B_{\mu} \right) L_L, \\
D_{\mu} e_R &= \left(\partial_{\mu} - i \frac{g'}{2} Y B_{\mu} \right) e_R,
\end{aligned}$$

where T_a

It is worth noting no terms are included for fermion masses. These would have the form of $m \bar{\psi} \psi$ but are forbidden as they would break the $SU(2)_L \times U(1)$ gauge invariance. Neither is it possible to add explicit mass terms for the U(1) and SU(2) gauge fields.

1.3.3 Brout-Englert-Higgs mechanism

To come to a viable description of the elementary particles one is required to introduce masses into a chiral theory. The masses of the W and Z bosons are explained by the use of the Brout-Englert-Higgs mechanism formulation. Introducing one or more scalar fields, the Higgs fields, which can acquire a vacuum expectation value, it is possible to spontaneously break a symmetry in the Lagrangian. We say that electroweak symmetry is broken down to electromagnetism. According to the Goldstone theorem REF, every spontaneously broken continuous symmetry results in a massless scalar particle, the Goldstone boson. Hence, the number of Goldstone bosons in a theory is equal to the number of broken generators of the symmetry group.

Since the electroweak theory after symmetry breaking should contain three massive gauge bosons (W^+, W^- and Z) the scalar fields of the Higgs fields should contain at least three degrees of freedom. The simplest approach to do this is by introducing a complex, scalar $SU(2)$ doublet Φ with positive hypercharge ($Y = \frac{1}{2}$),

$$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi_1 + i\phi_2 \\ \phi_3 + i\phi_4 \end{pmatrix}, \tag{1.5}$$

Similar to the $SU(2)$ symmetry of the EW theory four new gauge particles are introduced: H^+, H^-, H^0 and H .




Figure 1.2: In this example the Higgs potential is illustrated in function of a complex scalar field (2D). The principal is the same for a complex scalar doublet but a lot harder to visualize. Top left: the Higgs potential with $\mu^2 > 0$, there is no vacuum expectation value. Top right: the scalar field will move to the lowest possible energy state. Bottom left: a flat direction in the potential corresponds to a massless Goldstone mode (remember there are two extra scalar fields meaning there are a total of 3). Bottom right: the concave shape of the potential near the minimum defines the Higgs boson mass.

Massive bosons

Having introduced this scalar doublet one needs to add the corresponding Lagrangian term to the electroweak Lagrangian from eq. 1.4,

$$\mathcal{L}_S = (D^\mu \Phi)^\dagger (D_\mu \Phi) - V(\Phi), \quad \text{with } V(\Phi) = \mu^2 \Phi^\dagger \Phi + \lambda (\Phi^\dagger \Phi)^2. \quad (1.6)$$

The first term is the kinetic term and the second corresponds to the potential of the scalar field*. The quartic term in the potential needs to be positive to ensure an absolute minimum in the Lagrangian. The quadratic term can either be positive or negative, depending on if $\mu^2 > 0$ or $\mu^2 < 0$. This is illustrated in Fig. 1.2. In the former case the scalar potential has an absolute minimum at the origin:

$$\langle 0 | \Phi | 0 \rangle \equiv \Phi_0 = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \quad (1.7)$$

From eq. 1.6 and 1.7 one can see that it the kinetic term does not give rise to massive particles†. In the case of $\mu^2 < 0$, the minimum is no longer located at the origin of the fields: $\partial_{|\Phi|} V(|\Phi|) = 0$

*The form of the potential is not known from first principles but is the simplest form that can explain the spontaneous symmetry breaking mechanism.

†Substitute $v = 0$ in eq. 1.13.

for $|\Phi| = \sqrt{-\frac{\mu^2}{2\lambda}}$, hence one possible solution is*

$$\langle 0 | \Phi | 0 \rangle \equiv \Phi_0 = \begin{pmatrix} 0 \\ \frac{v}{\sqrt{2}} \end{pmatrix}, v = \sqrt{-\frac{\mu^2}{\lambda}}. \quad (1.8)$$

v is referred to as the *vacuum expectation value* to reflect that the Higgs field is always “on”. To investigate the terms we can expand the field around the minimum:

$$\Phi(x) = \begin{pmatrix} \theta_2(x) + i\theta_1(x) \\ \frac{v+H(x)}{\sqrt{2}} - i\theta_3(x) \end{pmatrix} = e^{i\theta_a \tau_a} \begin{pmatrix} 0 \\ \frac{v+H(x)}{\sqrt{2}} \end{pmatrix}. \quad (1.9)$$

Implementing this into eq. 1.6 would yield the existence of unphysical fields $\phi_{1,2,3}$ that give rise to three extra degrees of freedom that were not present in the original Lagrangian[†]. Since a change of variables cannot alter the number of d.o.f. of a system one can conclude that three fields do not represent physical fields. They can be removed by fixing a gauge, the unitary gauge, which breaks the original symmetry of the system!

$$\Phi(x) \rightarrow e^{-i\theta_a \tau_a} \Phi(x) = \begin{pmatrix} 0 \\ \frac{v+H(x)}{\sqrt{2}} \end{pmatrix}, \quad (1.10)$$

where we have introduced a new scalar field $H(x)$. After inserting this in the kinetic part of the scalar Lagrangian (eq. 1.6), and redifining the gauge fields as

$$\begin{aligned} W_\mu^\pm &= \frac{1}{\sqrt{2}} (W^1 \mp iW_\mu^2), \\ Z_\mu &= \frac{1}{g^2 + g'^2} (gW_\mu^3 - g'B_\mu), \\ A_\mu &= \frac{1}{g^2 + g'^2} (gW_\mu^3 + g'B_\mu), \end{aligned} \quad (1.11)$$

we find for the kinetic part of the scalar Lagrangian:

$$|D_\mu \Phi|^2 = \frac{1}{2} (\partial_\mu H)^2 + \frac{1}{2} g^2 (v + H)^2 W_\mu^+ W^{-\mu} + \frac{1}{8} (v + H)^2 (g^2 + g'^2) Z_\mu Z^\mu. \quad (1.12)$$

Since mass terms enter this equation in the general form of $M_W^2 W_\mu W^\mu$ for the W -bosons and $\frac{1}{2} M_Z^2 Z_\mu Z^\mu$ for the Z boson, the mass terms of the gauge bosons after spontaneous symmetry breaking can be written down as:

$$\begin{aligned} M_W &= \frac{1}{2} vg, \\ M_Z &= \frac{1}{2} v \sqrt{g^2 + g'^2}, \\ M_A &= 0, \end{aligned} \quad (1.13)$$

where it is clear that the photon remains massless[‡].

*It is not possible for the charged part of the fields to have a vev as this would not be in agreement with electromagnetism.

[†]In eq. 1.6 the vector fields are massless and each give rise to 2 d.o.f. The vev would make the three vector fields massive, thus adding 3 d.o.f. and introduce three unphysical fields.

[‡]Because the W and Z bosons are massive it costs energy to produce them and so the weak force is only really effective over a short distance. This is in contrast to the massless photons that result into a long range electromagnetic force. Thus the Higgs is responsible for the “weakness” of the weak force.

Corollary 1.3.1 — Eating the gauge bosons. A vev can give rise to massless Goldstone bosons. These particles correspond to the infinite possibilities of it's phase in the potential. However, this would also lead to extra degrees of freedom in the Lagrangian. Therefore one has to fix a gauge resulting in the disappearance of the bosons and making the vector bosons of the original fields massive.

Specifically, the two massless W^1, W^2 bosons in the electroweak theory (2×2^a polarizations = 4 d.o.f.) and two charged Higgses (2 d.o.f.) sum to a total of six degrees of freedom. In the broken theory, we have two massive W^+, W^- bosons (2×3 polarizations) which again total to six degrees of freedom.

Similarly, the W^3, B (2×2 d.o.f.) and H^0 (2 d.o.f.) combine into the neutral Z (massive, 3 d.o.f.), the γ (2 d.o.f.) and the scalar H (1 d.o.f.).

Similar to the nucleons having a mass which is much greater than the summed mass of it's constituents the Higgs fields gives rise to the mass of these gauge bosons. In the case of nucleons it is primarily the potential energy of the strong force that is responsible for the total mass following $E = mc^2$. The coupling of the gauge bosons to the Higgs field gives a sense of inertia to the particle. The particle does not float freely in vacuum but interacts with the ever present Higgs field (with a vev $\neq 0$) making it massive.

^aA massless particle cannot have a third polarization, similar to a photon: it is traveling at the speed of light so it cannot have a polarization in the direction of propagation, only longitudinal.

Using the potential term in eq. 1.6 together with the vev in eq. 1.10 we find for the Lagrangian of the Higgs boson:

$$\mathcal{L}_H = \frac{1}{2} (\partial_\mu H) (\partial^\mu H) - \lambda v^2 H^2 - \lambda v H^3 - \frac{\lambda}{4} H^4. \quad (1.14)$$

The first term again corresponds to the kinetic term whereas the third and forth refer to the three- and four-point self-interactions of the Higgs respectively. Scalar masses have the general form $\frac{1}{2}m\phi^2$; the Higgs boson mass is thus equal to

$$m_H = 2\lambda v^2 = -2\mu^2, \quad (1.15)$$

and needs to be determined experimentally.

Working through the interaction terms of the Lagrangian, one can show that the electric charge e is related to the couplings of the weak isospin g and hypercharge g' .

$$e = g \sin \theta_W = g' \cos \theta_W, \quad (1.16)$$

where the Weinberg angle is denoted as θ_W and indicates the magnitude of rotation of the boson planes after spontaneous symmetry breaking:

$$\begin{pmatrix} \gamma \\ Z \end{pmatrix} = \begin{pmatrix} \cos \theta_W & \sin \theta_W \\ -\sin \theta_W & \cos \theta_W \end{pmatrix} \begin{pmatrix} B^0 \\ W^0 \end{pmatrix}, \quad (1.17)$$

and is related to the weak isospin and hypercharge:

$$\cos \theta_W = \frac{g}{\sqrt{g^2 + g'^2}} \text{ and } \sin \theta_W = \frac{g'}{\sqrt{g'^2 + g^2}}. \quad (1.18)$$

references: <https://www.quantumdiaries.org/2011/11/21/why-do-we-expect-a-higgs-boson-part-i-electroweak-symmetry-breaking/> and <https://www.nikhef.nl/~ivov/HiggsLectureNote.pdf>

Massive fermions

A term like $-m\bar{\psi}\psi = -m [\bar{\phi}_L\psi_R + \bar{\psi}_R\phi_L]$, where we have decomposed the equation into the left- and right-handed chiral states[§] is not gauge in the Lagrangian. The left-handed fermions

[§] $\psi_L = P_L\psi = \frac{1-\gamma^5}{2}\psi$ and $\psi_R = P_R\psi = \frac{1+\gamma^5}{2}\psi$

from an isospin doublet and the right-handed fermions form isospin singlets. They transform differently under $SU(2)_L \times U(1)_Y$:

$$\begin{aligned}\chi_L &\rightarrow \chi'_L = \chi_L e^{i\tau_L^z \vec{W} + i\alpha Y} \\ \psi_R &\rightarrow \psi'_R = \psi_R e^{i\alpha Y}\end{aligned}. \quad (1.19)$$

It is possible for the fields to couple to the complex Higgs doublet defined in 1.5 by adding Yukawa couplings. This results into terms which are *singlets* under $SU(2)_L$ and $U(1)_Y$:

$$\mathcal{L}_{Yuk} = \lambda_e \overline{L_L} \Phi e_R - \lambda_d \overline{Q_L} \Phi d_R - \lambda_u \overline{Q_L} \tilde{\Phi} u_R + h.c. \quad (1.20)$$

where we have introduced the conjugate of Φ , $\tilde{\Phi} = i\tau_2 \Phi^*$ which has a negative hypercharge. After spontaneous symmetry breaking (eq. 1.10), we find:

$$\begin{aligned}L_{Yuk} &= -\frac{1}{\sqrt{2}} \lambda_e (\bar{\nu}_e, \bar{e}_L) \begin{pmatrix} 0 \\ v + H(x) \end{pmatrix} e_R + \dots \\ &= -\frac{1}{\sqrt{2}} \lambda_e (v + H(x)) \bar{e}_L e_R + \dots,\end{aligned} \quad (1.21)$$

where we highlighted only the electron part. Fermion mass terms have the general form $m_f \bar{f}_L f_R + h.c.$. Therefore one finds:

$$m_e = \frac{\lambda_e v}{\sqrt{2}}, \quad m_u = \frac{\lambda_u v}{\sqrt{2}}, \quad m_d = \frac{\lambda_d v}{\sqrt{2}}. \quad (1.22)$$

The mass of the fermions is again not predicted as the Yukawa coupling parameters are free parameters.

1.3.4 Particle mixing

In equation 1.20 we introduced Yukawa coupling constants to explain the mass of fermions. In its most general realizations these couplings are not constants but matrices. This will introduce possible mixing of *flavor? eigenstates* into different *mass eigenstates*. Let us write out the second term in equation 1.20

$$\begin{aligned}\lambda_d \overline{Q_L} \Phi d_R &= Y_{ij}^d Q_{Li}^I \bar{\Phi} d_{Rj}^I = Y_{ij}^d \overline{(\text{up-type down-type})_{Li}^I} \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} (\text{down-type})_{Rj}^I \\ &= \begin{pmatrix} Y_{11} \overline{(u\ d)_L^I} \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} & Y_{12} \overline{(u\ d)_L^I} \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} & Y_{13} \overline{(u\ d)_L^I} \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} \\ Y_{21} \overline{(c\ s)_L^I} \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} & Y_{22} \overline{(c\ s)_L^I} \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} & Y_{23} \overline{(c\ s)_L^I} \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} \\ Y_{31} \overline{(t\ b)_L^I} \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} & Y_{32} \overline{(t\ b)_L^I} \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} & Y_{33} \overline{(t\ b)_L^I} \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} \end{pmatrix} \cdot \begin{pmatrix} d_R^I \\ s_R^I \\ b_R^I \end{pmatrix}\end{aligned} \quad (1.23)$$

where the superscript I implies that the fermion fields are expressed in the *interaction (flavor)* basis. The subscript i stands for the three generations.

This means that after symmetry breaking the quark mass terms break down into

$$\begin{aligned}-\mathcal{L}_{Yuk}^{\text{quarks}} &= Y_{ij}^d \overline{d_{Li}^I} \frac{v}{\sqrt{2}} d_{Rj}^I + Y_{ij}^u \overline{u_{Li}^I} \frac{v}{\sqrt{2}} u_{Rj}^I + \dots \\ &= M_{ij}^d \overline{d_{Li}^I} d_{Rj}^I + M_{ij}^u \overline{u_{Li}^I} u_{Rj}^I + \dots\end{aligned} \quad (1.24)$$

where we have omitted the hermitian conjugate terms and the Higgs field interaction terms. Note that the u - and d -terms in the equation still each represent the three up- and down-type quarks

repectively. There is mixing between the flavor fields as there is no reason why the matrix M should be diagonal*.

To obtain proper mass terms one has to diagonalize the mass matrices M^u and M^d and find proper eigenstates. We introduce unitary matrices V as follows

$$\begin{aligned} M_{diag}^d &= V_L^d M^d V_R^{d\dagger} \\ M_{diag}^u &= V_L^u M^u V_R^{u\dagger} \end{aligned} \quad (1.25)$$

which can be done when the matrices V are unitary ($V_L^{d,u\dagger} V_L^{d,u} = \mathbb{1}$). Equation 1.24 can now be expressed as follows:

$$\begin{aligned} -\mathcal{L}_{Y_{uk}}^{\text{quarks}} &= \overline{d_{Li}^I} M_{ij}^d d_{Rj}^I + \overline{u_{Li}^I} M_{ij}^u u_{Rj}^I + \dots \\ &= \overline{d_{Li}^I} V_L^{d\dagger} V_L^d M_{ij}^d V_R^d V_R^{d\dagger} d_{Rj}^I + \overline{u_{Li}^I} V_L^{u\dagger} V_L^u M_{ij}^u V_R^u V_R^{u\dagger} u_{Rj}^I + \dots \\ &= \overline{d_{Li}^I} \left(M_{ij}^d \right)_{diag} d_{Rj}^I + \overline{u_{Li}^I} \left(M_{ij}^u \right)_{diag} u_{Rj}^I + \dots \end{aligned} \quad (1.26)$$

where the V matrices have been absorbed in the quark flavor eigenstates and have formed mass eigenstates. These mass eigenstates, which are the eigenstates one sees in experiments, couple differently to the gauge fields of the weak interaction. Working out one term from equation 1.4 the mixing of the flavor eigenstates is clearly visible

$$\begin{aligned} \mathcal{L}_{kinetic}(Q_L) &= i \overline{Q_{Li}^I} \gamma_\mu D^\mu Q_{Li}^I \\ &= \frac{g}{\sqrt{2}} \overline{u_{Li}^I} \gamma_\mu W^{-\mu} d_{Li}^I + \frac{g}{\sqrt{2}} \overline{d_{Li}^I} \gamma_\mu W^+ \mu u_{Li}^I + \dots \\ &= \frac{g}{\sqrt{2}} \overline{u_{Li}^I} \left(V_L^u V_L^{d\dagger} \right)_{ij} \gamma_\mu W^{-\mu} d_{Li}^I + \frac{g}{\sqrt{2}} \overline{d_{Li}^I} \left(V^d V^{u\dagger} \right)_{ij} \gamma_\mu W^+ \mu u_{Li}^I + \dots \end{aligned} \quad (1.27)$$

The combination of matrices $\left(V_L^d V_L^{u\dagger} \right)_{ij}$, a unitary 3×3 matrix is known as the *Cabibbo-Kobayashi-Maskawa (CKM)* mixing matrix. By convention, the interaction and flavor eigenstates of the up-type quarks are chosen to be equal. The down-type quarks are therefore chosen to be rotated:

$$\begin{aligned} u_i^I &= u_i \\ d_i^I &= V_{CKM} d_i \end{aligned} \quad (1.28)$$

or explicitly:

$$\begin{pmatrix} d^I \\ s^I \\ b^I \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}. \quad (1.29)$$

Because the choice of the global phases of the quark fields is arbitrary and the matrix is unitary, the nine unknown complex elements can be reduced to three real numbers and one phase[†]. The matrix is most often written as:

$$V = \begin{pmatrix} c_{12}c_{13} & & s_{12}c_{13}s_{13}e^{-i\delta} & \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} - c_{12}s_{23} - s_{12}s_{23}s_{13}e^{i\delta} & & c_{23}c_{13}, & \end{pmatrix} \quad (1.30)$$

*The question and answer of flavor/mass mixing can be put as: “Why is there mixing? Because it can.”

[†]This phase is responsible for *CP-violation*.

where $c_{ij} = \cos(\theta_{ij})$ and $s_{ij} = \sin(\theta_{ij})$ for $i < j = 1, 2, 3$ and δ is the CP -violating phase.

The mixing of the flavor quantum states is necessary to explain charged current interactions changing the strangeness with one [45] and CP -violating processes [32].

Without going into detail it is worth noting that a similar matrix exists that connects the lepton flavor and mass eigenstates. In contrast to the quarks, the down-type interaction doublet states are chosen to be the same as the mass eigenstates. The mixing of the mass and interaction eigenstates is in the neutrino sector. This matrix is known as the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix [53].

1.4 A success story

Over the course of multiple decades the Standard Model was built up into an extremely comprehensive theory. The first building blocks necessary for its construction came from experiments in the early 19th century when its quantum characteristics became more apparent. The theory was formulated as a gauge theory in the 1960s and 1970s and for decades it has been rigorously tested and checked, leading to extremely accurate experimental precision measurements which agree with the theory. Apart from precision measurements it has lead to predictions of particles and their interactions which could only be tested years or decades after they were first proposed. In the following I give a brief overview of some astonishing experimental results.

1.4.1 Wave-particle duality

One of the most striking features of the SM is that particles may be partly described in terms of not only particles, but also of waves. The classical concepts of “particles” or “waves” is insufficient in describing the behaviour of quantum-scale objects. In the late 18th and early 19th century physicist were puzzled by the successful approaches to well known problems of electromagnetic phenomena, which were widely accepted as fields, with quantizations (particles). Black body radiation and the photoelectric effect are described below. In QFT particles are defined as excited states of a field. The wave nature is evident in the calculated *probability distributions* for a given reaction.

Photoelectric effect

At the close of the 19th century J.J. Thomson found that electricity is caused by *particles* that can fly through vacuum. But, since electromagnetism was known to be a wave that is generated by a changing electric or magnetic *field*, a particle description of electricity and charge was, in a way, flawed by construction. A first step towards quantization came when the ultraviolet catastrophe seemed to be solved if black-body radiation is not explained by the classical equipartition theorem (Appendix B) but a quantization of the electromagnetic field: Planck’s law. Planck denounced these particles of light as a limitation of his approximation, not a property of reality. It was only later, when Einstein combined this theory with the photoelectric effect (which was not understood), that light quanta/photons became more accepted.

The photoelectric effect describes that when a material is shined upon by light, electrons or other free carriers can be emitted. In a classical theory perspective, an alteration in the intensity of light would induce changes in the kinetic energy of the particles emitted from the material. Instead, these particles are dislodged only by the impingement of photons when those photons reach or exceed a threshold frequency/energy. Changing the intensity of the light bundle only changes the amount of particles released, but does not alter its kinetic energy. This is easy to explain in the quantum view and resulted in the Einstein’s Nobel Prize in 1921.

Double-slit experiment

The double-slit experiment was first performed by Thomas Young in the early 19th century. If a beam of light is shined through two slits of a screen onto a second screen behind the first an interference pattern can be seen as illustrated in Fig. 1.3. This proved the wavelike nature of light.

In later years similar experiments for electrons were done to prove the wavelike properties of particles [39, 58]. The same double-slit experiment was finally performed in the 60s [50] and for




Figure 1.3: Left: Schematic view of interference pattern as in Young’s experiment[66]. Right: (top) a single 20 ms time frame showing several individual photon arrival points. Right: (bottom) image obtained after integrating for a few seconds: the diffraction pattern emerges[67].

single electrons in 1974 [54]. Single photon [67] (shown in Fig. 1.3) and single electron double-slit experiments show the same results, illustrating the wave probability interpretation of particles and fields.

1.4.2 Predictions

By 1932, scientists knew that atoms were made up by protons, neutrons and electrons. Together with the photon a total number of four particles were known. Four grew to five when Anderson REF discovered the existence of positrons (which were predicted by Dirac REF). Then came the pion REF and muon. By the 1960s there were hundreds of “fundamental particles” with no good guiding principles to link them together. They were often referred to as the “particle zoo”.

By a series of insights by several individuals, the Standard Model as a quantum field theory became more widely accepted. Since then, the SM has predicted the results of experiment after experiment. Some of them are:

- Neutral weak currents. Postulated by A. Salam, S. Weinberg and S. Glashow, the theory of electroweak interactions predicted the existence of a new type of weak interaction, in which the reacting particles do not change their charges. The first observation was made in 1973 at the European nuclear research laboratory, CERN.
- Weak gauge bosons. Again postulated by the abovementioned people. These particles were also discovered in CERN, 1984.
- Heavy quarks. To explain the CP violations in kaon decays, M. Kobayashi and T. Maskawa predicted the existence of a third generation of quarks: the *top* and *bottom* quarks. The bottom quark was discovered in 1977 at Fermilab. It took another 18 years for the top quark to be found in the same institute.
- Gluons. The gauge bosons of quantum chromodynamics were discovered in 1978 and 1979.
- Higgs boson. On July 4, 2012, physicists at CERN announced the discovery of the only fundamental particle predicted by the Standard Model which was not yet discovered.

1.4.3 Precision tests

Inconsistencies between experiment and theory can be signs of wrong or incomplete theories. One example is the *Lamb shift*, a difference in energy between two energy levels of the hydrogen atom which was not predicted by the Dirac equation. This phenomenon is explained with the

Parameter	Experimental value	Theoretical value	Standard deviation
m_t [GeV]	172.74 ± 0.46	172.96 ± 0.45	-0.5
m_W [GeV]	80.387 ± 0.016	80.358 ± 0.004	1.8
	80.376 ± 0.033		0.6
	80.370 ± 0.019		0.6
Γ_W [GeV]	2.046 ± 0.049	2.089 ± 0.001	-0.9
	2.195 ± 0.083		1.3
m_H [GeV]	125.14 ± 0.15	125.14 ± 0.15	0.0
$g_V^{\nu e}$	-0.040 ± 0.015	-0.0398 ± 0.0001	0.0
$g_A^{\nu e}$	-0.507 ± 0.014	-0.5063	0.0
τ_τ [fs]	290.75 ± 0.36	290.39 ± 2.17	0.1
$\frac{1}{2}(g_\mu - 2 - \frac{\alpha}{\pi})$	$(4511.18 \pm 0.77) \times 10^{-9}$	$(4508.63 \pm 0.03) \times 10^{-9}$	3.3
M_Z [GeV]	91.1876 ± 0.0021	91.1884 ± 0.0020	-0.4
Γ_Z [GeV]	2.4952 ± 0.0023	2.4942 ± 0.0008	0.4
$\Gamma_Z(had)$ [GeV]	1.7444 ± 0.0020	1.7411 ± 0.0008	-
$\Gamma_Z(inv)$ [MeV]	499.0 ± 1.5	501.44 ± 0.04	-
$\Gamma_Z(l^+l^-)$ [MeV]	83.984 ± 0.086	83.959 ± 0.008	-

Table 1.2: Observables compared with the SM best fit predictions. Errors are the total (experimental plus theoretical) uncertainties. Results are taken from Tables 10.4 and 10.5 in [59]

theory of quantum electrodynamics. REF. Another example.....

Because of this, experimentalists are continuously testing theoretical parameters of the Standard Model. These precision tests are most often done for the theory of QED. With the use of *renormalization theory* many parameters of the theory can be calculated in great detail. High-precision measurements of various observables have been performed at LEP 1 and SLC [5, 6, 7, 8, 64, 65] for physics at the Z-boson mass ($\sqrt{s} \approx M_Z$) and other observables at Tevatron [2, 37], LEP 2[37], ATLAS[1, 20] and CMS[35, 36]. Some of them are listed in Table 1.2. From this it is clear that the SM is very consistent with what we see in experiments.

ook hydrogen atom: anders verdwijnt electron binnen 10^{**-12} sec! precision measurements in icecube?

The most spectacular successes of renormalization theory were the calculations of the anomalous magnetic moment of the electron and the Lamb shift in the spectrum of hydrogen.

Neutrino cross-section: IceCube/Nature <https://www.nature.com/articles/nature24459>

Beautiful interplay of forces: i.e. strong force necessary to KEEP the neutrons as they are. Without it all neutrons would become protons! On the other hand; protons and neutrons would weigh the same if?

Fun fact: entire lagrangian somewhere in a footnote?

Gargamelle experiment: weak neutral currents: hint of Z boson discovery? Another decade for direct detection

Weak bosons: proof that there are 3 and only 3 families

Leuk: bestaat ook niet zoiets als negatieve zwaartekracht. Nochtans wel alle andere flavors negatief en positief

1.5 The need for physics beyond the Standard Model

Despite its incredible success many physicists believe the Standard Model is certainly not the full story. There are a number of features that seem arbitrary, but also some that cannot be explained by the theory alone. Below, I give a list of open questions:

- Why are there **three families** for both leptons and quarks?
- What is the **cause of the symmetries** we see in the Standard Model. Why is, for example, QCD not an SU(4) gauge theory?

- There are a number of **parameters** in the SM which cannot be explained by first principles. We have no good explanation why the top quark is 75 000 times heavier than the up quark. Why is the vev of the Higgs potential 246 GeV? Why is the Higgs mass 125 GeV? There are in total 19 parameters in the SM which are tuned by experiments and can be found in Table 1.3.
- Why does the Higgs potential have this **Mexican hat shape**? In other words, why is $\mu^2 < 0$ in $\lambda (\Phi^\dagger \Phi)^2 + \mu^2 (\Phi^\dagger \Phi)$ negative? Also, the vev, the mass of the Higgs boson and the mass of the fermions due to the Yukawa couplings all appear in Table 1.3. This makes us believe there is something we do not fully understand about the BEH mechanism.
- Right-handed neutrinos can be introduced into the SM. They are singlets with respect to the strong and weak interaction and would therefore not carry an electric charge, weak hypercharge or weak isospin. Due to this lack of charge, right-handed neutrinos would be extremely difficult to detect. They have Yukawa interactions with other leptons and the Higgs boson but its coupling would be extremely small. Neutrinos can become massive with Dirac mass terms in the same way charged leptons become massive in the BEH mechanism. Their **extremely small masses** suggest another mechanism in which the very light left-handed neutrinos are accompanied with extremely heavy right-handed neutrinos. This mechanism is called the Seesaw mechanism and requires the addition of Majorana mass terms*.

Aside from these, there are a number of unexplained phenomena that probably cannot be explained in a simple extension of the Standard Model but need a non-trivial approach.

For example:

- It is a natural assumption that the universe is neutral with all conserved charges. Both the SM and general relativity give no explanation on the **matter-antimatter imbalance** we see in the universe. The Big Bang is expected to produce equal amounts of matter and anti-matter, yet we see that the observable universe consists almost exclusively out of baryonic matter[†]. The most likely explanation is that in the early universe physical laws we know today were absent or have acted differently. The observed CP-violation is insufficient to account for the observed baryon asymmetry of the universe given the limits on baryon number violation.
- The stars, planets, interstellar clouds, etc. we see in space consist of baryonic matter. Assuming general relativity is the correct theory to describe gravity on cosmological scales, the Lambda-CDM model REF predicts that the matter we see is only around 15% of the total matter present in our visible universe REF PLANCK, zie Nadja. To explain the galaxy rotation curves [38], galaxy velocity dispersions [43], galaxy cluster masses [15], gravitational lensing [56] and many more, we predict that around 85% of the mass is not yet observed. This matter is referred to as **dark matter** as it cannot interact electromagnetically because it would have already been observed otherwise. No known particles in the SM can explain this phenomenon.
- Similar to dark matter the Lambda-CDM model predicts that the total energy in the visible universe should consist mostly out of a constant energy density for the vacuum called **dark energy**. 5% of the total energy consists of baryonic matter, 26% should be dark matter and the remaining 69% of dark energy is necessary to explain the expansion of the universe[‡].
- General relativity is generally accepted to describe gravity on cosmological scales. Thusfar is has not been possible to describe **gravity** on a quantum scale as is the case for the Standard Model and still be valid on very large scales. The inclusion of the graviton would

*Oscillation of the massive neutrinos is described in the PMNS matrix and adds 7 new parameters to the SM: the three mass terms $m_{\nu 1}, m_{\nu 2}, m_{\nu 3}$, the mixing angles θ_{12}, θ_{13} and θ_{23} and the CP-violating phase δ_{CP}

[†]Why are there protons, neutrons and electrons everywhere while it is perfectly possible for antiprotons and antineutrons to form atomic nuclei with positrons?

[‡]If there is only matter and the Big Bang acceleration only happened in the beginning of the creation of the universe, then one would expect the expansion to diminish due to the gravitational pull of matter. Measurements REF say the opposite is true: the universe is expanding and in an accelerating rate.

for example not recreate what is observed experimentally without other modifications to the SM which have not been observed REF. Contradictory to popular belief, it is not true that general relativity and quantum mechanics are incompatible. There is a need for a more complete theory beyond the range of their combined applicability[42].

- Why is the CP-violation in the strong interaction extremely small or even zero?
- Often referred to as a muon g-2 anomaly there are possible hints of new physics as the theoretical prediction of magnetic moment of the muon and experimental values have a small but significant offset [47].
- Why is there much more mixing in the lepton sector (PMNS) compared to the quark sector (CKM)?
- To explain the apparent quantum fluctuations on cosmological scales together with the horizon REF, flatnessREF and magnetic monopoleREF problems we have a theory of exponential expansion of space in the early universe: **cosmic inflation**. The theory states that there was between 10^{-36} and 10^{-32} seconds after the Big Bang a rapid exponential expansion happened. This could explain the apparent thermal equilibrium between parts of the visible universe which are not in causal contact with each other and the even distribution of the cosmic microwave background. The hypothetical field that is thought to be responsible for inflation, inflaton, is not yet observed and would be an extension of the Standard Model.
- With the use of renormalization theory REF it is possible to show that bare parameters should not be the same as parameters measured in experiments. These parameters, as the mass of particles, depends on the energy scale at which they are probed and physics far beyond the scope of the probed energy scale can influence these parameters. An example is the screening effect which is described in more detail in section 1.5.1. Similarly one-loop corrections to the Higgs boson mass* will have radiative corrections with a quadratic dependence on the cutoff scale. Virtual particles in one-loop corrections can have infinite momenta which should contribute to the total mass of the Higgs boson. Since we expect new physics to be present at energies close to the Planck mass ($\approx 10^{18}$ GeV) these loop corrections should push the Higgs mass to similar energy ranges. But, we see that the Higgs mass is around 125 GeV, which would mean that there are other parameters which should almost *exactly* cancel these absurdly large numbers. This is called *fine-tuning*, and it's the intuition of most physicists that this incredible fine-tuning has a deeper, yet unknown, meaning. This problem is often referred to as *the hierarchy problem*.

Many of these problems can be seen as “environmental”. Why there are three families and why so many parameters in the Standard Model have no fundamental explanation could just be because it's just the way it is. Maybe there is a multiverse, a plethora of universes with similar Standard Models, which have slightly or vastly different parameters. Some questions might even be impossible to answer because of a lack of statistics: we only have one universe and mankind has not been around very long in the timescale of the universe. This consideration might be valid but again not answers all our questions, it does not solve the question around dark matter for example.

This argument should not prevent us in trying to find a more general theory for the Standard Model and general relativity. A better explanation could be fairly simple, but infinitely hard as well. There is only one way in trying to find a better understanding: experiments.

Unification is the most popular approach in describing physics beyond the Standard Model. Unification would mean that well established theories are low-energy approximations of a more grand unified theory. Historically this has worked very well: the unification of celestial gravitation of Kepler with terrestrial gravitation of Galileo into universal gravitation, unification of electricity, magnetism and later optics into electromagnetism. Gravity was overhauled by the much broader theory of general relativity. Lastly, the birth of gauge theories have combined QED and the weak interaction into the combined electroweak theory. The similarities in QCD and the electroweak,

*Fermions and bosons are not effected by higher energy physics in the same way as a scalar particle is. There is a logarithmic dependence.

Parameter	Description	Value
m_e	Electron mass	511 keV
m_μ	Muon mass	105.7 MeV
m_τ	Tau mass	1.78 GeV
m_u	Up quark mass	1.9 MeV
m_d	Down quark mass	4.4 MeV
m_c	Charm quark mass	1.32 GeV
m_s	Strange quark mass	87 MeV
m_t	Top quark mass	173.5 GeV
m_b	Bottom quark mass	4.24 GeV
θ_{12}	CKM 12-mixing angle	13.1°
θ_{23}	CKM 23-mixing angle	2.4°
θ_{13}	CKM 13-mixing angle	0.2°
δ_{CP}	CKM CP violation phase	0.995
g_1 or g'	U(1) gauge coupling	0.357
g_2 or g	SU(2) gauge coupling	0.652
g_3 or g_s	SU(3) gauge coupling	1.221
θ_{QCD}	QCD vacuum angle	≈ 0
v	Higgs vacuum expectation value	246 GeV
m_H	Higgs mass	125 GeV

Table 1.3: Bla

both being gauge theories, has led people to believe a unification is possible. This would unify the forces and particles known from the Standard Model into a *Grand Unified Theory* or GUT. A theory which would add gravity is called a *Theory Of Everything* or TOE.

1.5.1 Running of the coupling constants

The coupling constants in the Standard Model are actually not constants. They depend on the energy of the system. Quantum fluctuations in vacuum have a non negligible contribution in the apparent charge of particles. The effects of screening and anti-screening are visualized in Fig. 1.4 and can be mathematically formulated as a beta function. The function encodes the dependence of a coupling parameter, g , on the energy scale μ :

$$\beta(g) = \frac{\partial g}{\partial \log(\mu)}. \quad (1.31)$$

More coupling in shorter distances (i.e. higher energies) would give rise to a positive beta function and is the case in QED. In QCD gluons carry a color charge and enter the beta function with a negative sign:

$$\beta(g) = - \left(11 - \frac{2n_f}{3} \right) \frac{g^3}{16\pi^2}, \quad (1.32)$$

with n_f the number of fermions that participate in the strong interaction. This decrease in coupling strength in function of energy scale is called *asymptotic freedom*. Alternatively we can write down the coupling constants in function of the energy

$$\alpha_i^{-1}(Q) = \alpha_i^{-1}(m_Z) + \frac{b_i}{2\pi} \log \frac{Q}{m_Z}, \quad (1.33)$$

with $b_1 = -\frac{4}{3}n_g - \frac{1}{2}\log m_Z$, $b_2 = \frac{22}{3} - \frac{4}{3}n_g - \frac{1}{6}$ and $b_3 = \frac{11}{-4}$ the U(1), SU(2) and SU(3) constants in which n_g denotes the number of quark and lepton generations and n_h is the number of Higgs doublet fields. These three couplings seem to converge leading people to believe a more universal theory at higher energies would unify these parameters into one and is shown in Fig. ??.




Figure 1.4: Left: virtual pairs of positrons and electrons can be produced in vacuum. Positrons are attracted to the bare electron resulting into an apparent lower charge of the bare particles from larger distances. This effect is referred to as *screening*. Right: quark and antiquark pairs have a similar screening effect in color charge as the case for electrical charges. There is an anti-screening effect due to gluons which “extract” color from the bare quark and is stronger than the screening effect of quark-antiquark pairs. At small distances the dilution of the initial color charge is larger than at large distance. This drawing should not be taken literally and the concept of self-interaction is very hard to illustrate. One could interpret the initial red color with a gluon emission as a conservation of the red color, thus the red quark can only become blue if a gluon with a red and anti-blue color is emitted (gluons need to be color+anti-color pairs). In the loop two more gluons are present but cannot have a combined color: anti-green with red and green with anti-blue is a possible where green and anti-green have a net zero color charge. Figures from [40].




Figure 1.5: Running of the coupling constants in the Standard Model. Figure from [57].

1.5.2 Unifying theories

Linking the seemingly arbitrary parameters in the Standard Model has been ongoing for the last couple of decades. Linking these theories is not easy since they exhibit very different behaviors. Electromagnetism is long-ranged, the weak force is short-ranged and the strong force is weak in high-energy environments such as the early universe and strong where the probing energy is low. Many GUTs predict that quarks and leptons are part of a single representation of a gauge group with one single hypercharge and would explain why the electric charge of electrons and protons seem to be exactly the same REF.

The simplest GUT is SU(5), which would break down into the Standard Model at lower energies due to spontaneous symmetry breaking. Other possible extensions are SO(10)REF , SU(8)REF and O(16)REF Lie groups.

Without experimental results there is still much ongoing debate into which theory is the correct one. Multiple viable theories still remain. A subset of a handful are given in the following sections.

Supersymmetry

Supersymmetric models impose a new symmetry, supersymmetry (SUSY), that relates fermions and bosons. Every fermion would have a, yet unseen, bosonic supersymmetric partner and vice versa. This theory is highly motivated due to it's ability in cancelling the quadratically divergent terms in the mass correction of the Higgs boson. Unbroken SUSY would lead to partners with the same mass as the particles we know from the SM and would have been discovered long ago. Because of this one assumes the symmetry to be broken.

The last couple of years SUSY was regarded as the most promising extension of the SM with tremendous efforts from big collaborations such as the experiments at the Large Hadron Collidor (LHC) in search for proof. To this date no evidence for SUSY has been found.

Little Higgs

String Theory

1.5.3 How to look for new physics

In general one could say there are two possible ways to look for new physics. Almost all of the physics in our Solar system can be explained with what we know from the Standard Model. Interactions in controlled laboratory environments are currently on the order of 10 TeV in the experiments at the LHC. This could still be well below the energy levels to produce new exotic particles. In the *energy frontier* it is the goal to reach the highest energies possible in order to get as close to the energy requirements where new physics become more prominent. As a consequence more and more cosmic ray experiments have found an interest in searches for physics beyond the Standard Model. This is sometimes called the *cosmic frontier*. This analysis tries to explore this possibility in more detail for the IceCube experiment. Cosmic ray experiments have the disadvantage that they are not fully contained experiments and information is lost as the primary interaction is unknown and important parameters as energy, direction, type,... of the particle have to be reconstructed.

The other approach tries to extract information from precision experiments and are therefor reliant on limiting statistical and systematical uncertainties. In these experiments the intensity of the beam of particle accelerators is pushed to their highest values and is therefor referred to as the *intensity frontier*. This strategy tries to generate huge numbers of particles needed to study rare or exotic subatomic processes. Rare processes could gain us a lot of information on unknown physics. Some parameters which can be calculated in the SM have a small offset in what is measured in experiment. It is possible that new physics enter in Feynman diagrams and have a non-negligible contribution, however rare they are.

Waarom fysica? De vraag van Higgs veld: als er een globaal minimum is kunnen we allemaal dood gaan. Ook zelfde filmpje

I'd like to finish this chapter with quotes from Steven Weinberg in an interview with NovaREF about his vision on string theory. It shows the apparent stalemate phycisist seems to find themselves into: there are no theoretical breakthroughs regarding long standing problems.

"I believe that there is a simple theory that governs everything—the four forces we know about, perhaps other forces as well. I'm not sure that's true. It may be that nature is irreducibly messy. I'm sure that we should assume it's not, because otherwise we're never going to find a fundamental theory. But even so, we're not guaranteed that we'll find it. We may not be smart enough. Dogs are not smart enough to understand quantum mechanics. I'm not sure that people are smart enough to understand the whatever-it-is that unifies everything. I think we probably are, because of our ability to link our minds through language, but I'm not certain.

There was a marvelous period from, I'd say, the mid-'60s until the late '70s when theoretical physicists actually had something to say that experimentalists were interested in. Experimentalists made discoveries that theoretical physicists were interested in. Everything was converging toward a simple picture of the known particles and forces, a picture that eventually became known as the Standard Model. I think I gave it that name. And it was a time when graduate students would run through the halls of a physics building saying they had discovered another particle and it fit the theories, and it was all so exciting.

Since the late '70s, I'd say, particle physics has been in somewhat of a doldrums. Partly it's just the price we're paying for the great success we had in that wonderful time then. I think cosmology now, for example, is much more exciting than particle physics. The string theorists are trying to push ahead without much support from relevant experiments, because there aren't any relevant experiments that can be done at the kind of scales that the string theorists are interested in. "

- Lorentz violation? Nature of Teppei: <https://www.nature.com/articles/d41586-018-05931-2>



2. Theoretical Motivation of the Analysis

As seen in Chapter 1 there is much ongoing debate which beyond the Standard-Model physics models could help explain questions we do not have answers for. Over the last decades this quest has proven to be non-trivial since many anticipated accelerator experiments have not given any clear hints towards physics which cannot be explained by the Standard Model. A big part of the physics community is trying it's best to help answer these riddles and dedicated experiments have been constructed in their search for new physics. Other collaborations try to make use of their detector in the most efficient way possible. These experiments most often try to look for beyond the Standard-Model physics by searching for signals in their detector which could not be explained by the particles we know today. One example, and also being the subject of this work, is to try to look for particles which have a lower electromagnetic non-zero charge than the charged particles of the Standard Model.

2.1 Introduction

As seen in Chapter 1 all free particles have an electromagnetic charge which is a multitude of the absolute electron charge, e , equal to $1.602 \times 10^{-19} C$. Elementary particles such as (anti)quarks have fractional charges equal to $\pm \frac{1}{3}e$ and $\pm \frac{2}{3}e$ but have never been seen as isolated particles due to *confinement* as explained in Section [sub:quarks]. No other particles are expected to have a charge $< e$ and non-zero and are perfect candidates for searches for beyond the Standard Model. Different experiments have sought for these anomalously charged particles and are referred to as *Lightly Ionizing Particles (LIPs)* or *Stable Massive Particles (SMPs)*. Throughout this work the latter denomination is used, indicating they do not rapidly decay and have masses significantly higher than the lightest leptons.

2.2 Theory

In Section 1.5.2 I have introduced possible extensions of the Standard Model. One of the simplest possible extensions of the $SU(3) \times SU(2) \times U(1)$ group is the $SU(5)$ gauge group. It is the smallest Lie group that can contain the group of the Standard Model without introducing any new fermions. It could explain charge quantization REF, has complex representations and can accommodate fractional charges. In this scheme new vector bosons, usually called X and Y bosons, occur with charges $\frac{4}{3}$ and $\frac{1}{3}$. Extensions of the $SU(5)$ models allow for color singlet particles with charges $\frac{1}{3}$ and $\frac{2}{3}$ [23]. Other possible extentions are the $SU(7)$ [44], $SU(8)$ [70],

$\text{SO}(14)$ [69], $\text{SO}(18)$ [41], $\text{SO}(10) \times \text{SO}(8)$ [49].

It should be noted that the simplest form of an $\text{SU}(5)$ gauge group is already highly constrained as proton decay is allowed but experimental results have shown the lifetime to be $> 1.67 \times 10^{34}$ years ($\tau(p \rightarrow \pi^0 e^+)$) and $> 6.6 \times 10^{33}$ years.

There are also some string theories massive particles with a fractional charge are also predicted [18, 68], which was later confirmed to occur very often in certain compactifications [19].

More recently there has been an increasing interest in searches for millicharged particles. New particles could couple to the Standard Model via a “kinetic mixing” or “hypercharge portal” [46, 48]. And in recent years, they were studied as possible candidates for dark matter [26, 27, 29, 61]. The charges of these particles are however often $< 10^{-3}e$ and no ideal candidates in neutrino Cherenkov experiments. It is possible to look for them in neutrino experiments [52] but are more targeted toward future experiments such as DUNE [10] and SHiP [17]. A more detailed explanation of these particles can be found in [24]

There are possibly many other possible extensions but go beyond the scope of this work. One could just keep in mind that no free particles with an anomalous charge less than e are expected, and if seen would give clear hints of beyond the Standard-Model physics and would help in finding a more clear picture of what is possibly lurking beyond the realms of our understanding.

https://ac.els-cdn.com/S0370269314001257/1-s2.0-S0370269314001257-main.pdf?_tid=7f8b3af4-8845-417fa219-703f382cb092&acdnat=1535556263_162a6333d0a615412be21aa5fc3e5720

2.3 Properties of the signal

Because there are many possible scenarios in what these particles are, originate from or are produced one has to make certain assumptions about the properties of the signal. A particle traveling at the speed of light with a lifetime < 0.1 seconds traversing a detector will not give the same signal properties as one that has a very long lifetime. Therefore, I have chosen that the particles I am looking for

- behave leptonically, similar to muons,
- have a long lifetime and will not decay within the detector, or have a very low probability,
- follow an energy spectrum with a spectrum of -2^* ,
- are assumed to produce a uniform flux in angle space close to the detector.

These assumptions are consistent with previous searches which are mentioned in Section 2.4. The behaviour of these particles in the detector will depend on the charge (see section???) and, to a lesser extent, the mass. In this work I have chosen to look for particles with a

- charge of $1/3$, $1/2$ and $2/3$,
- mass of 10 GeV, 100 GeV, 1 TeV, 10 TeV and 100 TeV,

where I have referred to the charge of the particles as relative to the absolute electron charge, e^\dagger . The possible combinations result into a total of 15 unique signal samples which will be searched for.

2.4 Previous searches

There are several ways on can assume to produce fractional charge particles. Different assumptions lead to different possible searches with previous and current detectors. In the following I will give the results of several experiments. More information and a very good overview can be found in

2.4.1 Searches with accelerators and fixed targets

The total energy of the interaction should be large enough to produce particles of a certain mass. The square of the centre of mass energy is given by:

*More information about spectra can be found in section???????

[†]This will be done throughout this work from this point on.

Energy (GeV)	Charges sought	Collider	Reference
1-1.4	$\frac{2}{3}$	VEPP-2M	[28]
29	$\frac{1}{3}, \frac{2}{3}$	PEP	[12]
130-209	$\frac{2}{3}, \frac{4}{3}, \frac{5}{3}$	LEP	[3]
130-136, 161 and 172	$\frac{2}{3}$	LEP	[9]
91.2 (m_Z)	$\frac{2}{3}, \frac{4}{3}$	LEP	[13]
91.2 (m_Z)	$\frac{4}{3}$	LEP	[30]

Table 2.1: Highest-energy fractional charge particle searches in electron-positron colliders. No evidence for fractionally charged particles was found. From ??.

$$\begin{aligned} s &= (p_1 c + p_2 c)^2 \\ &= m_1^2 c^4 + m_2^2 c^4 + 2E_1 E_2 - 2\vec{p}_1 \cdot \vec{p}_2 c^2, \end{aligned} \quad (2.1)$$

where $p_{1,2}$ are the four-momenta of the two particles and c is the speed of light. Assuming E is the energy of the incoming particle and m the mass of a target particle in rest, the maximum mass reach of a search is given by:

$$m_{max} \approx \sqrt{2mE}. \quad (2.2)$$

If I is the incoming particle from the input beam and N a nucleus then the production of exotic particles can be depicted as

$$I + N \rightarrow F + X, \quad (2.3)$$

where F stands for the fractional charged particle and X for the other particles which are produced in the interaction. No experiments that used accelerators and fixed targets found evidence for the existence of fractional charge particles [51]. The highest-energy search used muons with a muon beam of 200 GeV, resulting in an m_{max} of 19 GeV/c² [21].

2.4.2 Colliders

Particle colliders can reach much higher energies than most fixed-target experiments. The maximal mass of new particles in a storage ring which is colliding particles of energy E , Eq. 2.1 gives

$$s = 4E^2. \quad (2.4)$$

There is a big difference in lepton and hadron accelerator experiments as much less particles are being produced in the former due to the absence of strong interactions. The production is “cleaner” and the sought particles are easier to distinguish from other productions. But, it is more difficult to reach higher energies* for lepton accelerators. An overview of electron-positron colliders is given in Tab. 2.4.2. No evidence for fractionally charged particles was found.

Experiments that use proton-antiproton colliders have reached large masses but have also found no evidence of fractionally charged particles. An overview is given in Tab. 2.4.2.

A more recent search was performed at the LHC, a proton-proton collider, when operating at an energy of 7 TeV. No evidence of particles with fractional charge was found. An upper limit of 95% confidence level was set for particles with electric charge $\frac{2}{3}$ up to a mass of 310 GeV and 140 GeV for those with charge $\frac{1}{3}$ [31].

*The radiative power of synchrotron radiation scales with a factor of m^{-4} : particles with low mass lose much more energy in circular accelerators with a fixed radius.

Energy (TeV)	Charges sought	Collider	Reference
0.54	$\frac{1}{2}, \frac{2}{3}$	SPS	[22]
1.8	$\frac{3}{2}, \frac{4}{3}$	Tevatron	[4]
1.8	$\frac{1}{3}, \frac{2}{3}$	Tevatron	[11]

Table 2.2: Highest-energy fractional charge particle searches in proton-antiproton colliders. No evidence for fractionally charged particles was found. From ??




Figure 2.1: Upper limits on fluxed of particles close to the repective detectors. LIP stands for *Lightly Ionizing Particles*. From [16]

2.4.3 Searches for particles with telescopes

There are several ways particles with a fractional charge could be produced in cosmological events;


- The particles were produced early on in the Universe and are a stable component of the present material;
- The particles are rare but can be continuously produced in high-energetic astrophysical event; or
- The particles are produced in cosmic ray processes on Earth.

Because there is no clear preference in one of these possibilities most telescope experiments therefore express their search sensitivity in function of an incoming flux close to the detector in units of $\text{cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$. This analysis has adopted the same search strategy and aims to improve upon previous results. The most stringent upper limit was realized by the MACRO experiment found on the arXive which compares results from older searches and can be found in Fig. 2.4.3. The best published result is set by Kamiokande II [w]ith upper limits of $2.1 \times 10^{-15} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ and $2.3 \times 10^{-15} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ for particles with charges $\frac{1}{3}$ and $\frac{2}{3}$ respectively [55].

Hmmm... Deze heb je nog nooit bekeken: <https://arxiv.org/pdf/1601.04004.pdf> +references!

Particle physicists don't look for a needle in a haystack as what is sometimes used for an

analogy to make clear in what we are doing. We are in fact looking at hay in a haystack. The hay we are looking for has slightly different properties. It can be a bit drier or a bit longer, but it takes an enormous amount of time and clever thinking to be able to distinguish the hay from normal hay. But sometimes errors can slip in, or you can have a very long normal piece of hay while you could have smaller pieces of new hay so finding a new one doesn't give you a hundred percent assurance that you've indeed found something new. You need to find enough of the new ones to be able to say that there are too many found that could be obtained by pure chance from the normal set.



3. Cosmic Rays and Neutrinos

Cosmic rays almost exclusively refer to particles with a finite rest mass. The term *rays* was historically wrongly attributed to these particles as they were thought to be mostly electromagnetic radiation. The interest of cosmic rays within the field of particle physics and modern particle physics is clear: multiple new particles were discovered from the interactions at energies which were higher than most experiments could reach. Positrons, muons, pions and kaons were first discovered in cosmic ray experiments in the 1930s and 40s. Today, high-energy cosmic ray interactions are still of interest as the highest energies of these particles go beyond what is feasible at the most powerful accelerators such as the LHC. Neutrinos are expected to be produced together with cosmic rays near the source or close to Earth making neutrino astronomy a powerful and important part of modern day astronomy. In this chapter I will give BLA BLA BLA. For a more exhaustive description of cosmic rays I refer the reader to [Gaisser:2016uoy].

3.1 Discovery of cosmic rays

With the use of electrometers, Victor Hess performed multiple ground-breaking balloon flight experiments in 1912 to prove that the amount of radiation increases with altitude [63]. This was in strong contradiction with the widespread belief that radiation on Earth's surface mostly originates from radioactive substances in its crust. Hess concluded that an extremely penetrating radiation existed. He described this radiation to be coming from space which then enters Earth's atmosphere which proved to be correct but it was wrongfully attributed to electromagnetic radiation by Robert Millikan in the 1920s [PhysRev.32.533].

Hess later ruled out the possibility that cosmic rays originate from the Sun as his observations showed no particular differences in night and day and during solar eclipses. In the late 1920s, first evidence was found that cosmic rays were charged due to a variation of their intensity with latitude [33]. This indicated that they were deflected by the geomagnetic field.

3.2 What are cosmic rays?

Cosmic rays are, almost exclusively, the collection of nuclei which are stripped of their electrons, making them electrically charged, heavy particles. Around 90% of the particles are ionized hydrogen atoms, or protons. 9% are alpha particles and 1% are nuclei of heavier elements. There is a striking resemblance between the relative abundance of cosmic rays and elements in the Solar System as seen in Fig. 3.2. A much smaller fraction of incoming particles are electrons,




Figure 3.1: The cosmic ray elemental abundances measured on Earth compared to the solar system abundances, all relative to carbon = 100. Figure from ACE news archive [ISRAEL2005201].

positron and antiprotons.

There are however two important differences between cosmic rays and elements from our Solar System. Firstly, the two groups of elements Li, Be, B and Sc, Ti, V, Cr, Mn are many orders of magnitude more abundant in cosmic rays than in the solar system. This is due to their absence in stellar nucleosynthesis and are therefore not expected to be produced in large numbers. More massive cosmic rays (mainly C, O and Fe) can produce these nuclei in the process of *spallation*. They are produced by collisions of cosmic rays with the interstellar medium. Therefore, these nuclei are sometimes referred to as *secondary nuclei*. The second difference is that nuclei with $Z > 1$ are much more abundant with respect to hydrogen for cosmic rays. This phenomenon is not yet well understood but could be attributed to the difficulty to ionize hydrogen, necessary for acceleration processes.

The amount of cosmic rays seen on Earth is expressed in units of $[m^{-2}s^{-1}sr^{-1}]$. We can see in Fig. 3.2 that the cosmic ray flux follows a energy power law spectrum:

$$dN/dE \propto E^{-\gamma} dE, \quad (3.1)$$

where γ is called the *spectrum index*. Because of the steepness of the spectrum it is often multiplied by a higher power of energy*.

We can divide the global spectrum in four regions. Between 10 GeV and 1 PeV the differential spectrum index is around -2.7. From 10 PeV to 1 EeV it is around -3.1. Above 10 EeV the spectrum again flattens to an index around -2.6 and an apparent cutoff region is present at around 10^{20} eV. The transition of this first to second region is referred to as the *knee* at around 3 PeV. The second to third region transition is referred to as the *ankle*. It is possible to describe the full cosmic-ray spectrum with sources within our galaxy. A more generally accepted theory is that the knee in the spectrum originates from the end of a population of particles which are accelerated within our Milky Way [Gaisser:2013bla].

Second knee?

*The broad range in both energy and flux should convince the reader that many types of detectors are necessary to study the behaviour of cosmic rays. Low-energy particles are abundant and high-energy particles are much more rare. Both the energy and the incoming flux will determine the type and size of the detector.




Figure 3.2: Spectrum of cosmic rays at Earth. The all-particle spectrum measured by different experiments is plotted together with the proton-only spectrum. Subdominant contributions to the total flux from electrons, positrons and antiprotons as measured by the PAMELA experiment are also shown. Figure from [Blasi:2013rva].

The origin of cosmic rays has been a topic of discussion for many years. We know now that most particles originate from sources in the local galaxy, having spent on average 10^7 years in diffusive motion in the interstellar medium [Gaisser:2013bla]. This is consistent with the resemblance of the relative abundances of cosmic rays and elements from our Solar System. However, there is no general consensus about the origin of the cosmic rays with energies above 3×10^{18} eV. In the following, the abovementioned energy regions are discussed in more detail.

3.2.1 Solar modulation

In the Solar System a stream of charged particles is released from the Sun. This stream is mostly made up of electrons, protons and alpha particles with kinetic energies ranging between 0.5 and 10 keV. Within this solar wind plasma there is a magnetic field. Cosmic rays coming in to the Solar System interact with these particles and magnetic field. The influence is greatest on particles with the lowest charges. This effect is called *solar modulation*. In effect, we see a strong suppression of cosmic rays at energies of 10 GeV and below.

3.2.2 Galactic component

The most probable acceleration mechanism for cosmic rays originating from our Galaxy is by shocks driven by expanding supernova remnants [0034-4885-64-4-201]. From the ratio of primary to secondary nuclei it can be inferred that cosmic rays travel distances thousands of times greater than the thickness of the disk of the Galaxy. There is also an apparent decrease in the amount of matter that is traversed by cosmic rays with higher energies than with lower. Higher-energy cosmic rays seem to spend less time in the Galaxy than lower-energy ones and suggests that cosmic rays are accelerated before most propagation occurs [Gaisser:2016uoy].

The way the spectrum is fit is not set in stone. Here I will use the convention used by Gaisser, Stanev and Tilav described used in reference [Gaisser:2013bla]. The spectrum is subdivided in three populations. The first population corresponds to the particles accelerated by supernova remnants, with the knee signaling the cutoff of this population. The second population is a higher-energy galactic component of unknown origin. The third generation will be described in more detail in ???. Assuming that the primary spectrum depends on the *magnetic rigidity*^{*},

$$R = \frac{pc}{Ze}, \quad (3.2)$$

where Ze is the charge of a nucleus of total energy $E_{tot} = pc$ and relates to the gyroradius of a particle in a given magnetic field B as

$$r_L = \frac{R}{B}. \quad (3.3)$$

If there is a characteristic regidity, R_e above which a particular acceleration process reaches a limit, then the feature will show up in total energy first for protons, then for helium and so forth for heavier nuclei according to

$$E_{tot} = Ze \times R_e. \quad (3.4)$$

This effect is visualised in Fig. 3.2.2 and indicates that as one population reaches its maximum the composition becomes heavier. The second knee, reported by KASCADE-Grande [Apel:2011mi] and GAMMA [Garyaka:2008gs] could be explained with an “iron knee” bump.

3.2.3 Extragalactic component

The flux at the highest energies is exceedingly small. The number of events per year at energies above 5×10^{19} eV is around one per square kilometer per century. There are only two experiments in the world capable of detecting the highest-energy cosmic rays in a statistical meaningful way:

*An assumption which is experimentally favored over other assumptions. Rigidity is an appropriate variable for interpreting changes in the spectrum due to propagation and acceleration in magnetic fields.




Figure 3.3: Blub

Telescope Array, located in the Northern Hemisphere (area of $\approx 700 \text{ km}^2$) and the Pierre Auger Observatory in the Southern Hemisphere (area of $\approx 3000 \text{ km}^2$).

Both experiments see a suppression of the flux above $6 \times 10^{19} \text{ eV}$. The exponential cut-off is consistent with the expected Greisen-Zatsepin-Kuzmin (GZK) effect [[Greisen:1966jv](#), [Zatsepin:1966jv](#)] where cosmic rays interact with the cosmic microwave background radiation (CMB)

$$\gamma_{CMB} + p \rightarrow \Delta^+ \rightarrow p + \pi^0 \quad (3.5)$$

or

$$\gamma_{CMB} + p \rightarrow \Delta^+ \rightarrow n + \pi^+. \quad (3.6)$$

Particles with energies above $5 \times 10^{19} \text{ eV}$ would interact with the CMB, leading to an exponential cutoff (but if the incoming particles above these energies would be relatively young it is still possible for them to reach the detector). The Pierre Auger experiment reported to see higher compositions at the highest energies [[icrc2017:pa](#)]. If the particle is a nucleus with A nucleons, then the GZK limit applies to its nucleons, which carry only a fraction $1/A$ of the total energy. For iron nuclei this would for example result in a limit of $2.8 \times 10^{21} \text{ eV}$. In contrast, the TA experiment interpreted their data as implying a light primary composition (mainly p and He) at the highest energies. Both experiments use a different interpretations for crucial quantities of these measurements and a thorough joint analysis conducted by both experiments states that, at the current level of statistics and understanding of systematics, both data sets are compatible with being drawn from the same parent distribution [59].

The Pierre Auger Observatory also reported evidence in an anisotropic distribution of the arrival directions of the highest-energy cosmic rays [[Aab:2017tyv](#)] in a direction where the distribution of galaxies is relatively high and does not coincide with the galactic plane. These observations, together with our lack of known possible sources within our galaxy for these ultra-high energies shows compelling evidence that these particles have an origin from outside our galaxy. Dan misschien ook verwijzen naar paper van Waxman dat de hoogst energetische neutrinos en CRs linkt en zegt dat ze inconsistent zijn met galactische bronnen (2013)

"anti-matter but no anti-nuclei???" Ook nog onduidelijkheden en waarom compositie zo moeilijk is. Compositie ook afhankelijk van energieschaal waar je naar kijkt bv.

Copy paste van Wikipedia: Data from the Fermi Space Telescope (2013)[3] have been interpreted as evidence that a significant fraction of primary cosmic rays originate from the




Figure 3.4: Expanded view of the highest energy portion of the cosmic-ray spectrum from data of the Telescope Array and the Pierre Auger Observatory [pdg2018].

supernova explosions of stars.[4] Active galactic nuclei also appear to produce cosmic rays, based on observations of a neutrino and gamma rays from blazar TXS 0506+056 in 2018.[5][6]

extragalactic neutrinos: IC paper

To put it simply, understanding cosmic rays and where they originate can help us answer fundamental questions about the origins of the universe, our galaxy and ourselves.

3.3 Acceleration mechanism

How cosmic rays got their signature slope in the energy spectrum and its intricate details have been under discussion for multiple decades. To this date there is no clear picture how these particles are accelerated in full detail. It is beyond the scope of this work to give a comprehensive overview of all possible acceleration mechanisms or possible sources. Most calculations are left out and for a more detailed discussion the reader is referred to specialized books or the references in the text.

The acceleration of the particles can be subdivided into two questions. First, where are the particles accelerated? Does it happen on large scales, cosmological distances in galaxies or near specific sources? Secondly, how are these particles exactly accelerated? What is the driving mechanism? Since primary cosmic rays are all electromagnetically charged particles these mechanisms should clearly be sought for in places where electric and/or magnetic fields play a dominant role.

Galactic accelerators

With their approximate energy density around 0.5 eV/cm^3 in our local galaxy, the bulk of cosmic ray acceleration could very well be explained by **supernovae**. This density results into a total power of around

$$L_{CR} = 7 \times 10^{40} \text{ erg/s}, \quad (3.7)$$

where erg is a unit often used in astronomy*. If one assumes a supernova explosion of around one per every 30 years then the total power output of type II supernovae with a mass output of around ten times the mass of the Sun at a velocity close to $5 \times 10^8 \text{ cm/s}$ would result in a power of

$$L_{SN} \sim 3 \times 10^{42} \text{ erg/s}. \quad (3.8)$$

* $1 \text{ erg} = 10^{-7} \text{ J}$.




Figure 3.5: The Hillas plot of potential cosmic ray accelerators locates objects according to size and magnetic field. Objects to the left of the diagonal lines cannot accelerate particles to 10^{20} eV (proton: solid, iron: broken). Image obtained from [Bauleo:2009zz]

These numbers are not set in stone and hold large uncertainties, but it shows that with an acceleration efficiency on the order of a couple of percent supernova explosions are a prominent source of energetic cosmic rays, if not the dominant one.

Extragalactic accelerators

We will see in Section 3.3.1 that the maximum energy from shock acceleration by a supernova remnant is insufficient to explain UHECR. As explained in Section 3.3.1, particles can be accelerated if the trajectory of the particles can be changed and energy can be transferred multiple times. The magnetic fields responsible for the course change of these particles has to be sufficient in magnitude in order for these particles not to escape and go beyond the grasp of the source responsible for the acceleration. This limitation is expressed by the gyroradius in the accelerator, $r_L = E/ZeB$ similar to Eq. 3.3, requiring it to be smaller than the radius of the accelerator: $r_L < R$ or $E < ZeBR$.

Even if only qualitative, this relation provides an interesting criterion to identify possible sources of UHECRs by looking at the accelerator related term BR . This was done in a classic paper by Hillas [Hillas:1985is], illustrated in the more recent Fig. 3.3. Accelerators necessary to explain the amount of UHECRs are not populated (enough) in our galaxy, making them more likely to be of extra-galactic origin. **Active galactic nuclei, blazars and gamma ray bursts** (en andere als je die hebt toegevoegd later) are therefore also briefly explained.

3.3.1 Supernova (remnants)

Supernovae can be broadly subdivided into two categories: type I and type II. Type I supernova explosions happen in binary star systems. In those systems one of the two stars is a carbon-oxygen white dwarf which accretes matter from the second star. When the total mass of the white dwarf reaches the Chandrasekhar REF limit of around 1.44 solar masses it cannot longer hold itself under the gravitational pressure and collapses in on itself. Within seconds, the carbon




Figure 3.6: Left: the Crab Nebula is the supernova remnant approximately one thousand years old. The supernova was noted by Chinese astronomers in the year 1054 AD. Right: Chandra X-ray observatory picture of the Cassiopeia A supernova remnant (pictures from NASA).

component in the white dwarf starts nuclear fusion and enough energy is released to produce an explosion brighter than the Sun with a factor of around 5 billion. A resulting shock wave can reach up to around 3% the speed of light.

Type II supernova explosions differ by being single star systems. When a star reaches the end of its lifecycle the subsequent fusion reactions reach to a halt. If the star has enough mass (at least 8 times the mass of the Sun), it is possible for the inner core to again reach the Chandrasekhar limit and collapse in on itself due to the lack of *electron degeneracy*. Without the outward pressure of nuclear fusion reactions and the support of the core, the outer layers of the star collapse under the gravitational pressure. The compression of the electrons and protons into neutrons results into a very hot, dense, neutron core. The velocity of the inwards falling layers can reach to a staggering 23% of the speed of light and recoil when hitting the remaining core. Neutrinos are produced in this violent core collapse and the outward going shockwave hits the remaining outer layers forming the supernova explosion *.

Because of their brightness supernovae within our galaxy can be seen with the naked eye (provided they are not too far away). The last recorded supernova from our galaxy was by Johannes Kepler in 1604 but earliest recordings go back to 185 AD by Chinese astronomers[†]. The question remains how supernovae can serve as cosmic ray accelerators. In 1949, Enrico Fermi proposed a mechanism where particles can gain energy by collisions with moving interstellar ionized gas clouds. Only later, it was realized that a large, plane shock front moving with a certain velocity is able to accelerate charged particles much more efficiently. This first mechanism results into an energy transfer proportional to the squared velocity of the cloud and is thus called *second order Fermi acceleration*. Shock front acceleration energy transfer is proportional to the velocity and is called *first order Fermi acceleration*. Supernova remnants provide an explanation for the origin of these shock fronts.

* Astronomer Carl Sagan once said, "The nitrogen in our DNA, the calcium in our teeth, the iron in our blood, the carbon in our apple pies were made in the interiors of collapsing stars. We are made of starstuff."

[†]From observations of other galaxies supernovae are expected to occur, on average, once every thirty years. Not all of these will be visible to the naked eye, but would almost certainly be observable with modern astronomical telescopes.




Figure 3.7: Left: magnetic cloud. Right: shock waves typically have magnetic inhomogeneities both preceding (downstream) and following them (upstream). If a charged particles travels through the shock wave it can gain velocity through first-order Fermi acceleration. In the illustration a particle travels from upstream to downstream and back upstream. At every back and forth movement the particle effectively gains in energy. For a particle with a velocity u_1 relative to the shock front the front seems to come at him with velocity $-u_1$. The downstream medium has a velocity relative to the shock front of $u_2 < u_1$ making it seem coming towards the particle with velocity $u_1 - u_2$.

First- and second-order Fermi acceleration

Suppose we have a magnetic cloud in the interstellar medium travelling with a certain velocity \vec{V} and a particle with velocity \vec{v} enters the cloud under an angle θ_1 (see Fig. 3.3.1). If we assume collisionless scattering can occur (no energy is dissipated from the particle to the cloud) due to the magnetic fields in the cloud, the magnitude of the momentum in the rest frame of the cloud will not change ($E'_1 = E'_2$, where we the apostrophe denotes the cloud rest frame). From special relativity we know that:

$$\begin{aligned} E'_1 &= \gamma (E_1 - p_{1,\parallel} V) \\ &= \gamma E_1 (1 - \beta \cos \theta_1), \end{aligned} \tag{3.9}$$

with $\beta = V/c$ and γ the Lorentz factor. Similarly and using $E'_1 = E'_2$

$$\begin{aligned} E_2 &= \gamma E'_2 (1 + \beta \cos \theta'_2) \\ &= \gamma^2 E_1 (1 - \beta \cos \theta_1) (1 + \beta \cos \theta'_2) \end{aligned} \tag{3.10}$$

and

$$\frac{\Delta E}{E} = \frac{E_2 - E_1}{E_1} = \frac{1 - \beta \cos \theta_1 + \beta \cos \theta'_2 - \beta^2 \cos \theta_1 \cos \theta'_2}{1 - \beta^2} - 1. \tag{3.11}$$

By hypothesis, the escaping particles are isotropic in the cloud frame: $\langle \cos \theta'_2 \rangle = 0$. One can show that $\langle \cos \theta_1 \rangle = -\frac{\beta}{3}$ [Gaisser:2016uoy], leading to

$$\frac{\Delta E}{E} = \frac{4}{3} \frac{\beta^2}{1 - \beta^2} \approx \frac{4}{3} \beta^2, \tag{3.12}$$

showing that for molecular clouds the energy gain is indeed proportional to the square of β for second-order Fermi acceleration.

If a particle is incoming to an expanding shock (see Fig. 3.3.1) $\langle \cos \theta'_2 \rangle$ is equal to 2/3, leading to

$$\frac{\Delta E}{E} = \frac{\frac{4}{3}\beta + \frac{13}{9}\beta^2}{1 - \beta^2} \approx \frac{4}{3}\beta, \quad (3.13)$$

where β is now equal to $u_1 - u_2$ as explained in the caption of the figure. We have shown that for shock fronts the energy gain is indeed proportional to β for first-order Fermi acceleration. From both the outcome as the discussion it is clear that the energy gain enters through relativistic effects, making the intuitive approach not straightforward.

Power

The energy gain of a “single collision” results into the power law spectrum when considering a process in which a test particle increases its energy by an amount proportional to its energy with each encounter. Let us assume $\Delta E = \xi E$, after n encounters:

$$E_n = E_0 (1 + \xi)^n, \quad (3.14)$$

where E_0 is the energy when the particle first enters the accelerator medium. To reach a certain energy E' , the particles must encounter a number of collisions

$$n(E') = \frac{\ln\left(\frac{E'}{E_0}\right)}{\ln(1 + \xi)}. \quad (3.15)$$

To reach energies of E' or higher, the number of collisions will be proportional to

$$\begin{aligned} N(\geq E') &\propto \sum_{m=n}^{\infty} P_{present}(m) = \sum_{m=n}^{\infty} (1 - P_{esc})^m \\ &= (1 - P_{esc})^n \left((1 - P_{esc}) + (1 - P_{esc})^2 + \dots \right) \\ &= \frac{(1 - P_{esc})^n}{P_{esc}}, \end{aligned} \quad (3.16)$$

where $P_{present}$ is the probability of a particle still being present in the accelerator and P_{esc} the probability of the particle to escape after per collision. Making use of $a^{\ln b} = e^{\ln a \ln b} = b^{\ln a}$ and inserting Eq. B.4

$$N(\geq E') \propto \frac{1}{P_{esc}} \left(\frac{E'}{E_0} \right)^{-\gamma}, \quad (3.17)$$

with

$$\gamma = \frac{\ln\left(\frac{1}{1-P_{esc}}\right)}{\ln(1 + \xi)} \approx \frac{P_{esc}}{\xi}. \quad (3.18)$$

The power law spectrum becomes visible in the derivative of the number of particles in energy

$$\frac{dN}{dE} \sim E^{-(\gamma+1)}, \quad (3.19)$$

in agreement with Eq. 3.1. Shock wave fronts have an expected $\gamma \approx 1$ giving rise to a different spectrum to what is seen on Earth but which can be explained by propagation from the source to Earth (this is beyond the scope of this work).

Seems plausible all galactic CRs are accounted for by supernovae. This is supported with the realization that first-order Fermi acceleration naturally produces a spectrum of cosmic rays close to what is observed.

Maximum energy

The highest energies that particle can be accelerated to can be defined by:

- The differential energy gain per collision dE/dt
- The total time the particle can be accelerated
- . The energy gain is given by

$$\frac{dE}{dt} = \frac{\xi E}{T_{cycle}}, \quad (3.20)$$

where T_{cycle} is the characteristic time for one acceleration cycle. T_{cycle} depends on the diffusion coefficients and velocities of the upstream and downstream regions and is set to $T_{cycle} \geq 20E/(3u_1ZeB)$ by Lagage and Cesarsky [Lagage:1983zz] for a strong shock and arguing that the diffusion length, λ_D cannot be smaller than the Larmor radius of the particle. Particles with a Larmor radius greater than the irregularities holding a magnetic field are not prone to be heavily influenced by them. Lagage and Cesarsky therefore concluded that

$$E_{max} \leq \frac{3}{20} \frac{u_1}{c} ZeB(u_1 T_{ST}), \quad (3.21)$$

where T_{ST} is the Sedov-Taylor time where particles are less prone to escape and is ~ 1000 years. For $u_1 \sim 10^9$ cm/s [stanev2010high] and $B \sim 3\mu G$ the Lagage and Cesarsky limit reads

$$E_{max} \leq Z \times 2.4 \times 10^5 GeV. \quad (3.22)$$

3.3.2 Active Galactic Nuclei

3.3.3 Blazars

verwijzing naar txs!

3.3.4 Gamma Ray Bursts

3.4 Neutrinos

image with all kinds of neutrinos? Van Gary Binder Ook iets zeggen over dat we ook extreem veel gamma ray studies hebben? Pg 236-237 in Gaisser. Ook waarom neutrinos zoveel handiger zijn dan fotonen en CRs

3.5 Air shower

3.5.1 Cosmological

3.5.2 Solar

3.5.3 Atmospheric

Conventional
Prompt

3.5.4 Astrophysical

Gaisser fig 2.10.


3.6 Cosmic ray and neutrino detectors

Heel kort: pg 5 Gaisser boek. Zo IC introduceren.

Ook dat plotje waarbij je toont dat neutrinos interessanter zijn in hoge E omdat fotonen onzichtbaar worden!

The IceCube Collaboration instead tested the principle using neutrinos. Neutrinos interact with matter through the weak force — one of the four fundamental forces of nature. The influence of the weak force is limited to minute distances. As a result, interactions between neutrinos and matter are extremely improbable, and a neutrino can easily traverse the entire Earth unimpeded. This poses a challenge for physicists trying to study these elusive particles, because almost every neutrino will simply pass through any detector completely unnoticed.

In neutral-current reactions neutrinos lose energy, but are not absorbed
hier tot pg 50?



4. Relativistic Particles in Matter

hier tot pg 60?



5. The IceCube Detector

hier tot pg 75?

Simulation, Data Processing and Analysis

6	Event Generation, Simulation and Reconstruction	53
6.1	Line-Fit	
6.2	SPE and MPE	
6.3	MILLIPEDE	
6.4	Boosted Decision Tree Classifiers	
7	The SPACE Analysis	55
7.1	Motivation	
7.2	General strategy	
7.3	Event cleaning	
7.4	Variables	
7.5	BDT results	
7.6	Pull validation	
7.7	Systematic Uncertainties	
7.8	Results	
8	Summary and Discussion	57
	Appendices	59
A	Gauge symmetries	61
B	Planck's law	63
B.1	Electromagnetic waves in a cubical cavity	
9	Some useful things for LaTeX	65
9.1	Definitions	
9.2	Remarks	
9.3	Corollaries	
9.4	Propositions	
9.5	Examples	
9.6	Exercises	
9.7	Problems	
9.8	Vocabulary	
	Bibliography	67
	Articles	
	Books	
	Index	71



6. Event Generation, Simulation and Reconstruction


- 6.1 Line-Fit
- 6.2 SPE and MPE
- 6.3 Millipede
- 6.4 Boosted Decision Tree Classifiers



7. The SPACE Analysis


<https://arxiv.org/pdf/physics/0312102v1.pdf>
<https://arxiv.org/pdf/1310.1284.pdf>

- 7.1 Motivation**
- 7.2 General strategy**
- 7.3 Event cleaning**
- 7.4 Variables**
- 7.5 BDT results**
- 7.6 Pull validation**
- 7.7 Systematic Uncertainties**
- 7.8 Results**



8. Summary and Discussion

Appendices




A. Gauge symmetries

The difference between global and local symmetries are not straightforward for everybody. In this appendix I try to give a better view of the matter.

Imagine that at each point in space and time there is a circle attached to it. If one shifts all circles of all points with a fixed angle the underlying physics hasn't changed. If we look at the whole in a different angle, nothing seems to be changed as everything holds the same relative orientation. This is a global symmetry. For local symmetries we instead shift each circle through a different angle, but an angle that changes smoothly from point to point and in a way that we can say how that angle is varying between different nearby regions. Then it will turn out that we can describe that rotation angle by means of a so-called gauge field, which just lets us transport the charged scalar field from one point in space time to another, taking account of how the rotation angle of the circle is changing. A gauge is a kind of coordinate system that is varying depending on the location with respect to some underlying space. In physics we are almost always concerned with space-time as the underlying space, and we are typically interested in theories that are invariant with respect to the choice of gauge or coordinate system.

Dan wat uitleg vanuit je QFT boek en de dingen hieronder: Je wilt je derivative anders doen werken in je theory onder een transformatie, maar daarvoor heb je een veld nodig. M.a.w.: dankzij een veld heb je lokale ijktransformatie mogelijk!



B. Planck's law

bron: <http://hyperphysics.phy-astr.gsu.edu/hbase/quantum/rayj.html>

B.1 Electromagnetic waves in a cubical cavity

Suppose we have EM waves in a cavity at equilibrium with its surroundings. These waves must satisfy the wave equation in three dimensions:

$$\frac{\partial^2 \Psi}{\partial x^2} + \frac{\partial^2 \Psi}{\partial y^2} + \frac{\partial^2 \Psi}{\partial z^2} = \frac{1}{c^2} \frac{\partial^2 \Psi}{\partial t^2}. \quad (\text{B.1})$$

The solution must give zero amplitude at the walls. A non-zero value would mean energy is dissipated through the walls which is in contradiction to our equilibrium assumption. A general solution takes the form of

$$\Psi(x, y, z, t) = \Psi_0 \sin k_1 x \sin k_2 y \sin k_3 z \sin k_4 t, \quad (\text{B.2})$$

which, after requiring $k_n L = n\pi$ with $n = 0, 1, 2, \dots$ and $k_4 \frac{\lambda}{2c} = \pi$, leads to

$$\Psi(x, y, z, t) = \Psi_0 \sin \left(\frac{n_1 \pi x}{L} \right) \sin \left(\frac{n_2 \pi y}{L} \right) \sin \left(\frac{n_3 \pi z}{L} \right) \sin \left(\frac{2\pi c t}{\lambda} \right). \quad (\text{B.3})$$

From the wave equation it is easy to find that

$$n^2 = n_1^2 + n_2^2 + n_3^2 = \frac{4L^2}{\lambda^2}, \quad (\text{B.4})$$

which span up a sphere in “n-space” with a volume of $\frac{1}{8} \frac{4}{3} \pi n^{3/2}$, where the first term originates from the positive nature of $n_{1,2,3}$. Because there are two possible polarizations of the waves one has to multiply with an additional factor 2. The number of modes per unit wavelength is equal to

$$\frac{dN}{d\lambda} \times \frac{1}{L^3} = \frac{d}{d\lambda} \left[\frac{8\pi L^3}{3\lambda^3} \right] \times \frac{1}{L^3} = - \left[\frac{8\pi}{\lambda^4} \right]. \quad (\text{B.5})$$

B.1.1 Classical approach

Following the principle of equipartition of energy, each standing wave mode will have an average energy kT with k the Boltzmann constant and T the temperature in Kelvin. The energy density is then:

$$\frac{du}{d\lambda} = -kT \frac{8\pi}{\lambda^4}. \quad (\text{B.6})$$

In function of frequency $\nu = \frac{c}{\lambda}$:

$$\frac{du}{d\nu} = -\frac{c}{\lambda^2} \frac{du}{d\lambda} = \frac{8\pi k T \nu^2}{c^3}, \quad (\text{B.7})$$

also known as the Rayleigh-Jeans law*. Problem: divergence

B.1.2 Quantum approach

The energy levels from a quantized harmonic oscillator are equal to

$$E_r = h\nu \left(r + \frac{1}{2} \right) = \frac{hc}{\lambda} \left(r + \frac{1}{2} \right) \quad \text{with } r = 0, 1, 2, \dots \quad (\text{B.8})$$

Implementing eq. B.4

$$E = \left(r + \frac{1}{2} \right) \frac{hc}{2L} \sqrt{n_1^2 + n_2^2 + n_3^2} \quad (\text{B.9})$$

According to statistical physics the average energy is now not equal to kT but follows a probability distribution

$$p(\nu, r) = \frac{e^{-r h \nu}}{\sum_{r=0}^{\infty} e^{-r h \nu}}, \quad (\text{B.10})$$

where we reference to the ground state of the oscillator: $E'_r = E_r - E_0$.

The average energy is now:

$$\begin{aligned} \langle E(\nu) \rangle &= \sum_{r=0}^{\infty} E(\nu, r) \cdot p(\nu, r) = \frac{\sum_{r=0}^{\infty} r h \nu e^{-r h \nu}}{\sum_{r=0}^{\infty} e^{-r h \nu}} \\ &= \frac{h \nu}{e^{h \nu / k T} - 1} \end{aligned} \quad (\text{B.11})$$

Substituting this for kT in eq. B.7 we find Planck's equation:

$$\frac{du}{d\nu} = \frac{8\pi h \nu^3}{c^3} \frac{h \nu}{e^{h \nu / k T} - 1} \quad (\text{B.12})$$

*This is often quoted per unit of steradian, which results in $\frac{2kT\nu^2}{c^3}$

9. Some useful things for LaTeX

9.1 Definitions

This is an example of a definition. A definition could be mathematical or it could define a concept.

Definition 9.1.1 — Definition name. Given a vector space E , a norm on E is an application, denoted $\|\cdot\|$, E in $\mathbb{R}^+ = [0, +\infty[$ such that:

$$\|\mathbf{x}\| = 0 \Rightarrow \mathbf{x} = \mathbf{0} \quad (9.1)$$

$$\|\lambda\mathbf{x}\| = |\lambda| \cdot \|\mathbf{x}\| \quad (9.2)$$

$$\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\| \quad (9.3)$$

9.2 Remarks

This is an example of a remark.



The concepts presented here are now in conventional employment in mathematics. Vector spaces are taken over the field $\mathbb{K} = \mathbb{R}$, however, established properties are easily extended to $\mathbb{K} = \mathbb{C}$.

9.3 Corollaries

This is an example of a corollary.

Corollary 9.3.1 — Corollary name. The concepts presented here are now in conventional employment in mathematics. Vector spaces are taken over the field $\mathbb{K} = \mathbb{R}$, however, established properties are easily extended to $\mathbb{K} = \mathbb{C}$.

9.4 Propositions

This is an example of propositions.

9.4.1 Several equations

Proposition 9.4.1 — Proposition name. It has the properties:

$$|||\mathbf{x}|| - ||\mathbf{y}||| \leq ||\mathbf{x} - \mathbf{y}|| \quad (9.4)$$

$$|| \sum_{i=1}^n \mathbf{x}_i || \leq \sum_{i=1}^n ||\mathbf{x}_i|| \quad \text{where } n \text{ is a finite integer} \quad (9.5)$$

9.4.2 Single Line

Proposition 9.4.2 Let $f, g \in L^2(G)$; if $\forall \varphi \in \mathcal{D}(G)$, $(f, \varphi)_0 = (g, \varphi)_0$ then $f = g$.

9.5 Examples

This is an example of examples.

9.5.1 Equation and Text

■ **Example 9.1** Let $G = \{x \in \mathbb{R}^2 : |x| < 3\}$ and denoted by: $x^0 = (1, 1)$; consider the function:

$$f(x) = \begin{cases} e^{|x|} & \text{si } |x - x^0| \leq 1/2 \\ 0 & \text{si } |x - x^0| > 1/2 \end{cases} \quad (9.6)$$

The function f has bounded support, we can take $A = \{x \in \mathbb{R}^2 : |x - x^0| \leq 1/2 + \epsilon\}$ for all $\epsilon \in]0; 5/2 - \sqrt{2}[$. ■

9.5.2 Paragraph of Text

■ **Example 9.2 — Example name.** Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris. ■

9.6 Exercises

This is an example of an exercise.

Exercise 9.1 This is a good place to ask a question to test learning progress or further cement ideas into students' minds. ■

9.7 Problems

Problem 9.1 What is the average airspeed velocity of an unladen swallow?

9.8 Vocabulary

Define a word to improve a students' vocabulary.

Vocabulary 9.1 — Word. Definition of word.

Bibliography

Articles

- [1] Morad Aaboud et al. “Measurement of the W -boson mass in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”. In: *Eur. Phys. J.* C78.2 (2018), page 110. doi: [10.1140/epjc/s10052-017-5475-4](https://doi.org/10.1140/epjc/s10052-017-5475-4). arXiv: [1701.07240 \[hep-ex\]](https://arxiv.org/abs/1701.07240) (cited on page 21).
- [2] Timo Antero Aaltonen et al. “Combination of CDF and D0 W -Boson Mass Measurements”. In: *Phys. Rev.* D88.5 (2013), page 052018. doi: [10.1103/PhysRevD.88.052018](https://doi.org/10.1103/PhysRevD.88.052018). arXiv: [1307.7627 \[hep-ex\]](https://arxiv.org/abs/1307.7627) (cited on page 21).
- [3] G. Abbiendi et al. “Search for stable and longlived massive charged particles in e+ e- collisions at $s^{**}(1/2) = 130\text{-GeV}$ to 209-GeV ”. In: *Phys. Lett.* B572 (2003), pages 8–20. doi: [10.1016/S0370-2693\(03\)00639-7](https://doi.org/10.1016/S0370-2693(03)00639-7). arXiv: [hep-ex/0305031 \[hep-ex\]](https://arxiv.org/abs/hep-ex/0305031) (cited on page 31).
- [4] F. Abe et al. “Limits on the production of massive stable charged particles”. In: *Phys. Rev.* D46 (1992), R1889–R1894. doi: [10.1103/PhysRevD.46.R1889](https://doi.org/10.1103/PhysRevD.46.R1889) (cited on page 32).
- [5] K. Abe et al. “First measurement of the left-right charge asymmetry in hadronic Z boson decays and a new determination of $\sin^{**2} \theta_W(\text{eff})$ ”. In: *Phys. Rev. Lett.* 78 (1997), pages 17–21. doi: [10.1103/PhysRevLett.78.17](https://doi.org/10.1103/PhysRevLett.78.17). arXiv: [hep-ex/9609019 \[hep-ex\]](https://arxiv.org/abs/hep-ex/9609019) (cited on page 21).
- [6] Kenji Abe et al. “A High precision measurement of the left-right Z boson cross-section asymmetry”. In: *Phys. Rev. Lett.* 84 (2000), pages 5945–5949. doi: [10.1103/PhysRevLett.84.5945](https://doi.org/10.1103/PhysRevLett.84.5945). arXiv: [hep-ex/0004026 \[hep-ex\]](https://arxiv.org/abs/hep-ex/0004026) (cited on page 21).
- [7] Koya Abe et al. “First direct measurement of the parity violating coupling of the Z0 to the s quark”. In: *Phys. Rev. Lett.* 85 (2000), pages 5059–5063. doi: [10.1103/PhysRevLett.85.5059](https://doi.org/10.1103/PhysRevLett.85.5059). arXiv: [hep-ex/0006019 \[hep-ex\]](https://arxiv.org/abs/hep-ex/0006019) (cited on page 21).
- [8] Koya Abe et al. “An Improved direct measurement of leptonic coupling asymmetries with polarized Z bosons”. In: *Phys. Rev. Lett.* 86 (2001), pages 1162–1166. doi: [10.1103/PhysRevLett.86.1162](https://doi.org/10.1103/PhysRevLett.86.1162). arXiv: [hep-ex/0010015 \[hep-ex\]](https://arxiv.org/abs/hep-ex/0010015) (cited on page 21).
- [9] P. Abreu et al. “Search for stable heavy charged particles in e+ e- collisions at $s^{**}(1/2) = 130\text{-GeV}$ to 136-GeV , 161-GeV and 172-GeV ”. In: *Phys. Lett.* B396 (1997), pages 315–326. doi: [10.1016/S0370-2693\(97\)00152-4](https://doi.org/10.1016/S0370-2693(97)00152-4) (cited on page 31).

- [10] R. Acciarri et al. “Long-Baseline Neutrino Facility (LBNF) and Deep Underground Neutrino Experiment (DUNE)”. In: (2015). arXiv: 1512 . 06148 [physics.ins-det] (cited on page 30).
- [11] D. Acosta et al. “Search for long-lived charged massive particles in $\bar{p}p$ collisions at $\sqrt{s} = 1.8$ TeV”. In: *Phys. Rev. Lett.* 90 (2003), page 131801. doi: 10.1103/PhysRevLett.90.131801. arXiv: hep-ex/0211064 [hep-ex] (cited on page 32).
- [12] H. Aihara et al. “Search for $q = 2/3e$ and $q = 1/3e$ particles produced in e^+e^- annihilations”. In: *Phys. Rev. Lett.* 52 (1984), page 2332. doi: 10.1103/PhysRevLett.52.2332 (cited on page 31).
- [13] R. Akers et al. “Search for heavy charged particles and for particles with anomalous charge in e^+e^- collisions at LEP”. In: *Z. Phys.* C67 (1995), pages 203–212. doi: 10.1007/BF01571281 (cited on page 31).
- [14] M. Tanabashi et al. “Review of Particle Physics”. In: *Phys. Rev. D* 98 (2018), page 030001 (cited on page 7).
- [15] Steven W. Allen, August E. Evrard, and Adam B. Mantz. “Cosmological Parameters from Observations of Galaxy Clusters”. In: *Ann. Rev. Astron. Astrophys.* 49 (2011), pages 409–470. doi: 10.1146/annurev-astro-081710-102514. arXiv: 1103 . 4829 [astro-ph.CO] (cited on page 22).
- [16] M. Ambrosio et al. “Final search for lightly ionizing particles with the MACRO detector”. In: (2004). arXiv: hep-ex/0402006 [hep-ex] (cited on page 32).
- [17] M. Anelli et al. “A facility to Search for Hidden Particles (SHiP) at the CERN SPS”. In: (2015). arXiv: 1504 . 04956 [physics.ins-det] (cited on page 30).
- [18] Ignatios Antoniadis and K. Benakli. “Fractionally charged particles and supersymmetry breaking in 4-D strings”. In: *Phys. Lett.* B295 (1992). [Erratum: *Phys. Lett.* B407,449(1997)], pages 219–224. doi: 10.1016/S0370-2693(97)00755-7, 10.1016/0370-2693(92)91557-P. arXiv: hep-th/9209020 [hep-th] (cited on page 30).
- [19] Gregory G. Athanasiou et al. “Remarks on Wilson Lines, Modular Invariance and Possible String Relics in Calabi-yau Compactifications”. In: *Phys. Lett.* B214 (1988), pages 55–62. doi: 10.1016/0370-2693(88)90451-0 (cited on page 30).
- [21] J. J. Aubert et al. “A Search for Free Quarks in Deep Inelastic Muon Scattering”. In: *Phys. Lett.* 133B (1983), page 461. doi: 10.1016/0370-2693(83)90828-6 (cited on page 31).
- [22] M. Banner et al. “A New Search for Relativistic Particles With Fractional Electric Charge at the CERN $p\bar{p}$ Collider”. In: *Phys. Lett.* 156B (1985), pages 129–133. doi: 10.1016/0370-2693(85)91368-1 (cited on page 32).
- [23] Stephen M. Barr, D. B. Reiss, and A. Zee. “Fractional Charges, Monopoles and Peculiar Photons”. In: *Phys. Rev. Lett.* 50 (1983), page 317. doi: 10.1103/PhysRevLett.50.317 (cited on page 29).
- [24] Marco Battaglieri et al. “US Cosmic Visions: New Ideas in Dark Matter 2017: Community Report”. In: (2017). arXiv: 1707 . 04591 [hep-ph] (cited on page 30).
- [26] James D. Bjorken et al. “New Fixed-Target Experiments to Search for Dark Gauge Forces”. In: *Phys. Rev.* D80 (2009), page 075018. doi: 10 . 1103/PhysRevD . 80 . 075018. arXiv: 0906 . 0580 [hep-ph] (cited on page 30).
- [27] C. Boehm and Pierre Fayet. “Scalar dark matter candidates”. In: *Nucl. Phys.* B683 (2004), pages 219–263. doi: 10 . 1016/j.nuclphysb . 2004 . 01 . 015. arXiv: hep-ph/0305261 [hep-ph] (cited on page 30).
- [28] A. E. Bondar et al. “SEARCH FOR LIGHT PARTICLES WITH CHARGE 2/3 IN e^+e^- ANNIHILATION”. In: *JETP Lett.* 40 (1984). [*Pisma Zh. Eksp. Teor. Fiz.* 40,440(1984)], pages 1265–1268 (cited on page 31).

- [29] David E. Brahm and Lawrence J. Hall. “U(1)-prime DARK MATTER”. In: *Phys. Rev.* D41 (1990), page 1067. DOI: 10.1103/PhysRevD.41.1067 (cited on page 30).
- [30] D. Buskulic et al. “Search for particles with unexpected mass and charge in Z decays”. In: *Phys. Lett.* B303 (1993), pages 198–208. DOI: 10.1016/0370-2693(93)90066-Q (cited on page 31).
- [31] Serguei Chatrchyan et al. “Search for fractionally charged particles in pp collisions at $\sqrt{s} = 7$ TeV”. In: *Phys. Rev.* D87.9 (2013), page 092008. DOI: 10.1103/PhysRevD.87.092008. arXiv: 1210.2311 [hep-ex] (cited on page 31).
- [32] J. H. Christenson et al. “Evidence for the 2π Decay of the K20 Meson”. In: *Physical Review Letters* 13 (July 1964), pages 138–140. DOI: 10.1103/PhysRevLett.13.138 (cited on page 19).
- [37] “Combination of CDF and D0 Results on the Width of the W boson”. In: (2010). arXiv: 1003.2826 [hep-ex] (cited on page 21).
- [38] Edvige Corbelli and Paolo Salucci. “The Extended Rotation Curve and the Dark Matter Halo of M33”. In: *Mon. Not. Roy. Astron. Soc.* 311 (2000), pages 441–447. DOI: 10.1046/j.1365-8711.2000.03075.x. arXiv: astro-ph/9909252 [astro-ph] (cited on page 22).
- [39] C. Davisson and L. H. Germer. “Diffraction of Electrons by a Crystal of Nickel”. In: *Phys. Rev.* 30 (1927), pages 705–740. DOI: 10.1103/PhysRev.30.705 (cited on page 19).
- [40] Alexandre Deur, Stanley J. Brodsky, and Guy F. de Teramond. “The QCD Running Coupling”. In: *Prog. Part. Nucl. Phys.* 90 (2016), pages 1–74. DOI: 10.1016/j.ppnp.2016.04.003. arXiv: 1604.08082 [hep-ph] (cited on page 25).
- [41] Fang-xiao Dong et al. “Fractional Charges, Monopoles and Peculiar Photons in SO(18) GUT Models”. In: *Phys. Lett.* 129B (1983), pages 405–410. DOI: 10.1016/0370-2693(83)90129-6 (cited on page 30).
- [42] John F. Donoghue. “The effective field theory treatment of quantum gravity”. In: *AIP Conf. Proc.* 1483 (2012), pages 73–94. DOI: 10.1063/1.4756964. arXiv: 1209.3511 [gr-qc] (cited on page 23).
- [43] S. M. Faber and R. E. Jackson. “Velocity dispersions and mass to light ratios for elliptical galaxies”. In: *Astrophys. J.* 204 (1976), page 668. DOI: 10.1086/154215 (cited on page 22).
- [44] P. H. Frampton and T. W. Kephart. “Fractionally Charged Particles as Evidence for Supersymmetry”. In: *Phys. Rev. Lett.* 49 (1982), page 1310. DOI: 10.1103/PhysRevLett.49.1310 (cited on page 29).
- [45] S. L. Glashow, J. Iliopoulos, and L. Maiani. “Weak Interactions with Lepton-Hadron Symmetry”. In: *Phys. Rev.* D2 (1970), pages 1285–1292. DOI: 10.1103/PhysRevD.2.1285 (cited on page 19).
- [46] Bob Holdom. “Two U(1)’s and Epsilon Charge Shifts”. In: *Phys. Lett.* 166B (1986), pages 196–198. DOI: 10.1016/0370-2693(86)91377-8 (cited on page 30).
- [47] J. L. Holzbauer. “The Muon g-2 Experiment Overview and Status”. In: *PoS NuFact2017* (2018), page 116. DOI: 10.22323/1.295.0116. arXiv: 1712.05980 [hep-ex] (cited on page 23).
- [48] Eder Izaguirre and Itay Yavin. “New window to millicharged particles at the LHC”. In: *Phys. Rev.* D92.3 (2015), page 035014. DOI: 10.1103/PhysRevD.92.035014. arXiv: 1506.04760 [hep-ph] (cited on page 30).
- [49] Xiang-Dong Jiang and Xian-Jian Zhou. “SO(10) X SO(8) MODEL WITH FRACTIONAL CHARGES, MONOPOLES AND PECULIAR PHOTONS. (IN CHINESE)”. In: *HEPNP* 9 (1985), pages 421–429 (cited on page 30).
- [50] Claus Jönsson. “Elektroneninterferenzen an mehreren künstlich hergestellten Feinspalten”. In: *Zeitschrift für Physik* 161.4 (Aug. 1961), pages 454–474. ISSN: 0044-3328. DOI: 10.1007/BF01342460. URL: <https://doi.org/10.1007/BF01342460> (cited on page 19).

- [51] L. Lyons. “Quark Search Experiments at Accelerators and in Cosmic Rays”. In: *Phys. Rept.* 129 (1985), page 225. DOI: 10.1016/0370-1573(85)90011-0 (cited on page 31).
- [52] Gabriel Magill et al. “Millicharged particles in neutrino experiments”. In: (2018). arXiv: 1806.03310 [hep-ph] (cited on page 30).
- [53] Ziro Maki, Masami Nakagawa, and Shoichi Sakata. “Remarks on the Unified Model of Elementary Particles”. In: *Progress of Theoretical Physics* 28.5 (1962), pages 870–880. DOI: 10.1143/PTP.28.870. eprint: /oup/backfile/content_public/journal/ptp/28/5/10.1143/ptp.28.870/2/28-5-870.pdf. URL: <http://dx.doi.org/10.1143/PTP.28.870> (cited on page 19).
- [54] P G Merli, G F Missiroli, and G Pozzi. “Electron interferometry with the Elmiskop 101 electron microscope”. In: *Journal of Physics E: Scientific Instruments* 7.9 (1974), page 729. URL: <http://stacks.iop.org/0022-3735/7/i=9/a=016> (cited on page 20).
- [55] M. Mori et al. “Search for fractionally charged particles in Kamiokande-II”. In: *Phys. Rev.* D43 (1991), pages 2843–2846. DOI: 10.1103/PhysRevD.43.2843 (cited on page 32).
- [56] Priyamvada Natarajan et al. “Mapping substructure in the HST Frontier Fields cluster lenses and in cosmological simulations”. In: *Mon. Not. Roy. Astron. Soc.* 468.2 (2017), pages 1962–1980. DOI: 10.1093/mnras/stw3385. arXiv: 1702.04348 [astro-ph.GA] (cited on page 22).
- [58] G P. Thomson and Reid AA. “Diffraction of Cathode Rays by a Thin Film”. In: 119 (Jan. 1927), pages 890–890 (cited on page 19).
- [61] Maxim Pospelov, Adam Ritz, and Mikhail B. Voloshin. “Secluded WIMP Dark Matter”. In: *Phys. Lett.* B662 (2008), pages 53–61. DOI: 10.1016/j.physletb.2008.02.052. arXiv: 0711.4866 [hep-ph] (cited on page 30).
- [64] Sabine Riemann. “Precision electroweak physics at high energies”. In: *Rept. Prog. Phys.* 73 (2010), page 126201. DOI: 10.1088/0034-4885/73/12/126201 (cited on page 21).
- [65] S. Schael et al. “Precision electroweak measurements on the Z resonance”. In: *Phys. Rept.* 427 (2006), pages 257–454. DOI: 10.1016/j.physrep.2005.12.006. arXiv: hep-ex/0509008 [hep-ex] (cited on page 21).
- [67] Antoine Weis and Robert Wynands. “Three Demonstration Experiments on the Wave and Particle Nature of Light”. In: 1 (June 2003) (cited on page 20).
- [68] Xiao-Gang Wen and Edward Witten. “Electric and Magnetic Charges in Superstring Models”. In: *Nucl. Phys.* B261 (1985), pages 651–677. DOI: 10.1016/0550-3213(85)90592-9 (cited on page 30).
- [69] Katsuji Yamamoto. “Fractionally Charged Leptons in the SO(14) GUT”. In: *Phys. Lett.* 120B (1983), pages 157–160. DOI: 10.1016/0370-2693(83)90643-3 (cited on page 30).
- [70] Hong-Wei Yu. “SU(8) GUT MODELS WITH STABLE PROTON, N ANTI-N OSCILLATIONS, FRACTIONAL CHARGES AND LOW MASS MONOPOLES”. In: *Phys. Lett.* 142B (1984), pages 42–46. DOI: 10.1016/0370-2693(84)91132-8 (cited on page 29).

Books

- [25] Alessandro Bettini. *Introduction to elementary particle physics*. 2008. ISBN: 9780521880213. URL: <http://www.cambridge.org/catalogue/catalogue.asp?isbn=9780521880213> (cited on page 7).
- [33] J. Clay. *Penetrating radiation*. Volume 30. 1927, pages 1115–1127 (cited on page 35).
- [60] Michael E. Peskin and Daniel V. Schroeder. *An Introduction to quantum field theory*. Reading, USA: Addison-Wesley, 1995. ISBN: 9780201503975, 0201503972. URL: <http://www.slac.stanford.edu/~mpeskin/QFT.html> (cited on page 7).

- [62] B. Povh et al. *Particles and nuclei: An Introduction to the physical concepts*. 1995. doi: 10.1007/3-540-36684-9 (cited on page 7).

Index

C

Corollaries 59

D

Definitions 59

E

Examples 60

 Equation and Text 60

 Paragraph of Text 60

Exercises 60

P

Paragraphs of Text 5

Problems 60

Propositions 59

 Several Equations 60

 Single Line 60

R

Remarks 59

V

Vocabulary 60