

IceCube Neutrino Observatory is the world's largest neutrino detector, located at the geographic South Pole, close to the Amundsen-Scott South Pole Station.

IceCube consists of 5160 optical sensors deployed deep in the Antarctic Ice, covering a volume of 1 km³. On top of this, 81 IceTop detector stations spread over 1 km² are located on the Antarctic plateau.

DeepCore 8 strings with a denser spacing.

Eiffel Tower

UGent members:

S. Verpoest, A. Porcelli, D. Ryckbosch

IceCube Gen 2

Increased InIce volume $1 \text{ km}^3 -> 10 \text{ km}^3$

Additional new surface detectors:

- Scintillators above snowed-in IceTop tanks
 - → measure ionization losses from charged particles
- Imaging Air-Cherenkov Telescopes
- → measure Cherenkov radiation of air shower
- Radio antennas
 - → measure radio waves from air showers

Thesis subjects

IceTop + Scintillators

analysis on the muon component in hadronic interaction models with very inclined showers

 IceTop + IceACT + Radio Antennas analysis on the electromagnetic component (e[±] + γ) in hadronic interaction models with multiple detectors

Cosmic rays


Cosmic ray air shower detection with energies from (1 - 1000) x 10⁶ GeV

Hybrid detection technique:

- Ultra relativistic cosmic ray interacts with atmosphere → secondary particles
- Energy reconstruction using particle density distribution seen by IceTop
- Many relativistic muons can reach the detector simultaneously → muon bundle
- Mass sensitivity from high-energy muon bundle through IceCube
- Multiple hadronic interaction models used in air showers are inconsistent


Thesis subjects

- IceTop + InIce: influence of hadronic interaction models used in air shower simulations
- IceTop: Calibration of the absolute energy scale of the IceTop detector.




Bedrock

Cosmic ray event



Energy spectrum



Average mass

