

Atmospheric Manifestation of Tropical Instability Wave Observed by QuikSCAT and Tropical Rain Measuring Mission

W. Timothy Liu, Xiaosu Xie, Paulo S. Polito

Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA

Shang-Ping Xie

International Pacific Research Center, University of Hawaii, Honolulu

and Hiroshi Hashizume

Graduate School of Environment and Earth Science, Hokkaido University, Sapporo, Japan

Abstract. Observations from two new spaceborne microwave instruments in 1999 clearly reveal the atmospheric manifestation of tropical instability waves north of the Pacific equatorial cold tongue. A unique zonal-temporal band-pass filter enables the isolation of the propagating signals and the determination of their phase differences. The phase differences between the propagation of wind and sea surface temperature (SST) signals observed from space and the vertical wind profiles measured from a research ship are consistent with the hypothesis that the coupling between wind and SST is caused by buoyancy instability and mixing, which reduces the wind shear in the atmospheric boundary layer. The coupling causes higher evaporative cooling over the warm phase and infers a negative thermal feedback.

1. Introduction

Tropical instability waves (TIW) vary in exact location and phase velocity. Such waves were best observed by radiometers on geostationary satellites as meanders of the temperature front between the cold upwelling water of the Pacific equatorial cold tongue and the warm water to the north [e.g., Legeckis, 1977; Yoder et al., 1994]. The waves propagate westward, with period of approximately 30 days, wavelength of 1100 km, and phase speed of 0.5 m/s. The waves are stronger from June to November and during La Nina episodes. Sea surface temperature (SST), however, is often obscured from visible and infrared radiometers by cloud cover. When such sensors are on polar orbiting spacecraft, many days of data are needed to form a composite map of TIW.

The ocean circulation associated with TIW has been well studied using in situ measurements [e.g., Hansen and Paul, 1984; Halpern et al., 1988; Qiao and Weisberg, 1995; Flamant et al., 1996; Baturin and Niler, 1997]. The TIW is believed to be generated by the shear of ocean currents and not by local winds. The atmospheric manifestations of the TIW were observed and inferred in a number of studies [e.g., Hayes et al., 1989; Deser et al., 1993]. The sparsity of historical ship data may have limited their use to seasonal and

interannual variations of air-sea coupling. The studies of Halpern et al. [1988] and Hayes et al., [1989] were based only on mooring data at fixed locations.


Xie et al. [1998] identified TIW in the wind variations observed by the radar scatterometer on the European Remote Sensing (ERS-1) satellite. They found that the wind divergence patterns propagate westward with their centers sandwiched between the warm and cold poles of the SST. A scatterometer measures both wind speed and direction near the ocean surface. The ERS-1 scatterometer scans a 479 km swath and covers only 40% of the global ocean daily and the wind vector retrieved has a resolution of 50 km. The weekly averaged winds used by Xie et al. [1998] can barely resolve TIW and is not adequate for reliable determination of the relative phase between wind and SST, which is key to infer physical mechanism.

Because the atmosphere and the clouds are transparent to microwave, the coincident measurements of two spaceborne microwave instruments during the La Nina episode in the second half of 1999 provided the unprecedented opportunity to observe the influence of the SST front on atmospheric parameters, under both clear and cloudy conditions. The microwave imager on the joint US/Japan Tropical Rainfall Measuring Mission (TRMM) has provided measurements of SST, surface wind speed (WS), and integrated water vapor (WV) in the atmosphere since November 1997. The low-inclination orbit of TRMM is designed to give an optimal sampling rate for monitoring diurnal variations. A radar scatterometer called SeaWinds was launched in June 1999 on NASA's QuikSCAT mission. It has a continuous swath of 1,800 km, providing wind vector measurements at 25 km resolution over 90% of the global ocean daily.

The satellite data also aided the planning of a research cruise on Japan Fishery Agency's R/V Shoyo Maru to observe the vertical structures of the TIW-induced atmospheric waves for the first time. This paper reports briefly on the analysis of the satellite and in situ data with respect to hypotheses on the atmospheric manifestation of oceanic TIW.

2. Hypotheses

There are two hypotheses on the relation between SST and surface winds over tropical oceans. As illustrated by



satellite data. The standard deviation of the wind averages at the surface are 0.5 m/s over warm water and 0.8 m/s over cold water; standard deviations in the lowest 300 m are very consistent with these values. Without band-pass filtering, high-level winds are more susceptible to the influences of non-local circulation. The model simulation by Xie et al. [1988] indicated a vertical phase-shift in the profiles of wind anomalies. The one-time cruise section may not have sufficient sampling to reveal two-dimensional changes.

6. Conclusion

Coincident all-weather observations by a spacebased microwave scatterometer and a microwave radiometer during a La Nina episode reveal clearly the atmospheric manifestation of TIW in surface wind vectors and integrated water vapor. The phase differences between wind components and SST determined from the spacebased data and the wind profiles measured on a research cruise are consistent with the hypothesis that buoyancy instability over warm region of TIW reduces vertical wind shear in the atmospheric boundary layer. LH is found to be in phase with SST and WS with possible negative feedback.

The TIW-induced wind fluctuation is about 2 m/s. The ability to detect the spatial structure of this fluctuation on a daily basis as demonstrated in Section 4 attests unprecedented accuracy and time-space resolution of SeaWinds measurements.

This study was confined to the TIW north of the equator in the Pacific, but similar air-sea coupling is obvious south of the equator in Fig. 1, 3, 4 and 5. Atmospheric responses to the TIW in the South Pacific and the Atlantic were investigated by Hashizume et al. [2000].

Acknowledgments. This study was performed at the Jet Propulsion Laboratory, California Institute of Technology, under contract with NASA. It was supported jointly by the QuikSCAT Project, the Tropical Rain Measuring Mission, and the Earth Observing System Interdisciplinary Science Program of NASA. The TMI data were produced by Frank Wentz who generously made his TMI data openly accessible. Processing of cruise data was performed at University of Hawaii. T. Watanabe, M. Shiotani, M. Fujiwara and the crew of the Japan Fisheries Agency's R/V Shoyo Maru successfully conducted the cruise and acquired the in-situ measurements. TMI data are accessible at <http://www.ssmi.com> and SeaWinds data are available at <http://airsea-www.jpl.nasa.gov/seafux>.

References

Baturin, N.G., and P.P. Niiler, Effects of instability waves in the mixed layer of the equatorial Pacific. *J. Geophys. Res.*, **102**, 27,771-27,793, 1997.

Chelton, D. B., F. J. Wentz, C. L. Gentemann, R. A. de Szoeke, M. G. Schlax, Satellite microwave SST observations of transequatorial tropical instability waves. *Geophys. Res. Lett.*, **27**, 1239-1242, 2000.

Deser, C., J.J. Bates, and S. Wahl, The influence of sea surface temperature gradient on stratiform cloudiness along the equatorial front in the Pacific Ocean. *J. Climate*, **6**, 1172-1180, 1993.

Flamant, P., S.C. Kennean, R.A. Knox, P.P. Niiler, and R. Bernstein, The three-dimensional structure of an upper ocean vortex in the tropical Pacific ocean. *Nature*, **383**, 610-613, 1996.

Halpern, D., R. Knox, and D.S. Luther, Observations of 20-day period meridional current oscillations in the upper ocean along the Pacific Equator. *J. Phys. Oceanogr.*, **18**, 1514-1534, 1988.

Hansen, D.V., and C.A. Paul, Genesis and effects of long waves in the equatorial Pacific. *J. Geophys. Res.*, **89**, 10,431-10,440, 1984.

Hashizume, H., S.-P. Xie, W. T. Liu, and K. Takeuchi, Local and remote atmospheric response to tropical instability waves: a global view from space. *J. Geophys. Res.*, submitted.

Hayes, S.P., M.J. McPhaden, and J.M. Wallace, The influence of sea surface temperature on surface wind in the eastern equatorial Pacific: weekly to monthly variability. *J. Climate*, **2**, 1500-1506, 1989.

Legeckis, R., Long waves in the eastern equatorial Pacific; a view of a geostationary satellite. *Science*, **197**, 1177-1181, 1997.

Lindzen, R.S., and S. Nigam, On the role of sea-surface temperature gradients in forcing low-level winds and convergence in the tropics. *J. Atmos. Sci.*, **44**, 2440-2458, 1987.

Liu, W.T., Moistures and latent heat flux variabilities in the tropical Pacific derived from satellite data. *J. Geophys. Res.*, **93**, 6749-6760, 1988.

Liu, W.T., Remote Sensing of surface turbulence flux. *Surface Waves and Fluxes, Vol. II*, G.L. Geernaert and W.J. Plant (eds), Kluwer Academic, Chapter 16, 293-309, 1990.

Liu, W.T., W. Tang, and P.P. Niiler, Humidity profiles over oceans. *J. Climate*, **4**, 1023-1034, 1991.

Liu, W.T., W. Tang, and P.S. Polito, NASA Scatterometer provides global ocean-surface wind fields with more structures than numerical weather prediction. *Geophys. Res. Lett.*, **25**, 761-764, 1998.

Polito, P.S., O.T. Sato, W.T. Liu, Characterization and validation of the heat storage variability from Topex/Poseidon at four oceanographic sites. *J. Geophys. Res.*, in press, 2000.

Qiao, L., and R.H. Weisberg, Tropical instability wave kinematics: observations from the Tropical Instability Wave Experiment. *J. Geophys. Res.*, **100**, 8677-8693, 1995.

Wallace, J.M., T.P. Mitchell, and C. Deser, The influence of sea-surface temperature on surface wind in the eastern equatorial Pacific: seasonal and interannual variability. *J. Climate*, **2**, 1492-1499, 1989.

Wentz, F.J., A well-calibrated ocean algorithm for special sensor microwave / imager. *J. Geophys. Res.*, **102**, 8703-8718, 1997.

Wentz, F.J., C. Gentemann, D. Smith, D. Chelton, Satellite Measurements of Sea Surface Temperature Through Clouds. *Science*, **288**, 847-850, 2000.

Xie, S.P., M. Ishiwatari, H. Hashizume, and K. Takeuchi, Coupled ocean-atmospheric waves on the equatorial front. *Geophys. Res. Lett.*, **25**, 3863-3966, 1988.

Yoder, J.A., S.G. Ackleson, R.T. Barber, P. Flament, and W. M. Balch, A line in the sea. *Nature*, **371**, 689-692, 1994.

W. T. Liu, X. Xie, Paulo S. Polito, Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, MS 300-323, Pasadena, CA 91109. (e-mail: liu@pacific.jpl.nasa.gov)
Shang-Ping Xie, International Pacific Research Center, University of Hawaii, Honolulu
and Hiroshi Hashizume, Graduate School of Environment and Earth Science, Hokkaido University, Sapporo, Japan

(Received February 29, 2000; revised June 6, 2000; accepted June 14, 2000.)