


Human oscillatory activity during virtual navigation: a comparison between scalp and intracranial recordings


I.O. Korolev; J. Jacobs; M.V. Mollison; M.J. Kahana
Depts. of Psychology and Neuroscience, University of Pennsylvania, Philadelphia, PA



Introduction


- We used electroencephalographic (EEG) recordings to study the modulation of brain oscillations with respect to navigational behavior within a 3D environment.
 - We present a novel comparison of intracranial EEG (iEEG) and scalp EEG (sEEG) data from humans performing the same task.
- Previous findings:**
- Increased oscillatory power during virtual **movement** vs. **standing still** in both cortical (Caplan et al., 2003) and hippocampal regions (Ekstrom et al., 2005).
 - Increased frontal and parieto-temporal theta oscillations at the scalp during goal-directed navigation (Nishiyama & Yamaguchi, 2001; Nishiyama et al., 2002).

Intracranial EEG




- 12 patients (ages 14 - 42) with refractory epilepsy.
- Long-term invasive monitoring to localize seizure onset for subsequent resection.
- Unique opportunity to study electrophysiology of human cognition.

Topographic maps of all cortical electrodes



	Left		Right	
	# Elec.	# Subj.	# Elec.	# Subj.
Frontal	158	7	184	9
Temporal	123	8	154	9
Parietal	34	3	67	7
Occipital	9	3	17	5

Scalp EEG



- 128-channel system from Electrical Geodesics, Inc.
- 200 M Ω high-impedance amplifier.
- Recorded at 500 Hz.

- 16 participants (8 female; ages 19-27).
- Right-handed; normal or corrected-to-normal vision.
- Two participants excluded due to insufficient data.

The YellowCab Task




- Participants played the role of a taxi-driver in a virtual town, looking for passengers ("Searching") and delivering them to requested destinations ("Goal-seeking") (Caplan et al., 2003; Newman et al., 2005; Ekstrom et al., 2005).

Hypotheses


- An increase in oscillations during **movement**, particularly during the "Searching" phase (Caplan et al., 2003).
- Similarities between iEEG and sEEG topographies.

Analytical Methods

Detecting Oscillatory Episodes






- $P_{episode}$** (Caplan et al., 2001):
 - Method of quantifying oscillatory activity
 - Indicates proportion of time that an oscillation at frequency f appears above background EEG activity.






$P_{episode}$ at a right temporal electrode in an iEEG patient during the "Searching" phase.

- Region of Interest (ROI) Analysis:**
 - Data combined across participants at each frequency band (5 bands; 24 bins; 1 - 54 Hz) and ROI (Right/Left Frontal, Parietal, Temporal, Occipital).

iEEG Results



sEEG Results



Discussion

- Unidirectional increase in oscillations at both scalp and cortical surfaces during **movement**.
 - More powerful increase during "Searching" in iEEG.
- Difference between "Searching" and "Goal-seeking" topographies.
- There is not a perfect concordance between iEEG and sEEG topographies.
 - iEEG oscillations are often locally generated (Raghavachari et al., 2005).
 - Observable activity at the scalp requires synchronous neuronal activity.

ROI Analysis Details

- For each participant:**
- P -values (non-parametric test) at each electrode and frequency bin converted to z-scores.
 - Average z-scores within each ROI.
 - Average z-scores across frequency bins within each band for each ROI.
- Across participants:**
- One-sample t -test on mean z-scores for each ROI and frequency band.

Acknowledgements

- This poster was typeset in L^AT_EX₂ using the posterboxen style and TikZ.
- This work was sponsored by NIH grants (R01-MH055687 and R01-MH061975) to MJK, and by an NSF CELEST grant to Boston University (SBE0354378).
- Members of the Computational Memory Lab.
- Patients and their families.

References

- Caplan, J. B., Madsen, J. R., Raghavachari, S., & Kahana, M. J. (2001). Distinct patterns of brain oscillations underlie two basic parameters of human maze learning. *Journal of Neurophysiology*, 86, 368-380.
- Caplan, J. B., Madsen, J. R., Schulze-Bonhage, A., Aschenbrenner-Scheib, R., Newman, E. L., & Kahana, M. J. (2003). Human theta oscillations related to sensorimotor integration and spatial learning. *Journal of Neuroscience*, 23, 4726-4736.
- Ekstrom, A., Caplan, J. B., Ho, E., Shattuck, K., Fried, I., & Kahana, M. (2005). Human hippocampal theta activity during virtual navigation. *Hippocampus*.
- Newman, E. L., Caplan, J. B., Kirschen, M. P., Korolev, I. O., Sekuler, R., & Kahana, M. J. (2005). Learning your way around town: How virtual taxicab drivers learn to use both layout and landmark information. *Cognition*, in press.
- Nishiyama, N., Mizuhara, H., Mawakechi, F., & Yamaguchi, Y. (2002). Theta episodes observed in human scalp eeg during virtual navigation - spatial distribution and task dependence. *Proceedings of the 9th International Conference on Neural Information Processing*, 1.
- Nishiyama, N., & Yamaguchi, Y. (2001). Human eeg theta in the spatial recognition task. *Presented at the 5th World Multi-Conference on Systems, Cybernetics and Informatics*, Orlando, Florida.
- Raghavachari, S., Lisman, J. E., Tully, M., Madsen, J. R., Bromfield, E. B., & Kahana, M. J. (2005). Theta oscillations in human cortex during a working memory task: Evidence for local generators. Manuscript submitted for publication.

65.16