



Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Development of an RF resonator for a double junction ion trap

Semester Project

Konstantin Nesterov

Sunday 30th December, 2018

Advisors: Prof. Dr. J. Home, Chiara Decaroli

Department of Physics, ETH Zürich

Abstract

This paper shows complete process of modeling, designing and testing an RF helical resonator suitable for supporting a double-junction ion trap in a cryogenic environment.

Contents

Contents	ii
1 Introduction	1
1.1 Why do we need resonators?	1
1.2 Context of a project	2
1.3 Kinds of resonators	2
1.3.1 Helical	2
1.3.2 RLC	2
1.3.3 Crystal	2
2 Theory	3
2.1 Helical resonator models	3
2.2 Comparing Macalpine's vs Hensinger's	3
3 Design	4
4 Validation	5
5 External circuits & additional features	6
Bibliography	7

Chapter 1

Introduction

Quantum computing is an exciting and rapidly evolving field of a modern science. One of the popular implementations of a quantum computer is based on an ability to control and measure systems of trapped ions. Those ions are typically confined in Paul [4] traps by applying static and high voltage RF signals to trap's electrodes, which on average generates a corresponding stabilizing electric potential.

1.1 Why do we need resonators?

One could potentially couple a radio frequency source directly to an ions' trap. However it imposes the following challenges:

- noise from a source may contribute to heating of trapped ions [5]
- in order to maintain efficient cryostat cooling all internal connections must be manufactured out of materials with low thermal conductivity. Unfortunately following the Wiedemann-Franz law [1] it leads to low electrical conductivity thus resulting in high dissipation of power
- impedance mismatch between source and trap leads to an additional dissipation of RF power

These issues can be avoided by placing an amplifier close to Paul trap, which would filter incoming signal and output it with voltage suitable for operating the trap. Two available options are active and passive amplifiers. Active amplifiers perform better in terms of voltage gain at room temperatures, but cryogenic temperatures do not allow underlying semiconductor technology to function, which leaves us with passive amplifiers (resonators).

1.2 Context of a project

This semester project aims to be a part of an attempt to create a scalable quantum computing architecture by Chiara Decaroli. It provides the following benefits compared to existing solutions:

- using subtractive laser writing to manufacture wafers eliminates misalignment effects
- double junction ion trap was designed for parallel operations, Decoherence Free Subspace (DFS) ion transport across the junctions, and manipulation of long chains of ions
- integrated laser delivery through optical lensed fibers eliminates the need for bulky optics and custom objectives which do not provide an easy scalability

[TRAP PICTURE HERE]

1.3 Kinds of resonators

Required frequency range limits our selection to the following types of resonators: helical [3] or, for higher frequencies, coaxial, RLC [2], and crystal oscillators.

1.3.1 Helical

1.3.2 RLC

1.3.3 Crystal

Chapter 2

Theory

2.1 Helical resonator models

2.2 Comparing Macalpine's vs Hensinger's

Chapter 3

Design

Chapter 4

Validation

Chapter 5

External circuits & additional features

Bibliography

- [1] R. Franz and G. Wiedemann. Über die Wärme-Leitungsfähigkeit der Metalle. *Annalen der Physik*, 165(8):497–531, 1853.
- [2] Davide Gandolfi. *Compact RF Amplifier for Scalable Ion-Traps*. PhD thesis, Università degli Studi di Trento and Universität Innsbruck, 2010.
- [3] S. Gulde. *Experimental realization of quantum gates and the Deutsch-Josza algorithm with trapped ^{40}Ca -ions*. PhD thesis, University of Innsbruck, 2017.
- [4] Wolfgang Paul. Electromagnetic Traps for Charged and Neutral Particles. *Reviews of Modern Physics*, 62(3):531–540, 1990.
- [5] Q. A. Turchette, D. Kielpinski, B. E. King, D. Leibfried, D. M. Meekhof, C. J. Myatt, M. A. Rowe, C. A. Sackett, C. S. Wood, W. M. Itano, C. Monroe, and D. J. Wineland. Heating of trapped ions from the quantum ground state. *Physical Review A - Atomic, Molecular, and Optical Physics*, 61(6):8, 2000.



Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Declaration of originality

The signed declaration of originality is a component of every semester paper, Bachelor's thesis, Master's thesis and any other degree paper undertaken during the course of studies, including the respective electronic versions.

Lecturers may also require a declaration of originality for other written papers compiled for their courses.

I hereby confirm that I am the sole author of the written work here enclosed and that I have compiled it in my own words. Parts excepted are corrections of form and content by the supervisor.

Title of work (in block letters):

Authored by (in block letters):

For papers written by groups the names of all authors are required.

Name(s):

First name(s):

With my signature I confirm that

- I have committed none of the forms of plagiarism described in the '[Citation etiquette](#)' information sheet.
- I have documented all methods, data and processes truthfully.
- I have not manipulated any data.
- I have mentioned all persons who were significant facilitators of the work.

I am aware that the work may be screened electronically for plagiarism.

Place, date

Signature(s)

For papers written by groups the names of all authors are required. Their signatures collectively guarantee the entire content of the written paper.