VIJAYA DIAGNOSTIC CENTRE®

H No. 2-137/10, Plot No.42, NH65, Opposite R.S. Brothers, Gangaram, Chanda Nagar, Hyderabad - 500050, Telangana

TEST REPORT

Name	: **Ms. TAYYABA SIDDIQUA**		Registered on : 11-Feb-2026 17:41
Age/Gender	: **23 Years / Female**	BirthDate : 10-Apr-2002	Collected on : 11-Feb-2026 17:48
Registration ID	: **260660003332**		Released on : 12-Feb-2026 09:38
Ref. By	: Dr AMIT GOEL		Printed on : 12-Feb-2026 09:54
Sample Type	: Serum		Regn Centre : **Gachibowli-66**

THYROID PROFILE

TEST NAME		RESULT	UNIT	BIOLOGICAL REFERENCE INTERVAL
T3 Total *Method: Chemiluminiscence Immunoassay*	:	0.73	ng/mL	Non pregnant : 0.60 - 1.81 Pregnant 1st Trimester : 0.81 - 1.90 2nd & 3rd Trimester : 1.0 - 2.60
T4 Total *Method: Chemiluminiscence Immunoassay*	:	9.70	µg/dL	Adult : 3.2 - 12.6 Pregnant : 6.4 - 10.7
TSH - Ultrasensitive *Method: Chemiluminiscence Immunoassay*	:	1.575	µIU/mL	Non pregnant : 0.55 - 4.78 Pregnant 1st Trimester : 0.3 - 4.5 2nd Trimester : 0.5 - 4.6 3rd Trimester : 0.8 - 5.2

Interpretation / Comments :

- Patient preparation is particularly important for hormone studies, results of which may be markedly affectedby many factors such as stress, position, fasting state, time of the day, preceding diet and drug therapy.
- The levels of T3 helps in the diagnosis of T3 thyrotoxicosis and monitoring the course of hyperthyroidism.T3 is not recommended for diagnosis of hypothyroidism as decreased values have minimal clinical significance.
- Values below the lower limits can be caused by a number of conditions including non-thyroidal illness, acute and chronic stress and hypothyroidism.
- Elevated level of T4 are seen in hyperthyroidism, pregnancy, euthyroid patients with increased serum thyroxine binding globulin.
- Decreased levels are noted in hypothyroidism, hypoproteinemia, euthyroid sick syndrome, decrease in thyroxine binding globulin.
- TSH levels are increased in primary hypothyroidism, insufficient thyroid hormone replacement therapy,Hashimotos thyroiditis, use of amphetamines, dopamine antagonists, iodine containing agents, lithium and iodine induced or deficiency goiter.
- Decreased levels of TSH may be seen in graves disease, toxic multinodular goitre, thyroiditis, excessivetreatment with thyroid hormone replacement and central hypothyroidism.

MC-7476

DR. R PAVAN KUMAR
MD BIOCHEMISTRY
Registration No: 70314

---------- End of Report ----------

* Suggested Clinical Correlation, If Necessary Kindly Discuss with signatory

Customer Care ✆ 040-2100 0000

The Pioneers in Diagnostic Medicare...
Terms and Conditions of Reporting mentioned overleaf