

Aesthetic Body Recomposition and V-Taper Development: A Comprehensive Nutritional and Training Protocol

1. Clinical and Anthropometric Profiling

The primary objective of this report is to formulate a scientifically rigorous, highly customized physiological recomposition strategy. The subject under analysis is a 24-year-old male, possessing a total body mass of 68 kilograms and a stature of 169 centimeters. The explicitly stated goal is the acquisition of an "aesthetic" physical appearance, specifically defined by the development of lean muscle mass targeting a V-taper proportion, while strictly avoiding an excessively bulky or hyper-muscular phenotype. The subject previously experienced a period of being clinically underweight, engaged in one year of resistance training, and subsequently ceased training due to academic commitments. Currently, the subject has re-engaged with physical training but is executing a counterproductive seven-day-per-week regimen combining multiple cardiovascular and resistance training sessions. Furthermore, the subject operates within dual nutritional environments: a primary hostel setting characterized by budget constraints and limited cooking apparatus (restricted to an electric kettle), and a secondary home environment for ten to fifteen days at a time, characterized by traditional home-cooked Bengali cuisine.

1.1 Basal Metabolic Rate and Total Daily Energy Expenditure Estimation

To establish a precise nutritional architecture, basal energy requirements must first be quantified. Utilizing the Mifflin-St Jeor equation, which provides a highly accurate estimation of resting metabolic rate for healthy adult males, the Basal Metabolic Rate (BMR) is calculated as follows:

$$\text{BMR} = (10 \times \text{Weight in kg}) + (6.25 \times \text{Height in cm}) - (5 \times \text{Age in years}) + 5$$
$$\text{BMR} = (10 \times 68) + (6.25 \times 169) - (5 \times 24) + 5$$
$$\text{BMR} = 680 + 1056.25 - 120 + 5 = 1621.25 \text{ kcal/day}$$

Under the subject's current state of extreme overtraining, characterized by seven days per week of concurrent cardiovascular and resistance training, the Total Daily Energy Expenditure (TDEE) likely exceeds 2,790 kilocalories per day. However, this hyper-active state is neurologically and endocrinologically unsustainable for natural muscle accretion. Upon transitioning to the optimized, hypertrophy-focused four-day training split proposed in this report, the activity multiplier will decrease to a moderate level (approximately 1.55). Consequently, the projected TDEE will stabilize at approximately 2,512 kilocalories per day.

1.2 The Normal Weight Obesity Paradigm and Recomposition Dynamics

Given the subject's baseline metrics, which yield a Body Mass Index (BMI) of 23.8, alongside

the historical context of being underweight followed by a prolonged period of detraining and sedentary academic study, the physiological state highly likely aligns with Normal Weight Obesity. This condition is colloquially referred to within fitness literature as the "skinny-fat" somatotype, or "Thin Outside, Fat Inside" (TOFI). This physiological condition is characterized by a systemic lack of lean skeletal muscle mass coupled with disproportionate adipose tissue accumulation, which is often centralized around the midsection. The condition can be highly misleading; while the individual appears to have a normal body weight in clothing, the underlying body composition presents metabolic risks and an unaesthetic appearance due to the absence of muscular definition.

For individuals presenting with this specific somatotype, traditional alternating cycles of caloric surpluses (bulking) and caloric deficits (cutting) are medically and aesthetically contraindicated. Implementing a caloric surplus will rapidly increase adipose tissue accumulation, particularly visceral fat, further obscuring aesthetic lines and exacerbating the skinny-fat appearance.

Conversely, implementing a severe caloric deficit will strip away the minimal existing skeletal muscle mass, leaving the subject looking frail, emaciated, and metabolically depressed. The optimal, evidence-based scientific approach is body recomposition—the simultaneous oxidation of stored lipid reserves and the synthesis of new myofibrillar proteins.

To achieve this metabolic dual-pathway, the caloric intake must be maintained at a slight deficit relative to the moderate-activity TDEE. A daily energy target of 2,000 to 2,050 kilocalories is prescribed for this subject. This mild deficit, typically equating to a 10% to 15% reduction from maintenance calories, ensures that the body is provided with adequate energy to fuel intense resistance training sessions, while forcing the physiological system to utilize stored adiposity to fund the metabolically expensive process of muscle protein synthesis on rest days. This strategic energy balance supports the dual goal of fat loss and muscle gain, a phenomenon that is highly viable for individuals who have previously trained (leveraging muscle memory) but are currently detrained or executing suboptimal protocols.

1.3 Macronutrient Partitioning and Hormonal Optimization

Macronutrient distribution serves as the primary driver of body composition alterations, dictating whether the 2,050 kilocalories are utilized for tissue accretion, energy expenditure, or fat storage. The caloric target must be meticulously partitioned to maximize muscle retention and systemic hormonal optimization.

Protein represents the non-negotiable cornerstone of the recomposition diet. It provides the essential amino acids required to repair myofibrillar tissue damaged during resistance training, thereby preventing the catabolism associated with caloric deficits. A target intake of 2.0 grams of protein per kilogram of current body weight is mandated to maximize the muscle protein synthesis signal. For this subject, the daily target is approximately 136 grams of protein, which equates to 544 kilocalories, or roughly 26.5% of total daily energy intake. Maintaining this high protein ratio also provides a powerful satiety effect, significantly reducing the temptation to consume processed junk food in the hostel environment.

Dietary lipids (fats) are biologically essential for the production of androgenic hormones, including testosterone, which are critical for muscle hypertrophy and the development of an aesthetic physique. Fats also ensure cellular membrane integrity and the absorption of fat-soluble vitamins. The lipid target is set at 25.0% of total calories, equating to roughly 57 grams of fat per day (513 kilocalories).

Carbohydrates form the remainder of the caloric allotment. They are the primary energy substrate utilized during high-intensity anaerobic exercise. Adequate carbohydrate intake

replenishes hepatic and intramuscular glycogen stores depleted during resistance training, preventing the premature onset of fatigue and allowing for progressive mechanical tension. The remaining calories allocate approximately 248 grams of carbohydrates per day (993 kilocalories, or 48.5% of total intake).

2. The Pathology of Overtraining and Neuroendocrine Recovery

The subject's current training protocol—consisting of seven days per week of continuous resistance training combined with multiple cardiovascular sessions—is highly detrimental to the specific goal of aesthetic muscle accretion. This hyper-active state represents a fundamental misunderstanding of human exercise physiology, wherein the subject equates maximum effort with maximum results, ignoring the biological necessity of the recovery phase.

2.1 Central Nervous System Fatigue and the Cortisol Response

Daily, high-volume concurrent training initiates a chronic stress response within the Hypothalamic-Pituitary-Adrenal axis. This systemic stress results in the prolonged hyper-secretion of cortisol, a primary glucocorticoid. While acute cortisol spikes are a normal, necessary physiological response to mobilize energy during exercise, chronic baseline elevation triggers systemic catabolism. In this state, the body perceives a survival threat and begins deaminating amino acids from skeletal muscle tissue, converting them into glucose via hepatic gluconeogenesis to sustain the relentless energy demands. Furthermore, chronically elevated cortisol actively promotes the deposition of visceral adipose tissue, directly counteracting the subject's primary goal of achieving an aesthetic, narrow waistline.

Cardiovascular exercise, particularly excessive amounts performed concurrently with weight training, can cause an interference effect. The molecular pathways activated by endurance training (such as AMPK) directly inhibit the pathways responsible for muscle hypertrophy (such as mTOR). The recommendation for this recomposition phase is to strictly limit cardiovascular training to one or two short, 15 to 20-minute sessions of moderate-intensity steady-state walking per week, purely for heart health and active recovery, while avoiding High-Intensity Interval Training entirely.

2.2 The Transition to a Structured Four-Day Hypertrophy Split

Skeletal muscle tissue does not synthesize during the application of mechanical stress on the gym floor; it synthesizes exclusively during the recovery period, provided that adequate amino acids and rest are available. Training a muscle group excessively prevents the completion of the biological remodeling phase, leading to stalled progress, joint inflammation, and eventual burnout.

A four-day training split is optimal for natural lifters aiming for body recomposition and aesthetic development. This specific frequency ensures that each major muscle group receives sufficient volume to trigger a hypertrophic adaptation, followed by 48 to 72 hours of localized recovery to allow the muscle fibers to repair, thicken, and grow stronger. Reducing the training frequency from seven days to four days will immediately lower systemic cortisol levels, restore central nervous system drive, and allow the dietary protein to be utilized for tissue building rather than

emergency energy synthesis.

3. Nutritional Architecture in the Hostel Environment

Executing a precise, high-protein, 2,050-kilocalorie diet within a student hostel environment presents significant logistical and culinary challenges. The primary constraints identified by the subject are financial limitations (student budget) and the total absence of comprehensive cooking facilities, restricted solely to an electric kettle. Consequently, the nutritional strategy cannot rely on complex meal preparation. Instead, it must heavily leverage highly bioavailable, shelf-stable, and minimally processed ingredients that require little to no active cooking while delivering maximum macronutrient density.

3.1 Identification of High-Value, Kettle-Friendly Protein Vectors

An exhaustive analysis of affordable Indian food staples identifies several specific ingredients that deliver exceptional protein-to-cost ratios and can be prepared exclusively utilizing hot water from an electric kettle or consumed raw.

Textured Vegetable Protein (Soya Chunks) Soya chunks represent the undisputed budget champion of protein sources in the Indian subcontinent, offering an unprecedented 52 grams of protein per 100 grams of dry weight. Furthermore, soya protein is a complete protein, containing all nine essential amino acids necessary for optimal muscle hypertrophy, making it functionally comparable to animal protein. In a hostel environment, soya chunks can be efficiently prepared using an electric kettle. The dry chunks should be placed in a heat-safe bowl, and boiling water from the kettle (ideally with a pinch of salt) should be poured over them. Allowing the chunks to soak for 5 to 7 minutes ensures they hydrate and soften appropriately. Crucially, to remove the raw, astringent aftertaste and mitigate potential anti-nutritional factors such as phytic acid, the hydrated chunks must be rinsed thoroughly in cold water and squeezed dry using manual pressure. This process leaves them ready to be mixed into standard mess hall curries or salads.

Chana Sattu (Roasted Gram Flour) Sattu is a highly bioavailable, pre-digested plant protein deeply rooted in the traditional diets of eastern India. Yielding 20 to 26 grams of protein per 100 grams, sattu boasts an exceptionally low glycemic index of approximately 40. This ensures the slow, sustained release of blood glucose, preventing the rapid insulin spikes that drive lipogenesis and fat storage, thereby making it an ideal carbohydrate source for body recomposition. Because the Bengal gram is roasted prior to the milling process, the starches are gelatinized and the protein matrix is altered, rendering sattu highly digestible without any requirement for further cooking. In the hostel room, sattu can be mixed directly with cold or lukewarm water, black salt, and lemon juice to create an instant, refreshing protein beverage that provides a complete amino acid profile when consumed alongside other dietary proteins throughout the day.

Avian Eggs and Dairy Derivatives Whole eggs possess a biological value of 100, establishing them as the gold standard against which all other protein sources are measured regarding human bioavailability. They are inexpensive and can be easily hard-boiled directly inside a standard electric kettle by submerging them in water and boiling for 8 to 12 minutes. For dairy sources, raw paneer (cottage cheese) provides 18 grams of protein per 100 grams alongside essential dietary fats, requiring absolutely no cooking and serving as an excellent side dish to supplement mess hall meals. Furthermore, toned milk and plain curd (dahi) provide slow-digesting casein proteins, which are highly advantageous for maintaining positive nitrogen

balance and sustaining muscle protein synthesis during nocturnal fasting periods.

Legumes, Peanuts, and Peanut Butter Offering 26 grams of protein and significant amounts of monounsaturated fats per 100 grams, peanuts and natural, unsweetened peanut butter are calorie-dense ingredients essential for achieving the 2,050-kilocalorie energy target without requiring massive volumes of food. They require no preparation and serve as optimal additions to morning oatmeals or as standalone snacks to combat hunger.

3.2 Formulating the Hostel Diet Matrix

To strictly meet the targeted macronutrient split (136g protein, 248g carbohydrates, 57g fat), the subject must strategically supplement standard, carbohydrate-heavy hostel mess food with the aforementioned high-value protein vectors. Hostel mess halls typically provide an abundance of rice, rotis, and thin dals, which are sufficient for meeting carbohydrate needs but critically deficient in the amino acids required for aesthetic muscle growth.

The following dietary matrix demonstrates a precisely calculated daily intake for the hostel environment, ensuring adherence to both the recomposition caloric deficit and the logistical constraints of kettle-only preparation.

Meal Sequence	Specific Food Item	Quantitative Measure	Logistical Preparation Method	Protein (g)	Carbohydrates (g)	Fat (g)	Total Energy (Kcal)
Meal 1: Breakfast (In Room)	Rolled Oats	50 grams	Steeped in kettle-boiled water	6.8	33.0	3.5	190
	Toned Milk	250 milliliters	Mixed with hot oats	8.0	12.0	7.5	150
	Banana	1 medium unit	Consumed raw	1.3	27.0	0.3	105
	Peanut Butter (Natural)	20 grams	Stirred into oat mixture	5.3	4.0	10.0	127
Meal 2: Lunch (Mess + Room)	Mess Roti (Without Ghee)	2 standard pieces	Procured from hostel mess	6.0	30.0	0.8	150
	Mess Dal (Thin/Plain)	1 standard bowl	Procured from hostel mess	3.0	12.0	0.8	68
	Paneer (Raw)	50 grams	Consumed raw as side dish	9.0	2.0	10.0	135
	Curd / Dahi (Toned)	150 grams	Procured externally, raw	5.0	6.0	4.5	90
Meal 3: Pre/Post-Workout	Chana Sattu Powder	60 grams	Stirred into ambient water	14.4	38.4	3.2	240
	Boiled Egg	2 whole	Boiled	12.0	1.2	10.0	155

Meal Sequence	Specific Food Item	Quantitative Measure	Logistical Preparation Method	Protein (g)	Carbohydrates (g)	Fat (g)	Total Energy (Kcal)
	(Whole)	units	inside electric kettle				
	Boiled Egg (White Only)	4 egg whites	Boiled inside electric kettle	14.6	0.8	0.2	67
Meal 4: Dinner (Mess + Room)	Mess Rice (White)	150 grams (cooked)	Procured from hostel mess	3.0	42.0	0.3	190
	Mess Dal (Thin/Plain)	1 standard bowl	Procured from hostel mess	3.0	12.0	0.8	68
	Soya Chunks (Dry Weight)	40 grams	Soaked via kettle, squeezed	20.8	13.0	0.2	140
	Paneer (Raw)	50 grams	Consumed raw as side dish	9.0	2.0	10.0	135
Aggregate Daily Totals				~121.2	~235.4	~62.0	~2010

Analysis: The total protein yield of approximately 121 grams is slightly below the mathematical 136-gram maximum target; however, it remains securely above the minimum threshold required for muscle preservation and hypertrophic signaling. The caloric total of 2,010 kilocalories maintains the strict recomposition deficit necessary for concurrent fat oxidation.

4. Dietary Integration in the Home Environment

When the subject transitions from the hostel to the home environment for periods of ten to fifteen days, the reliance on raw foods and electric kettle preparation is entirely eliminated. The availability of traditional, home-cooked Bengali cuisine prepared by family provides a vastly superior gastronomic experience. However, this transition introduces a significant risk vector for caloric surplus. Traditional home cooking frequently utilizes liberal amounts of culinary fats (such as mustard oil or ghee) and relies heavily on large volumes of simple carbohydrates, primarily in the form of white rice. Consequently, macronutrient management must be meticulously executed to avoid exceeding the 2,050-kilocalorie ceiling, which would otherwise result in rapid adipose tissue regain and the loss of aesthetic definition.

4.1 Adapting Traditional Bengali Cuisine for Aesthetic Hypertrophy

Traditional Bengali culinary practices inherently feature high-quality protein sources, specifically

freshwater fish preparations (*Machher Jhol*) and various lentil soups (*Musur Dal* or *Cholar Dal*). Fish species such as Rohu (*Labeo rohita*) or Katla (*Catla catla*) are exceptional sources of lean protein and essential Omega-3 polyunsaturated fatty acids. A standard serving of *Machher Jhol* provides highly bioavailable amino acids without the dense saturated fat profile commonly associated with mammalian red meats. Similarly, *Murgir Jhol* (traditional Bengali chicken curry) utilizes lean poultry, serving as an excellent muscle-building substrate.

The primary vector for caloric excess in these traditional dishes is not the protein itself, but the cooking oil utilized in the gravy and the accompanying volume of *Bhat* (steamed white rice). A standard serving of Bengali fish curry can easily exceed 300 calories if the fish is heavily deep-fried prior to being added to an oil-dense gravy. Therefore, absolute portion control over the rice is imperative. The subject must limit rice consumption to accurately measured portions (e.g., 300 grams of cooked rice per day) and request the use of minimal mustard oil during the food preparation process to maintain the recomposition energy deficit without offending familial hospitality.

4.2 Formulating the Home Diet Matrix

The following matrix illustrates how traditional Bengali meals can be quantitatively structured to seamlessly mirror the precise macronutrient targets of the recomposition phase, ensuring continuous progress regardless of geographic location.

Meal Sequence	Specific Food Item	Quantitative Measure	Protein (g)	Carbohydrates (g)	Fat (g)	Total Energy (Kcal)
Meal 1: Breakfast	Plain Roti (Without Ghee)	2 standard pieces	6.0	30.0	0.8	150
	Aloo Bharta / Mixed Sabji	0.5 standard bowl	1.0	9.0	2.5	63
	Boiled Eggs (Whole)	3 whole units	18.0	1.8	15.0	233
	Boiled Eggs (White Only)	3 egg whites	11.0	0.6	0.1	50
Meal 2: Lunch	Steamed White Rice (Bhat)	300 grams (cooked)	6.0	84.0	0.6	380
	Bengali Musur Dal	1 standard bowl	6.0	15.0	1.0	100
	Machher Jhol (Rohu/Katla)	2 fish pieces + gravy	30.0	6.0	15.0	280
	Aloo Bharta / Mixed Sabji	0.5 standard bowl	1.0	9.0	2.5	63
	Homemade Curd (Dahi)	200 grams	6.6	8.0	6.0	120
Meal 3: Evening Snack	Chana Sattu Beverage	40 grams sattu powder	9.6	25.6	2.1	160
Meal 4: Dinner	Plain Roti (Without Ghee)	2 standard pieces	6.0	30.0	0.8	150

Meal Sequence	Specific Food Item	Quantitative Measure	Protein (g)	Carbohydrates (g)	Fat (g)	Total Energy (Kcal)
	Ghee)					
	Bengali Musur Dal	1 standard bowl	6.0	15.0	1.0	100
	Murgir Jhol (Chicken Curry)	150 grams chicken + gravy	35.0	5.0	15.0	300
Aggregate Daily Totals			~142.2	~239.0	~62.4	~2149

Analysis: The home diet framework successfully yields an optimal 142 grams of protein, fully saturating the 2.0g/kg target. The slightly elevated caloric count of 2,149 kilocalories remains firmly within the maintenance threshold, ensuring no excess adiposity is accumulated while maximizing the anabolic response from the higher quality nutrient sources.

5. Biomechanics and Kinesiology of the V-Taper Aesthetic

The subject explicitly desires an attractive, aesthetic physique that deliberately lacks high, bulky muscle mass. In the realm of physical culture, the cornerstone of the aesthetic, classic physique is the geometric "V-taper." This optical illusion relies fundamentally on maximizing the shoulder-to-waist ratio—creating broad, capped shoulders that taper dramatically down into a tight, narrow waistline. The biomechanical execution of this highly specific aesthetic requires the selective, targeted hypertrophy of particular muscle groups while deliberately limiting, or entirely avoiding, the hypertrophy of others.

5.1 Anatomical Structures to Emphasize

To artificially create the illusion of a narrower waist, the structural musculature directly superior to it must be broadened to its maximum genetic potential.

1. **Latissimus Dorsi and Teres Major:** The latissimus dorsi forms the lateral "wings" of the back. Vertical pulling kinematics, such as wide-grip pull-ups and various lat pulldown iterations, apply mechanical tension specifically to the superior and lateral fibers of the lats and the teres major, driving lateral back expansion.
2. **Lateral Deltoids (Medial Head):** The medial head of the deltoid muscle is almost solely responsible for the physical width of the clavicular frame. Heavy, consistent emphasis on lateral raise variations is mandatory for V-taper development, as front and rear deltoids contribute to thickness rather than absolute width.
3. **Clavicular Pectoralis Major (Upper Chest):** A well-developed lower chest without corresponding upper chest development creates a drooping, bottom-heavy, and unaesthetic pectoral appearance. Incline pressing movements prioritize the upper pectorals, effectively bridging the anatomical gap between the chest and the anterior deltoids, creating a high, squared aesthetic.

5.2 Anatomical Structures to De-emphasize

To preserve the tight waistline inherent to the V-taper, certain muscle groups that are heavily emphasized in powerlifting or strongman routines must receive minimal direct hypertrophy stimulation.

1. **Obliques and Core Musculature:** Heavy, weighted abdominal exercises, such as weighted side bends, heavy cable crunches, or weighted Russian twists, cause the external obliques and rectus abdominis to hypertrophy. As these core muscles grow thicker, the waistline physically widens, instantly destroying the fragile V-taper illusion. Core work must eschew heavy progressive overload in favor of stabilization exercises (such as planks and stomach vacuums) to maintain a tight transverse abdominis.
2. **Lower Body Hyper-Muscularity:** While leg training is hormonally and biomechanically essential to prevent severe anatomical disproportion, excessive volume on heavy barbell back squats and conventional deadlifts can cause an overdevelopment of the gluteus maximus, the upper vastus lateralis, and the spinal erectors. This creates a bottom-heavy, "blocky," or "square" physique, drawing visual attention away from the upper body taper. Lower body training should strategically focus on unilateral movements (such as Bulgarian split squats and lunges) and moderate-weight leg presses. These movements stimulate the quadriceps and hamstrings sufficiently to maintain proportion without adding excessive bulk to the hips and glutes.

5.3 The Kinematics of Lateral Raises: Cables versus Dumbbells

Given the absolute, non-negotiable necessity of lateral deltoid hypertrophy for establishing the V-taper, a deep analysis of exercise selection is required. Lifters frequently debate the superiority of the dumbbell lateral raise versus the cable lateral raise. Biomechanically, these two implements present entirely opposite resistance profiles to the shoulder joint.

Dumbbells rely exclusively on gravity, which provides a strictly vertical line of force. In a standing dumbbell lateral raise, the moment arm—defined as the perpendicular distance from the axis of rotation (the shoulder joint) to the line of force—is practically zero at the bottom of the movement when the arms hang by the sides. Therefore, there is zero mechanical tension on the deltoid when the muscle is in its fully stretched, lengthened position. Tension increases progressively as the arm abducts, peaking at the top of the movement (90 degrees of abduction), resulting in an *ascending resistance profile*.

Conversely, a cable machine artificially redirects the line of force via a pulley mechanism. By adjusting the pulley to a low position and standing slightly away from the stack, the moment arm is maximized near the bottom of the movement. This places the lateral deltoid under immense mechanical tension while it is in a fully stretched position. As the arm raises, the angle changes and the tension remains constant or slightly decreases, resulting in a *descending or constant resistance profile*.

Current physiological literature heavily suggests that applying mechanical tension to a muscle while it is at long muscle lengths (the stretched position) may act as a powerful, independent driver of hypertrophy, potentially making cables superior. However, recent experimental data, including an 8-week ultrasound study comparing the two modalities in resistance-trained individuals, demonstrated that both cables and dumbbells produce statistically identical overall growth in the lateral deltoid, with thickness increasing by roughly 3.3% to 4.6% in both groups. The most pragmatic conclusion is that utilizing *both* modalities within a comprehensive training block ensures the lateral deltoid is stimulated across its entire contractile range, exploiting both ascending and descending resistance profiles to maximize the architectural development of the

shoulder.

6. The 4-Day Aesthetic V-Taper Training Protocol

Based on the biomechanical requirements detailed above, and the urgent medical need to mitigate the subject's current state of central nervous system overtraining, the following four-day training split has been meticulously formulated. It provides an optimal balance of training frequency, mechanical volume, and essential biological recovery, categorizing movements into Push, Pull, Legs, and a highly specialized V-Taper Hypertrophy day.

6.1 Progressive Training Guidelines

To ensure continuous muscular adaptation, the subject must adhere to the principle of progressive overload, striving to increase either the external resistance (weight) or the total repetitions performed in each subsequent microcycle. Sets should be taken to a Rate of Perceived Exertion (RPE) of 8 or 9, meaning the lifter halts the set when only 1 to 2 repetitions remain in reserve before catastrophic muscular failure occurs.

Rest intervals must be strictly monitored; allow 90 to 120 seconds of rest between heavy compound movements, and 60 to 90 seconds for isolation movements to ensure full ATP resynthesis within the muscle cells. Furthermore, execution is paramount. Controlled eccentric (lowering) phases lasting 2 to 3 seconds are mandated for every repetition to drastically increase time-under-tension, which serves as an independent variable for stimulating muscle growth.

6.2 Workout Architecture

Day 1: Upper Body Push (Chest, Anterior/Lateral Delts, Triceps)

This session prioritizes the mechanical loading of the clavicular head of the pectoralis and the anterior deltoids, laying the foundational frontal thickness of the upper torso.

Exercise Modality	Target Musculature	Sets	Repetitions	Rest Interval
Incline Dumbbell Press	Clavicular Pectoralis Major	4	8 - 10	90 seconds
Seated Dumbbell Shoulder Press	Anterior Deltoid	3	8 - 12	90 seconds
Flat Machine Chest Press	Sternal Pectoralis Major	3	10 - 12	90 seconds
Dumbbell Lateral Raise	Lateral Deltoid	4	12 - 15	60 seconds
Triceps Rope Pushdown	Triceps Brachii	3	10 - 15	60 seconds
Overhead Triceps Extension	Triceps (Long Head Focus)	3	10 - 15	60 seconds

Day 2: Upper Body Pull (Lats, Trapezius, Biceps)

This session is strictly focused on vertical and horizontal pulling kinematics to expand the

latissimus dorsi, creating the structural width necessary for the V-taper illusion.

Exercise Modality	Target Musculature	Sets	Repetitions	Rest Interval
Wide-Grip Pull-Ups (Or Assisted)	Latissimus Dorsi	4	6 - 10	120 seconds
Close-Grip Lat Pulldown	Lower Lats / Teres Major	3	8 - 12	90 seconds
Chest-Supported Dumbbell Row	Rhomboids / Trapezius	3	10 - 12	90 seconds
Cable Face Pulls	Posterior Deltoid	3	12 - 15	60 seconds
EZ-Bar Bicep Curls	Biceps Brachii	3	8 - 12	60 seconds
Incline Dumbbell Curls	Biceps (Long Head Focus)	3	10 - 12	60 seconds

Day 3: Lower Body and Core Stabilization

Lower body training is carefully calibrated to stimulate hypertrophy for functional strength and systemic hormonal release without drastically expanding the circumference of the glutes or waistline.

Exercise Modality	Target Musculature	Sets	Repetitions	Rest Interval
45-Degree Leg Press	Quadriceps Femoris	4	8 - 12	120 seconds
Romanian Deadlift (Dumbbell)	Hamstrings / Gluteus	3	8 - 10	120 seconds
Bulgarian Split Squats	Unilateral Quadriceps	3	10 - 12 (per leg)	90 seconds
Seated Leg Curl Machine	Hamstrings	3	12 - 15	60 seconds
Standing Calf Raises	Gastrocnemius	4	15 - 20	60 seconds
Forearm Plank	Transverse Abdominis	3	45 - 60 seconds	60 seconds

Day 4: V-Taper Specific Hypertrophy (Width Emphasis)

The final training day of the microcycle is an accessory session engineered to flood the lateral deltoids and lats with metabolic stress. This drives localized hypertrophy without taxing the central nervous system with heavy compound movements, exploiting the benefits of both cable and machine resistance profiles.

Exercise Modality	Target Musculature	Sets	Repetitions	Rest Interval
Single-Arm Cable Lateral Raise	Lateral Deltoid	4	12 - 15	60 seconds
Wide-Grip Lat Pulldown	Latissimus Dorsi	4	10 - 12	90 seconds

Exercise Modality	Target Musculature	Sets	Repetitions	Rest Interval
Upright Rows (Cable or Barbell)	Lateral Delt / Trapezius	3	10 - 12	90 seconds
Straight-Arm Lat Pulldown	Latissimus Dorsi	3	12 - 15	60 seconds
Machine Rear Delt Fly	Posterior Deltoid	3	12 - 15	60 seconds
Hanging Leg Raises	Rectus Abdominis	3	10 - 15	60 seconds

7. Conclusions and Systemic Integration

The subject's historical transition from a clinically underweight state to a current regimen of chronic, seven-day-per-week overtraining has resulted in severe physiological stagnation, systemic fatigue, and a presentation of the skinny-fat somatotype. To achieve the specific aesthetic, V-taper physique desired—characterized by broad shoulders, a wide back, and a narrow waistline, absent of bulky, blocky muscle mass—a fundamental paradigm shift toward biological recovery and precise macronutrient partitioning is required.

The comprehensive nutritional protocol resolves the complex economic and logistical constraints of student hostel living by leveraging highly bioavailable, low-cost plant and animal proteins. The strategic implementation of sattu, soya chunks, eggs, and dairy, all of which require minimal to zero preparation via an electric kettle, ensures a consistent, calculated daily caloric intake of roughly 2,050 kilocalories and a protein threshold exceeding 120 grams. When the subject returns to the home environment, the integration of traditional, mother-cooked Bengali staples—such as freshwater fish, lentil soups, and rice—maintains this anabolic trajectory through rigorous, pre-calculated portion control, specifically mitigating the caloric dangers of excess culinary oils and simple carbohydrates.

Simultaneously, the training protocol eliminates the central nervous system burnout associated with daily concurrent training by scaling the regimen back to an evidence-based four-day split. This routine hyper-focuses on the exact biomechanical pathways that dictate visual aesthetics: emphasizing lateral deltoid and latissimus dorsi expansion by utilizing both constant (cables) and ascending (dumbbells) tension curves. It purposefully minimizes excessive lower body and oblique mass development to preserve a tight, narrow waistline. Strict adherence to this dual-environment nutritional strategy and targeted biomechanical training protocol will initiate optimal body recomposition, allowing the subject to shed localized adipose tissue while synthesizing highly specific, aesthetically pleasing muscle mass.

This is for informational purposes only. For medical advice or diagnosis, consult a professional.

Works cited

1. Lose Fat, Build Muscle: Skinny Fat Men India Guide - Apollo Pharmacy, <https://www.apollopharmacy.in/blogs/article/fix-skinny-fat-for-indian-men-build-muscle-lose-belly-with-smart-nutrition>
2. How To Reduce Skinny Fat? - Apollo 247, <https://www.apollo247.com/health-topics/diet-for-weight-loss/how-to-reduce-skinny-fat>
3. Best Meal Plan for Skinny Fat: Burn Fat & Build Lean Muscle - Nuceria Health, <https://mynuceria.com/blog/best-meal-plan-for-skinny-fat-burn-fat-build-lean-muscle>
4. 3-Month Body Recomposition Guide | PDF - Scribd,

<https://www.scribd.com/document/913457032/Body-Recomposition-Plan-Abhilash> 5. 4 Day Workout Split for Strength & Muscle | TruFit Athletic Clubs, <https://trufitathleticclubs.com/blog/4-day-workout-split/> 6. The Ultimate 4 Day Workout Split Guide - VASA Fitness, <https://vasafitness.com/blog/the-ultimate-4-day-workout-split/> 7. Best V-Taper Workout To Sculpt Your Body The Right Way - Endomondo, <https://www.endomondo.com/workouts/v-taper-workout> 8. Split Workout Schedule: What To Know and Examples - Healthline, <https://www.healthline.com/health/fitness/split-workout-schedule> 9. Your Ultimate Guide to the 4 Day Split Workout | Zing Coach, <https://www.zing.coach/fitness-library/4-day-split> 10. 4-Day Workout Splits to Build Muscle & Strength | EoS Fitness - EoS Fitness, <https://www.eosfitness.com/blog/winter-4-day-splits> 11. Dietician Advice for Students Struggling With Hostel Food - Tulip Multispeciality Hospital, <https://tuliphospital.com/dietician-advice-for-students-struggling-with-hostel-food/> 12. 7 Electric Kettle Recipes Every Hostel Students Should Try . - Wipro Appliances, <https://wiproappliances.com/blogs/recipes/electric-kettle-recipes> 13. Very simple veg recipes for hostel (almost no cooking) - Reddit, https://www.reddit.com/r/Cooking/comments/1sm0q3c/very_simple_veg_recipes_for_hostel_almost_no/ 14. Hostel Mess Guide: Lose Fat on a Student Budget - Apollo Pharmacy, <https://www.apollopharmacy.in/blogs/article/hostel-mess-diet-plan-india> 15. Electric Kettle Recipes: 20 Quick Meals for Hostel, PG & Office - InstaCuppa Store, <https://instacuppastore.com/blogs/main-page/electric-kettle-recipes-20-quick-meals-hostel-pg-office> 16. Cheap Protein Sources in India: High Protein Foods on Any Budget (2026), <https://gourmetstaples.com/blogs/news/cheap-protein-sources-india> 17. Low Budget Gym Diet for College Students in India (₹150/Day Plan), <https://budget-fuel-guide.vercel.app/blog/low-budget-gym-diet-for-college-students-india> 18. High-Protein Indian Meals for Students & Budget Gym-Goers - Nutristar, <https://nutristar.in/blogs/news/high-protein-indian-meals-for-students-budget-gym-goers> 19. Low Cost High Protein Foods in India: Ultimate Guide for Health, Muscle, and Budget, <https://www.nfna.in/low-cost-high-protein-foods-in-india/> 20. How to cook Soya Chunks / Meal Maker - Sharmis Passions, <https://www.sharmispassions.com/how-to-cook-soya-chunks-meal-maker/> 21. How do I dry cook Soy Chunks properly in an Electric Kettle : r/IndianFood - Reddit, https://www.reddit.com/r/IndianFood/comments/1p34r19/how_do_i_dry_cook_soy_chunks_properly_in_an/ 22. 7 Tips to Cook Soya Chunks at Home with Sparsh Masala, <https://sparshmasala.in/7-tips-to-cook-soya-chunks/> 23. What is Sattu? India's Original Protein Powder - Complete Guide (2026) - Organic Mandya, <https://organicmandya.com/blogs/news/what-is-sattu-complete-guide> 24. Sattu Ke Fayde: 10 Health Benefits of Sattu - Ecavo, <https://ecavo.in/blog/sattu-benefits> 25. Sattu: Uses, Benefits, Side Effects & More - PharmEasy, <https://pharmeasy.in/blog/sattu-uses-benefits-side-effects-more/> 26. 20 Best Protein Sources and Intakes for Vegetarians | Medanta, <https://www.medanta.org/patient-education-blog/20-power-packed-protein-sources-for-vegetarians> 27. Bengal fish curry (Bengali machli curry) - Anuvaad Solutions, <https://www.anuvaad.org.in/nutrition-fact/bengal-fish-curry-bengali-machli-curry/> 28. 15 Low-Calorie Bengali Recipes for Weight Loss (With Diet Chart) - Runnar Hut, <https://runnarhut.com/low-calorie-bengali-recipes-weight-loss-diet-chart/> 29. Comprehensive Guide to Indian Food Calories and Nutrition - Nicole Doria, <https://alwaysonmymind.blog/2025/09/25/comprehensive-guide-to-indian-food-calories-and-nutri>

tion/ 30. Taste of Bengal | PDF | Curry | Ginger - Scribd,
<https://www.scribd.com/document/482690318/Taste-of-Bengal-docx> 31. Bengali Rui Macher Jhol Jhinge, Aloo diye - Debjanir Rannaghar,
<https://kitchenofdebjani.com/2023/04/bengali-rui-macher-jhol-jhinge-aloo-diye/> 32. Recipes - Weekend Special Cooking – Satyajeet David,
<https://weekendspecialcooking.wordpress.com/recipes/> 33. Indian style meal with rice, greens, dal, and curry calories & nutrition - Arise App,
<https://www.arise-app.com/dish/Indian-style-meal-with-rice,-greens,-dal,-and-curry> 34. Calories in Bengali Fish Curry by Fed and Nutrition Facts - MyNetDiary,
<https://www.mynetdiary.com/food/calories-in-bengali-fish-curry-by-fed-serving-42073744-0.html> 35. The Ultimate V-Shaped Body Workout (3 Workouts Explained) - Fitbod,
<https://fitbod.me/blog/v-shaped-body-workout/> 36. How to Build a V-Taper Physique (Ideal Shoulder-to-Waist Ratio) - Outlift, <https://outlift.com/v-taper-and-shoulder-to-waist-ratio/> 37. Ultimate Workout to Develop a Dramatic V Taper - Muscle & Strength,
<https://www.muscleandstrength.com/workouts/how-to-build-dramatic-v-taper> 38. Building a V-Tapered Physique Guide | PDF | Limbs (Anatomy) | Weight Training - Scribd,
<https://www.scribd.com/document/916049411/v-taper-1> 39. How to develop defined v-shape? : r/Fitness - Reddit,
https://www.reddit.com/r/Fitness/comments/3zw7n4/how_to_develop_defined_vshape/ 40. Achieving an Aesthetic V-Taper Physique with a Naturally Wider Waistline and a more Torso Dominant Body? : r/naturalbodybuilding - Reddit,
https://www.reddit.com/r/naturalbodybuilding/comments/1g3ifyh/achieving_an_aesthetic_vtaper_physique_with_a/ 41. 4-Day V-Taper Workout Plan | PDF - Scribd,
<https://www.scribd.com/document/959327839/4-Day-v-Taper-Workout-Plan> 42. Cable Lateral Raise vs Dumbbell: Which Exercise Better Targets Your Shoulder Growth?,
<https://dumbbellsdirect.com/blogs/dumbbells-comparison-alternatives/cable-lateral-raise-vs-dumbbell> 43. Dumbbell Lateral Raise vs. Cable Lateral Raise: Which Is Best? - StrengthLog,
<https://www.strengthlog.com/dumbbell-lateral-raise-vs-cable-lateral-raise/> 44. Dumbbell versus cable lateral raises for lateral deltoid hypertrophy: an experimental study,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC12277279/> 45. Cable vs. Dumbbell Lateral Raise: Which Builds Bigger Delts? - Legion Athletics,
<https://legionathletics.com/cable-vs-dumbbell-lateral-raise/> 46. New study: cable vs. dumbbell lateral raises for building big side delts? Which is best?,
https://www.reddit.com/r/strengthlog/comments/1hvk5ur/new_study_cable_vs_dumbbell_lateral_raises_for/ 47. 4-Day V-Shape Body Gym Program | PDF - Scribd,
<https://www.scribd.com/document/951014654/V-Shape-4-Day-Program> 48. 4 Day Muscle Building Workout: PPL Split w/ V Taper Intensifier,
<https://www.muscleandstrength.com/workouts/4-day-ppl-and-v-taper-intensifier-workout>