Exploring the Limits of Machine-Learning Classification of Neutron-Star Matter Models:

Populations of Equations of State, Literature-Anchored Errors and Calibrated Uncertainty

Wasif Husain
School of Science, Technology and Engineering, University of Sunshine Coast, Adelaide 5000

Revised manuscript (supersedes arXiv:2512.23758v3)

Abstract

Machine-learning classification of neutron-star composition is usually demonstrated on synthetic data in which every composition is a single equation of state (EoS) and test points are random draws along the same sequences. We show first that the headline 97% accuracy of a released benchmark of 770 such models is largely interpolation between neighbouring models (the MLP accuracy falls from 97% to 94% and 69% when blocks of six and 24 models are withheld) and that its tabulated surface redshift leaks class information. We then replace the benchmark by a population study of 360 EoS, 90 in each of four composition scenarios (nucleonic, hyperonic-like, strange, and neutron-decay dark-matter-admixed), solved with a full general-relativistic TOV, tidal and f-mode code that reproduces six independent reference f-mode frequencies to 0.007%. With whole-EoS hold-out, six classifiers (logistic regression, k-NN, random forest, gradient boosting, SVM and an MLP ensemble) reach only 64% accuracy even without measurement error, and a kernel-Bayes classifier that uses the exact error model does no better: within these populations the plateau appears to be set by the information in the observables, not by the learner. Strange matter is separable (100% recall when f is measured accurately), hyperonic-like matter partly (74%), but nucleonic and dark-matter-admixed matter are indistinguishable once the nuclear EoS is unknown (pairwise accuracy 52%); with the nuclear EoS known exactly the dark component, which shifts f by about 0.56%, would need a $\sim$0.51% f-mode measurement to be detected at 75%. With errors anchored on published measurements and forecasts (NICER-like 10% in M and 9% in R, and a 20% f-mode measurement) the accuracy is 34–37% (chance 25%) and conformal prediction sets contain 3.22 of four classes; 5%, 5% and 2% errors would give 53%, and the damping time adds nothing. Probabilities are well calibrated and conformal coverage is about 90%. This is a controlled identifiability and forecasting study: all results are conditional on the phenomenological EoS families, priors and error models defined here, and no inference about real stars is claimed.

1 Introduction

Neutron stars probe matter at densities well above nuclear saturation (Lattimer & Prakash, 2004; Özel & Freire, 2016), and a wide range of compositions—purely nucleonic matter, hyperon-rich matter, deconfined quark matter and neutron decay into a dark sector (Husain et al., 2022)—has been proposed. Machine learning has been applied to neutron-star inference, for example to map observed masses and radii to the EoS (Fujimoto et al., 2018). A natural question is how far a supervised classifier can distinguish composition from observables, and where the answer is limited by physics rather than by the algorithm.

A previous version of this work (arXiv:2512.23758v3) trained a small multilayer perceptron (MLP) on seven observables of 770 synthetic stellar models, each class being one EoS, and reported a test accuracy of 97.4% on 116 held-out models. That number does not support the conclusions drawn from it, and a single sequence per class cannot say how well compositions can be told apart when the EoS itself is uncertain. This revision therefore does the following.

Table 1: Composition of the released single-sequence data set (770 models). $N_{\geq 0.5}$ and $N_{\geq 1}$ are the numbers of models with $M \geq 0.5\,M_\odot$ and $M \geq 1\,M_\odot$.

Class	N	M range ($M_\odot$)	R range (km)	$M < 0.5\,M_\odot$	$N_{\geq 0.5}$	$N_{\geq 1}$
Nucleonic	222	7.2×10^{-8}–2.16	0.06–11.11	47%	118	83
Strange	220	4.2×10^{-6}–1.94	0.11–11.06	45%	122	80
Dark-matter	122	1.7×10^{-3}–0.68	0.51–6.56	67%	40	0
Hyperonic	206	2.0×10^{-4}–1.74	0.31–10.56	44%	115	69




Figure 1: (a) Mass–radius sequences of the four classes. (b) Distribution of masses; the dotted line is $1\,M_\odot$. (c) Fractional difference between the tabulated surface redshift and the value implied by M and R; the hyperonic and dark-matter classes agree to $\sim$0.03%, the nucleonic and strange classes do not.

1. **Audits the single-sequence benchmark** (Sec. 2): class sizes, mass coverage, a class-dependent redshift inconsistency, and the dependence of the accuracy on how test points are chosen.

2. **Builds a population of EoS** for each composition scenario and solves every one with a validated full-GR code (Sec. 3); the validation against six independent f-mode frequencies is given in Table 4.

3. **Evaluates with whole-EoS hold-out**, six learners and a kernel-Bayes reference classifier (Secs. 4, 5), under error scenarios taken from published measurements and forecasts as well as a scale that shows what precision would be needed.

4. **Quantifies uncertainty**: calibration, temperature scaling and conformal prediction sets with exchangeability at the level of whole EoS.

Appendix A keeps the single-sequence noise study, and Appendix C lists each claim of the earlier version that has been withdrawn or changed.

2 What the single-sequence benchmark can and cannot show

2.1 Audit of the released data set

The released data (`DataML.xlsx`) hold 770 models of four classes: a nucleonic and a hyperonic quark–meson-coupling EoS (Rikovska Stone et al., 2007), a neutron-decay dark-matter scenario (Husain et al., 2022) and MIT-bag strange matter; the oscillation quantities come from the earlier oscillation code of Husain (2026a) and are used as released here. Table 1 and Fig. 1 summarise the data.

1. *Class sizes.* The classes contain 222, 220, 122 and 206 models (not $\approx$ 190 each).

Table 2: Random-split test accuracy (%) over 100 seeds (stratified 70/15/15, $N_{\rm test} = 116$), noise-free.

Setup	Mean	s.d.	Min–max
Published setup: 7 columns incl. z, MLP	97.7	1.6	92.2–100.0
Published setup, random forest	99.2	1.1	94.0–100.0
Redshift removed, MLP	97.5	1.9	90.5–100.0
Redshift removed, random forest	98.9	1.2	94.0–100.0
Redshift removed, k-NN	98.7	1.0	96.6–100.0
M, R, f, τ only, MLP	93.6	3.5	79.3–99.1
M, R only, k-NN	60.1	4.1	50.9–69.8
M, R only, logistic regression	27.1	4.0	19.0–41.4
M only, logistic regression	25.3	3.4	15.5–31.9




Figure 2: (a) Test accuracy over 100 random splits for several setups (star: seed 42, used in the earlier version). (b) Distribution for the original setup.

2. *Sampling.* Models lie on a regular radius grid ($\Delta R = 0.05\,$km) along each sequence, so neighbouring models are almost identical: for 96.5% of the models the nearest neighbour in standardised $(\ln M, \ln R, \ln f, \ln \tau)$ belongs to the same class.

3. *Mass coverage.* 49% of the models have $M < 0.5\,M_\odot$ and 70% have $M < 1\,M_\odot$, down to $7.2 \times 10^{-8}\,M_\odot$.

4. *The dark-matter sequence ends at* $0.68\,M_\odot$, so at $M \gtrsim 0.7\,M_\odot$ the class is excluded by the mass alone.

5. *Redshift inconsistency.* For the dark-matter and hyperonic classes the tabulated z equals the Schwarzschild value implied by (M, R) to ≈0.031%; for the nucleonic and strange classes the median deviation is +18% and +11%. A logistic regression on the redshift residual alone separates {nucleonic, strange} from {dark matter, hyperonic} with 91% cross-validated accuracy (57% for M alone, 72% for M and R). The redshift is therefore a class-dependent artefact and is not used in any analysis here.

2.2 Replicating the earlier protocol

Table 2 and Fig. 2 repeat the earlier random-split protocol for 100 seeds. The published 97.4% is typical ($97.7 \pm 1.6\%$), so it is reproducible, but it is not specific to the MLP: a random forest reaches 99.2%. Removing the redshift changes the MLP result only from 97.7% to 97.5%. With M and R only, a k-nearest-neighbour classifier reaches 60.1%, and with M alone logistic regression is at chance (25.3%).




Figure 3: Noise-free test accuracy versus the length of the held-out contiguous mass block (mean and 95% t-interval over 15 repetitions). Dotted: chance. A block length of one is equivalent to the random split of the earlier version.

Table 3: Parametrisations and priors. PP: piecewise polytrope (Read et al., 2009) with core break densities $\rho_1 = 10^{14.7}$ and $\rho_2 = 10^{15}\,\mathrm{g\,cm^{-3}}$ on a four-piece crust; CSS: constant speed of sound, $\varepsilon = \varepsilon_0 + p/c_s^2$.

Scenario	Parameters (uniform priors)
Nucleonic	PP: $\log_{10} P_1 \in [34.15, 34.60]$ ($\mathrm{dyn\,cm^{-2}}$ at ρ_1), $\Gamma_1 \in [2.4, 3.8]$, $\Gamma_2 \in [2.4, 3.9]$, $\Gamma_3 \in [2.2, 3.9]$
Hyperonic-like	PP with $\log_{10} P_1 \in [34.20, 34.60]$, $\Gamma_1 \in [2.6, 3.8]$ and a soft segment $\Gamma_2 \in [1.8, 2.6]$ above $\rho_1 \approx 2n_0$ (hyperon onset), then a causal CSS segment with $c_s^2 \in [0.5, 1]$ above $\rho_t = 10^u$, $u \in [14.95, 15.25]$
Strange (self-bound)	CSS with $\varepsilon_0 = 4B$, $B^{1/4} \in [130, 160]\,\mathrm{MeV}$, $c_s^2 \in [1/3, 1]$ ($c_s^2 = 1/3$ is the MIT bag model)
Dark-matter admixed	nucleonic PP prior plus a self-interacting dark Fermi gas in chemical equilibrium with the baryons (neutron-decay scenario): $m_\chi \in [937.9, 939.2]\,\mathrm{MeV}$, repulsive self-interaction $G \in [50, 500]\,\mathrm{fm^2}$; effective one-fluid barotrope

2.3 Leakage from neighbouring models

Figure 3 holds out contiguous blocks of increasing length (noise-free, 70/30 split, 15 repetitions). With all six observables (M, R, f, Q, τ, h_c) the MLP accuracy decreases from 97% (single-model holdout, equivalent to a random split) to 94% at six-model blocks and 69% at 24; the random forest drops from 99% to 82% and 42%. With the four plausible observables M, R, f, τ the MLP values are 78%, 69% and 45%. A large part of the apparent skill is therefore interpolation between near-duplicate neighbours of the same sequence. Q and h_c are model-derived amplitudes rather than measurable quantities and are not used further. Even with the leakage removed, a single sequence per class cannot say whether the *composition* is identifiable when the EoS is uncertain: "classifying the family" and "recognising one EoS" are the same task. The remainder of the paper addresses this with populations.

3 A population of equations of state

3.1 Composition scenarios as EoS populations

Each scenario is a family of EoS defined by a small number of parameters with a uniform prior (Table 3); one draw from the prior is one EoS. The families are phenomenological parametrisations *inspired by* the four scenarios of the earlier version, not microphysical calculations. Draws are accepted only if they satisfy broad observational and consistency screens: $M_{\max} \geq 1.97\,M_\odot$, a causal speed of sound ($c_s^2 \leq 1$) up to the centre of the maximum-mass star, $70 \leq \Lambda_{1.4} \leq 800$

and $10 \leq R_{1.4} \leq 14\,\mathrm{km}$. The acceptance rates are 21%, 57%, 24% and 26% for the nucleonic, strange, dark-matter and hyperonic-like priors, so the screens remove most of the prior volume (mainly through $M_{\max}$ and causality, and for strange matter through $\Lambda_{1.4}$ and $R_{1.4}$). Observables are tabulated for stars of 1.2–1.9 $M_\odot$ and evaluated on a mass grid over 1.2–1.85 $M_\odot$, where all accepted EoS have stable stars.

The dark-matter-admixed scenario follows the effective-barotrope construction of the earlier work: baryons (nucleonic PP, $\mu_N = (\varepsilon+p)/n_B$) and a dark Fermi gas with $\mu_\chi = \sqrt{m_\chi^2 + k_F^2} + \hbar c G n_\chi$ share one chemical potential, and the total pressure and energy density are summed at that potential. Because the dark component appears only above $\mu_N = m_\chi$, the stellar surface is purely baryonic and the dark component is at most a few per cent of the energy density even at the highest densities. By construction this scenario shares the nuclear prior with the nucleonic one, so the question the classifier faces is whether the dark component can be recognised against the spread of nuclear EoS.

3.2 Solvers and validation

TOV and tidal deformability. Stars are integrated with the Tolman–Oppenheimer–Volkoff equations (Tolman, 1939; Oppenheimer & Volkoff, 1939) and the tidal-deformability equation (Hinderer, 2008).

Full-GR f-mode. The $l = 2$ quasi-normal f-mode of the non-rotating, barotropic star is computed from the linearised Einstein equations in the Regge–Wheeler gauge. The interior equations were derived symbolically (not taken from the literature) from the linearised Einstein equations and the conservation law, reduced to four first-order equations, and solved with regular boundary conditions at the centre and a vanishing Lagrangian pressure perturbation at the surface; this is the same physical system as the Lindblom–Detweiler formulation (Detweiler & Lindblom, 1985). The exterior solution is matched to the Zerilli equation (Zerilli, 1970), and the complex eigenfrequency is the zero of the ingoing-wave amplitude; the real part gives f and the imaginary part τ. The exterior equations were checked against an independent symbolic derivation (residual 10^{-12}). The complex eigenfrequency is found by a secant search on the analytic ingoing amplitude, starting within 3% of the root.

Numerical error budget. Varying one numerical setting at a time (interior and exterior integration tolerances from 10^{-8} to 10^{-13}, the TOV tolerance, the surface cut-off, the starting radius at the centre and the outer matching radius from 1500 to 8000 km) changes f by at most 3e-08 in relative terms and τ by at most 5e-07, for SLy and for SLy with $G = 100\,\mathrm{fm}^2$ at $1.4\,M_\odot$. These are about seven orders of magnitude smaller than the dark-matter shift of interest (median 0.56%). The remaining sensitivity is physical: changing the crust pressure scale by ±10% shifts f of the nuclear EoS by 0.20% but changes the dark-matter shift only from 1.313% to 1.338% (SLy, $G = 100\,\mathrm{fm}^2$), so the *shift* is robust to the crust treatment even though absolute frequencies are not.

Validation. The crust is a single calibrated parameter (the absolute pressure scale of a four-piece SLy-like crust, 1.135 times the electron-gas value), fitted only to the radius and tidal deformability of six published EoS fits at $1.4\,M_\odot$ (Table 4); the radii are then reproduced to 0.0006 km and Λ to 0.05%. The f-mode frequencies were *not* used in any fit, and are reproduced to 0.007% (maximum over the six EoS). The reference values are those of the finite-EoS pilot of the earlier work (Husain 2026b), which used the separately developed RelModPy full-GR code (`https://github.com/fgittins/RelModPy`); the six values are listed in Table 4 so that the comparison can be repeated with any other code. The pilot manuscript is unpublished, so this is a comparison with an independent code rather than with published benchmark tables. In addition, the dark-matter frequency shift at fixed nuclear EoS (SLy, $G = 100\,\mathrm{fm}^2$, $1.4\,M_\odot$) is 25.7 Hz against 25.8 Hz in the reference, and a tabulated version of the same EoS agrees with the analytic one to

Table 4: Validation at $1.4\,M_\odot$ against six published piecewise-polytropic EoS fits. R and Λ were used to calibrate one crust parameter; f was held out.

EoS	R (km)	$R_{\rm ref}$	Λ	$\Lambda_{\rm ref}$	f (Hz)	$f_{\rm ref}$	$\Delta f/f$ (%)	τ (s)
SLy	11.7055	11.7061	295.5	295.6	1935.20	1935.06	+0.0072	0.1920
AP3	11.9712	11.9717	367.1	367.2	1848.51	1848.40	+0.0059	0.2094
AP4	11.3192	11.3197	247.5	247.6	2007.63	2007.50	+0.0062	0.1793
ENG	11.9695	11.9699	369.9	370.1	1845.65	1845.53	+0.0066	0.2100
MPA1	12.4339	12.4343	478.9	479.1	1745.68	1745.58	+0.0057	0.2341
WFF2	11.1245	11.1249	224.9	225.0	2047.47	2047.34	+0.0062	0.1729

Table 5: Population data set (90 EoS per scenario after screening): ranges of $M_{\rm max}$ ($M_\odot$), $R_{1.4}$ (km), $\Lambda_{1.4}$ and f at $1.4\,M_\odot$ (Hz).

Scenario	N	$M_{\rm max}$	$R_{1.4}$	$\Lambda_{1.4}$	$f_{1.4}$
Nucleonic	90	1.97–2.59	10.9–13.8	218–682	1614–2061
Strange	90	2.02–3.34	10.0–11.7	265–797	1558–1979
Dark-matter	90	1.97–2.66	10.8–13.7	199–685	1613–2098
Hyperonic	90	1.97–2.30	10.1–13.5	104–663	1625–2380

0.013% in f. Damping times of 0.17–0.23 s are obtained for the six EoS, as expected for a $1.4\,M_\odot$ f-mode; no independent reference values for τ were available.

3.3 The data set

Table 5 and Fig. 4 summarise the 360 EoS (2886 computed modes in total; about 202 s of computation per EoS). The strange family lies on a distinct f–R relation, whereas the nucleonic, dark-matter and hyperonic-like families overlap heavily; the hyperonic-like family extends to higher frequencies.

4 Methods

4.1 Whole-EoS validation

Every EoS belongs to exactly one of five folds (stratified by scenario). In each of five rounds one fold is the test set, the next fold is the calibration set and the other three are the training set; no EoS appears in more than one role, so every reported number is for EoS the classifier has never seen. The partition is repeated for three random seeds; we report the mean over folds within a repetition and Student-t 95% intervals over the three repetitions. Observations are (M, R, f, τ) at masses drawn uniformly over 1.2–1.85 $M_\odot$ (a mass prior shared by all classes), with independent noise draws, and classes are balanced (chance 25%).

4.2 Observational-error scenarios

Observables are perturbed multiplicatively, $x_j^{\rm obs} = x_j(1 + \sigma_j \epsilon_j)$, $\epsilon_j \sim \mathcal{N}(0,1)$ independent. *Literature-anchored scenarios:* **NICER-like**: $\sigma_M = 10\%$, $\sigma_R = 9\%$, no f or τ; these are the 68% uncertainties of the NICER analyses of PSR J0030+0451 ($M = 1.34^{+0.15}_{-0.16}$ and $1.44^{+0.15}_{-0.14}\,M_\odot$, $R = 12.71^{+1.14}_{-1.19}$ and $13.02^{+1.24}_{-1.06}$ km; Riley et al. 2019; Miller et al. 2019). **NICER-like plus f-mode**: in addition $\sigma_f = 20\%$, motivated by the forecast that third-generation detectors could measure the f-mode frequency of a GW170817-like inspiral to within a few hundred Hz (Williams et al., 2022), i.e. roughly 10–25% of f. *Precision scale:* to show what would be required, the same observables are also evaluated with errors $s \times (5\%, 5\%, 2\%, 20\%)$ in (M, R, f, τ) for $s \in \{0.25, 0.5, 1, 2, 4\}$, and in the noise-free limit. These values are not forecasts for any instrument; they define a scale on




Figure 4: Population data set. (a) Radius and (b) f-mode frequency versus mass; lines are medians and bands the 5–95% range across EoS of each scenario. (c) f versus R at $1.5\,M_\odot$, one point per EoS.

which the literature-anchored scenarios can be placed. Three observable sets are used: $\{M, R\}$, $\{M, R, f\}$ and $\{M, R, f, \tau\}$.

4.3 Classifiers

Learners: logistic regression, k-nearest neighbours ($k = 25$), a random forest (200 trees, minimum leaf 5), histogram gradient boosting, an RBF support-vector machine (8000 training samples) and an MLP deep ensemble (Lakshminarayanan et al., 2017) of three 2×64 ReLU networks (the architecture of the earlier version) on log-transformed, standardised inputs. Training data contain three noise realisations of 33 masses for each training EoS.

4.4 Kernel-Bayes reference classifier

For a reference set of (EoS, M) points the posterior of class k for an observation $\mathbf{x}^{\mathrm{obs}}$ is

$$P(k \mid \mathbf{x}^{\mathrm{obs}}) \;\propto\; \frac{1}{N_k} \sum_{i \in k} \prod_{j} \mathcal{N}\Big(x_j^{\mathrm{obs}};\, x_{ij},\, (\sigma_j x_{ij})^2\Big)\,, \tag{1}$$

which is the Bayes-optimal classifier for the stated error model and a uniform prior over the reference EoS and masses. We use the training EoS as reference (the same information as the learners) and, separately, all EoS except the test fold (a larger reference). Unlike the single-sequence oracle of Appendix A, this is not an upper bound, because the test EoS are not in the reference set; agreement between learners and kernel-Bayes indicates that the remaining error is set by the information in the observables rather than by the learner.

4.5 Uncertainty quantification

Calibration: expected calibration error (ECE, 15 bins), negative log-likelihood (NLL) and Brier score, before and after temperature scaling fitted on the calibration EoS. *Epistemic uncertainty*: mutual information across ensemble members. *Conformal prediction*: split conformal prediction with the score $1 - \hat{p}(y \mid x)$ at target miscoverage $\alpha = 0.1$ (Vovk et al., 2005; Angelopoulos & Bates, 2023), calibrated on the calibration EoS. The exchangeable unit is the EoS: the scores of all stars of a calibration EoS (33 masses, two noise draws each) are pooled, the finite-sample quantile level $\lceil (n+1)(1-\alpha) \rceil / n$ uses n equal to the number of calibration EoS (about 72), and the target is the marginal coverage of a new star (EoS, mass, noise draw) from the same population. Stars of one EoS are not independent, so the coverage guarantee is approximate; it is verified empirically on held-out EoS. It is marginal, not conditional on class or mass.




Figure 5: Held-out-EoS accuracy without measurement error for each learner and observable set (95% t-intervals over three repetitions; dotted: chance).

5 Results

5.1 Noise-free separability of EoS populations

Figure 5 shows the held-out-EoS accuracy without measurement error, and the left panel of Fig. 7 the confusion matrix. With M and R only the accuracy is 39% (MLP ensemble; logistic regression 36%, random forest 39%); adding the f-mode raises it to 64% and adding τ changes nothing (64%). All learners agree within a few points. The confusion matrix shows why the ceiling is so low: strange matter is recognised perfectly (100%), hyperonic-like matter in 74% of cases, but nucleonic and dark-matter-admixed stars are assigned to each other almost at random (recall 44% and 40%; accuracy between the two classes 52%, chance 50%). Merging them into one class gives 84% three-class accuracy; the baseline for this merged problem is 50% (always predicting the merged class, which holds half of the stars), not the 33% of three balanced classes. The reason is physical: the dark component shifts f by only 0.56% at fixed nuclear EoS (median over the population; 25.7 Hz or 1.3% for SLy with $G = 100\,\mathrm{fm}^2$; Sec. 5.5), far less than the spread of f at fixed mass produced by the uncertain nuclear EoS (about ±10%; Table 5).

5.2 Errors anchored on published measurements

Table 6 gives the accuracy under the literature-anchored scenarios and at selected points of the precision scale. With NICER-like errors (M 10%, R 9%) the accuracy is 34% (MLP ensemble; kernel-Bayes 34%), barely above the 25% chance level; adding a 20% f-mode measurement raises it to 37%. The three-class accuracy after merging nucleonic and dark-matter-admixed matter is 51%, no better than the 50% majority-class baseline of the merged problem. Errors of 5%, 5% and 2% in (M, R, f) would give 53%, and 1.3%, 1.3%, 0.5% would give 61%, which approaches but never exceeds the noise-free ceiling of 64%.

5.3 Precision scale and which observable matters

Figure 6 shows the accuracy against the precision scale. For the MLP ensemble, $\{M, R\}$ gives 39%, 37% and 29% at $s = 0.25$, 1 and 4, whereas $\{M, R, f\}$ gives 61%, 53% and 38%; the damping time with a 20% error (scaled with s) adds nothing within the uncertainty. Learners and kernel-Bayes agree to a few points at every scale (at $s = 1$: MLP 53%, kernel-Bayes 53%, kernel-Bayes with all other EoS as reference 53%; logistic regression 51%, random forest 52%, gradient boosting 53%, SVM 53%), so none of the learners tried leaves a measurable amount of information unused, and

Table 6: Held-out-EoS accuracy (%, mean ± 95% t-interval over three repetitions) by error scenario and learner. Chance is 25%.

Scenario	LogReg	k-NN	RF	HistGB	SVM	MLP ens.	Kernel Bayes
NICER-like: M, R at 10%, 9%	33±1	32±1	31±1	33±1	33±1	34±1	34±1
+ f-mode at 20%	36±1	35±0	36±1	37±1	37±0	37±0	37±1
$s = 1$: 5%, 5%, 2%	51±1	52±1	53±1	53±2	53±2	53±1	53±2
$s = 1$ with τ 20%	51±1	52±0	52±1	53±1	53±2	53±1	53±2
$s = 0.25$ (1.3%, 1.3%, 0.5%, 5%)	59±0	60±1	60±1	60±2	60±2	61±2	60±2




Figure 6: Held-out-EoS accuracy versus precision scale s ($1 = 5\%, 5\%, 2\%, 20\%$ in M, R, f, τ) for the MLP ensemble and the kernel-Bayes classifier, for three observable sets (95% t-intervals; dotted: chance). The $\{M, R, f\}$ and $\{M, R, f, \tau\}$ curves coincide.

enlarging the reference sample from the training EoS to all other EoS does not change the result (this is evidence for an information plateau within these priors, not a proof of an upper bound).

5.4 Degeneracies

Figure 7 shows row-normalised confusion matrices. With accurate f ($s = 0.25$) strange matter is still identified perfectly (100%) and the hyperonic-like class in 63%, but the nucleonic and dark-matter rows remain mixed (40% and 40% correct, 41% and 40% assigned to each other). With a 20% f-mode error all rows are confused with all columns, strange matter being the best recovered (66%).

5.5 Dark matter when the nuclear EoS is known

To separate "the dark component is small" from "the nuclear EoS is unknown", the observables of each of the 90 dark-matter-admixed EoS were also computed for its own nuclear base EoS without the dark component, at the same masses. At fixed nuclear EoS and mass the dark component raises f by a median of 0.56% (5–95% range 0.27–2.18%; positive for 100% of the pairs) and lowers R by a median of -0.40% (-1.46 to -0.20%); the shift falls steeply with the self-interaction (Fig. 8a; correlation of $\ln G$ with $\Delta f/f$ at $1.4\,M_\odot$ is -0.95). If the observer were told the nuclear EoS exactly, the Bayes accuracy for dark matter versus no dark matter (equal priors, independent Gaussian errors, mass identical by construction) would reach 75% at $\sigma_f = 0.51\%$ when R is known to 1% (0.45% for 5%) and 90% at $\sigma_f = 0.20\%$ (Fig. 8b); at the points of the precision scale it is 75% at $s = 0.25$ (0.5% in f, 1.25% in R), 58% at $s = 1$ and 52% at $s = 4$, and with a 20% f-mode error and 9% in R it is 51%. With the nuclear EoS *unknown* the accuracy between the nucleonic and dark-matter classes is 52% even without any measurement error (Sec. 5). The degeneracy therefore comes from the uncertainty of the nuclear EoS, not from the smallness of the shift alone:




Figure 7: Row-normalised confusion matrices (%) of the temperature-scaled MLP ensemble, pooled over folds and repetitions. Left to right: noise-free, $s = 0.25$, and NICER-like errors plus a 20% f-mode measurement.




Figure 8: Dark matter at fixed nuclear EoS (90 paired EoS). (a) Fractional f-mode shift at $1.4\,M_\odot$ versus the self-interaction G. (b) Bayes accuracy for dark matter versus no dark matter, averaged over EoS and masses, as a function of the f-mode error for three radius errors (dotted: chance).

detecting the dark component needs both an f-mode measurement at the 0.5% level and a nuclear EoS known to better than the shift itself.

5.6 Calibration and prediction sets

The probabilities are well calibrated because the learners are close to the information limit (Fig. 9a): at $s = 1$ the ECE is 0.038 for the raw ensemble, 0.037 after temperature scaling and 0.039 for kernel-Bayes, with NLL 0.92 nats for all three (Table 7). Conformal sets have empirical coverage between 0.91 and 0.92 at the 0.90 target for every scenario, with the finite-sample correction applied at EoS level. The price of validity is set size: 3.29 of four classes under NICER-like errors, 3.22 when a 20% f-mode is added, 2.24 at $s = 1$ and 1.85 at $s = 0.25$. Temperature scaling and kernel-Bayes give the same sets. At realistic precision the data therefore support a short list containing most of the classes, not a single label.

6 Discussion and limitations

What the results support. (i) A high accuracy on single-sequence data does not carry over to populations: with whole-EoS hold-out the highest accuracy reached by any method for these four scenarios is about 64% even without measurement error. (ii) The plateau looks informational: six different learners (including the MLP ensemble) and a kernel-Bayes classifier built on the exact error model agree, so within these priors a better learner is unlikely to help (the kernel-Bayes

Table 7: Calibration and conformal sets (mean over three repetitions): ECE and NLL (nats) for the raw MLP ensemble, the temperature-scaled ensemble ($+T$) and kernel-Bayes, and coverage and set size (target coverage 0.90).

	ECE			NLL			MLP $+T$		Kernel Bayes	
Scenario	Raw	$+T$	KB	Raw	$+T$	KB	Cov.	Size	Cov.	Size
NICER-like	0.021	0.023	0.019	1.32	1.32	1.32	0.91	3.29	0.91	3.29
$+\ f$ at 20%	0.021	0.022	0.020	1.29	1.29	1.29	0.92	3.22	0.92	3.21
$s = 0.25$	0.053	0.043	0.093	0.71	0.71	0.83	0.91	1.85	0.92	1.93
$s = 0.5$	0.043	0.039	0.054	0.75	0.75	0.77	0.91	1.93	0.91	1.97
$s = 1$	0.038	0.037	0.039	0.92	0.92	0.92	0.91	2.24	0.91	2.24
$s = 2$	0.030	0.032	0.026	1.16	1.16	1.15	0.92	2.84	0.92	2.83
$s = 4$	0.019	0.026	0.018	1.29	1.29	1.28	0.92	3.26	0.92	3.25




Figure 9: (a) Reliability diagram ($s = 1$, pooled). (b) Conformal coverage and (c) mean prediction-set size versus precision scale (95% t-intervals).

classifier uses a finite reference population and is not a rigorous upper bound). (iii) With the errors of current X-ray measurements and a plausible third-generation f-mode measurement, composition is essentially unidentifiable (34–37% against 25% chance); an accurate f-mode is the observable that would help, and the damping time adds nothing. (iv) Dark-matter-admixed matter is degenerate with nucleonic matter unless the nuclear EoS is known to better than the shift in f that the dark component produces (median 0.56%, up to about 2% for weak self-interaction); with the nuclear EoS known exactly, an f-mode precision of about 0.51% would give 75% accuracy (Sec. 5.5), in line with the EoS-conditional analysis of the earlier pilot. (v) Probabilities and conformal sets are reliable and turn the degeneracy into an explicit statement about which classes remain plausible.

Limitations. *Phenomenological families.* The EoS families are parametrisations with uniform priors, screened by $M_{\max}$, causality, $\Lambda_{1.4}$ and $R_{1.4}$; conclusions depend on these priors. The nucleonic–dark-matter degeneracy in particular follows from the dark-matter scenario sharing the nuclear prior. Microphysical models (for example a population of QMC EoS) would change the priors and might make some classes easier or harder to separate. The hyperonic-like family is a soft-then-stiff parametrisation, not a hyperonic calculation, and the strange family uses a constant speed of sound; the classifier therefore separates parametrisation families, and the perfect recognition of strange matter in particular may depend on how that family is built (a self-bound surface and a constant speed of sound give a distinct f–R relation). No out-of-family test with independently constructed microphysical EoS has been made, and the sensitivity of the conclusions to other priors and functional forms has not been quantified. *Dark matter.* The dark component is treated as an effective one-fluid barotrope in chemical equilibrium; the check against the two-fluid pilot is a single number (25.7 versus 25.8 Hz). Independent displacement of the two fluids, other dark-sector models and other dark fractions are not explored. *Solver.* Non-rotating stars, $l = 2$ only,

adiabatic perturbations equal to the equilibrium barotrope (no composition gradients, entrainment or crust elasticity); the damping time is not independently validated; the crust has one calibrated parameter. *Errors.* Independent multiplicative Gaussian errors are used; the NICER and f-mode values are anchored on published results and forecasts for individual sources, not on a population of detections, and systematic theory error and correlated errors are not modelled. *Statistics and sampling.* 90 EoS per scenario and three fold partitions that reuse the same EoS, so the t-intervals reflect partition and noise variability but not the finite size of the EoS sample; EoS-level bootstrap intervals and the dependence on population size were not computed. A uniform mass prior over 1.2–1.85 $M_\odot$ is used; conformal coverage is marginal only. *Observational realism.* The errors are independent and Gaussian; real NICER posteriors are correlated and non-Gaussian, and the GW170817 and NICER posteriors themselves were not used. *Not comparable to earlier data.* The new data come from a new solver and cannot be compared point by point with the single-sequence data set.

7 Conclusions

Evaluated on populations of EoS with whole-EoS hold-out, machine-learning classification of neutron-star composition is far less decisive than the 97% single-sequence benchmark suggested: the accuracy plateau is about 64% even without measurement error, it appears to be set by the information in the observables rather than by the learner, strange matter is the only scenario that is cleanly separable (in the family defined here), and nucleonic and dark-matter-admixed matter are not distinguishable when the nuclear EoS is unknown. With errors anchored on current X-ray measurements and a forecast f-mode precision the accuracy is 34–37%, and calibrated prediction sets contain about 3.22 of four classes.

Data, code and reproducibility

The single-sequence analysis uses `audit.py`, `repro.py`, `blocksize.py`, `main.py`, `extrap.py`, `interp_check.py` and `report.py`; the population study uses the solver and generator in `population/` (`eos.py`, `tov.py`, `fmode2.py`, `families.py`, `gen_pop.py`, `gen_base.py`, `paired_analysis.py`), the derivation scripts (`derive_vac2.py`, `derive_matter.py`, `elim.py`), the experiment (`pop_experiment.py`) and the report scripts. The code and data are on the `revised-analysis` branch of the repository below (a tagged release with a DOI will be deposited at submission); the convergence tests are in `convergence_tests.py`. The 360 generated EoS and all results are included as JSON/CSV; figures, tables and every number in the text are generated from them. Fixed seeds are used throughout. The environment is Python 3.13.0 with scikit-learn 1.7.2. The released single-sequence data are at `https://github.com/wasif553/ML-NeutronStar-Classification`.

A Single-sequence noise study (supplement)

This appendix keeps the noise, oracle and uncertainty analysis performed on the single-sequence benchmark, with blocked validation. It is superseded by the population study for questions about composition, but it isolates the effect of measurement error on the original data and provides an exact oracle bound.

Mass regimes, errors, learners, oracle

Three mass regimes are used: **A**, all models ($N = 770$); **B**, $M \geq 0.5\,M_\odot$ ($N = 395$, with 40 dark-matter models); **C**, $M \geq 1\,M_\odot$ ($N = 232$, three classes). Contiguous mass blocks of six models are assigned at random to training (60%), calibration (20%) and test (20%) roles; the analysis is repeated for 10 random assignments. Observables are perturbed as $x_j(1 + s\sigma_j\epsilon_j)$ with $(\sigma_M, \sigma_R, \sigma_f, \sigma_\tau) = (5, 5, 2, 20)\%$ at $s = 1$. The learner is a three-member MLP ensemble, with




Figure 10: Single-sequence data: test accuracy versus noise scale s for the MLP ensemble (solid, 95% t-intervals over 10 repetitions) and the oracle Bayes bound (dashed). Dotted: chance.




Figure 11: Single-sequence data: (a) reliability diagram (regime B, $s = 1$), (b) NLL, (c) conformal coverage versus noise scale.

logistic regression and a random forest as baselines. The oracle Bayes classifier knows the exact sequences (including the test blocks) and the error model,

$$P(k \mid \mathbf{x}^{\mathrm{obs}}) \;\propto\; \frac{1}{N_k} \sum_{i \in k} \prod_{j} \mathcal{N}\Big(x_j^{\mathrm{obs}};\, x_{ij},\, (s\sigma_j x_{ij})^2\Big)\,, \tag{2}$$

and is an upper bound for learners trained on finite data.

Results

At $s = 1$ in regime B the ensemble reaches 47% with M, R, f, τ against an oracle ceiling of 65% (chance 25%); M and R alone give 29% (oracle 49%). In regime C (three classes) the values are 50% (ensemble) and 65% (oracle). Adding f raises the oracle from 49% to 64%; adding a 20% τ gives 65%. Strange and hyperonic matter are the most confused pair (36% of strange models labelled hyperonic and 37% the reverse). The raw ensemble is overconfident (ECE 0.188, NLL 1.13 at $s = 1$, regime B; oracle 0.040, 0.74); temperature scaling reduces them to 0.096 and 1.00. Conformal sets have about 90% marginal coverage and contain 2.35 of four classes at $s = 1$ (oracle 1.85). When the heaviest 25% of each class is withheld, the ensemble reaches 50% (oracle 80%) and the conformal sets cover only 0.81, since exchangeability fails; the ensemble's mutual information rises from 0.010 to 0.025 nats, an incomplete warning.

Table 8: Single-sequence data: test accuracy (%, mean ± 95% t-interval over 10 repetitions) by regime, observable set and noise scale s. Chance is 25% (A, B) or 33% (C).

Regime	Observables	s	MLP ensemble	Logistic reg.	Random forest	Oracle bound
A	M, R	0.25	69.4±4.0	35.9±5.4	21.3±3.5	84.4±2.7
A	M, R	1	41.8±4.2	37.1±4.5	24.8±3.5	61.3±4.2
A	M, R	4	25.4±4.2	29.1±3.7	23.1±2.5	39.6±4.1
A	M, R, f	0.25	80.2±5.1	38.7±4.9	56.0±4.9	95.5±1.1
A	M, R, f	1	67.7±4.4	38.7±5.2	56.3±3.5	81.1±2.8
A	M, R, f	4	43.9±3.8	38.3±4.3	39.5±2.7	54.8±3.3
A	M, R, f, τ	0.25	81.8±4.0	44.8±4.8	58.9±3.7	96.0±1.0
A	M, R, f, τ	1	68.9±4.0	40.3±4.3	59.6±3.8	81.9±2.8
A	M, R, f, τ	4	43.6±3.5	38.1±4.2	39.7±2.9	55.4±3.1
B	M, R	0.25	49.7±4.8	40.6±6.0	30.9±4.7	71.8±3.9
B	M, R	1	29.2±5.2	35.9±5.7	28.1±3.5	48.8±2.4
B	M, R	4	30.3±4.4	32.1±4.3	30.1±1.6	36.7±2.5
B	M, R, f	0.25	74.1±6.0	59.5±5.2	40.8±5.8	91.5±1.9
B	M, R, f	1	46.0±6.3	45.5±4.3	38.8±3.8	63.9±2.5
B	M, R, f	4	31.3±3.9	35.1±4.3	30.6±2.6	37.6±2.2
B	M, R, f, τ	0.25	76.3±5.2	63.6±3.9	43.4±5.7	92.7±1.8
B	M, R, f, τ	1	47.3±5.8	46.4±4.4	39.9±4.3	64.6±2.7
B	M, R, f, τ	4	31.9±3.8	35.2±4.1	31.6±2.1	37.8±2.2
C	M, R	0.25	61.2±7.7	62.6±7.5	37.7±8.7	78.7±4.2
C	M, R	1	34.0±5.7	40.7±7.1	33.1±5.5	49.0±4.8
C	M, R	4	32.7±3.5	33.9±6.2	33.2±2.5	37.5±5.8
C	M, R, f	0.25	70.8±5.6	69.0±6.7	44.1±10.0	89.8±1.6
C	M, R, f	1	48.9±6.8	51.4±6.6	41.9±6.7	64.0±3.5
C	M, R, f	4	34.2±4.2	36.0±5.1	34.3±3.4	40.2±4.3
C	M, R, f, τ	0.25	73.9±5.6	72.5±6.4	50.0±9.6	91.7±1.4
C	M, R, f, τ	1	49.9±5.9	52.5±7.0	44.1±6.9	65.0±3.1
C	M, R, f, τ	4	35.3±4.7	36.0±5.2	35.2±2.7	42.1±5.1

B Experimental settings

Table 9: Settings.

Item	Setting
Population folds	5 folds by EoS (stratified), test / calibration / training = 1 / 1 / 3 folds, 3 random partitions
Population samples	training: 33 masses × 3 noise draws per EoS; calibration: 33 × 2; test: 33 × 3 (single draw when noise-free)
MLP	3 members, 2×64 ReLU, Adam, lr 10^{-3}, $\alpha = 10^{-4}$, batch 128, early stopping (max 120 iterations)
Other learners	logistic regression (L2); k-NN, $k = 25$; random forest, 200 trees, min leaf 5; histogram gradient boosting (150 iterations); RBF SVM, $C = 3$, 8000 samples
Kernel Bayes	reference: 22 masses per EoS; training EoS, or all non-test EoS
Conformal	LAC score, $\alpha = 0.1$, correction with the number of calibration EoS
Single-sequence study	blocks of six models, 60/20/20, 10 repetitions; 20 noise draws per training model; random forest 150 trees; 15 repetitions for the block-length study, 100 for the random-split replication

C Changes relative to arXiv:2512.23758v3

Table 10: Statements of the earlier version that are withdrawn or changed.

Earlier statement	Status
Each class is one EoS; the classifier "separates composition"	Replaced by populations of EoS with whole-EoS hold-out; the accuracy ceiling is about 64% and nucleonic and dark-matter-admixed matter are not distinguishable.
≈ 190 models per class, 770 in total	Total is correct; classes contain 222, 220, 122 and 206 models.
Models span $\sim 1\,M_\odot$ to the maximum mass	The data span 7×10^{-8} to $2.16\,M_\odot$; 70% have $M < 1\,M_\odot$.
97.4% test accuracy	Reproducible (97.7% average over 100 splits) but inflated by neighbouring models; falls to 94–69% with six- to 24-model blocks.
Per-class accuracy ranges 82–97% "across dataset realizations"	Not reproduced; removed.
"Mild degradation near the high-mass end"	Not supported by the earlier figure; removed.
"Not merely overfitting noise"	The inputs were noise-free; replaced by explicit error scenarios.
Permutation importance shows f and τ dominate	Replaced by observable-set ablations: f carries the information, τ adds nothing.
Tabulated redshift used as an input	Inconsistent with (M, R) for two classes; removed.
Oscillation quantities from the earlier code	Population data generated with a new solver validated against six independent f-mode frequencies.

References

A. N. Angelopoulos and S. Bates, Found. Trends Mach. Learn. **16**, 494 (2023) [arXiv:2107.07511].

S. Detweiler and L. Lindblom, Astrophys. J. **292**, 12 (1985).

Y. Fujimoto, K. Fukushima, K. Murase, Phys. Rev. D **98**, 023019 (2018).

T. Hinderer, Astrophys. J. **677**, 1216 (2008).

W. Husain, S. Motta, A. W. Thomas, arXiv:2203.02758 (2022).

W. Husain, oscillation (f-mode and gravitational-wave) calculations supplying f, τ, Q, h_c of the single-sequence data set; cited as "Husain (2026)" in arXiv:2512.23758v3 (full citation to be completed by the author).

W. Husain, *Testing f-mode information beyond static structure in neutron-decay dark-matter stars: a finite-EoS pilot and working research manuscript*, 15 October 2026 (unpublished working manuscript).

B. Lakshminarayanan, A. Pritzel, C. Blundell, Adv. Neural Inf. Process. Syst. **30** (2017).

J. M. Lattimer and M. Prakash, Science **304**, 536 (2004).

M. C. Miller et al., Astrophys. J. Lett. **887**, L24 (2019).

J. R. Oppenheimer and G. M. Volkoff, Phys. Rev. **55**, 374 (1939).

F. Özel and P. Freire, Annu. Rev. Astron. Astrophys. **54**, 401 (2016).

J. S. Read, B. D. Lackey, B. J. Owen, J. L. Friedman, Phys. Rev. D **79**, 124032 (2009).

T. E. Riley et al., Astrophys. J. Lett. **887**, L21 (2019).

J. Rikovska Stone, P. A. M. Guichon, H. H. Matevosyan, A. W. Thomas, Nucl. Phys. A **792**, 341 (2007).

R. C. Tolman, Phys. Rev. **55**, 364 (1939).

V. Vovk, A. Gammerman, G. Shafer, *Algorithmic Learning in a Random World* (Springer, 2005).

N. Williams, G. Pratten, P. Schmidt, Phys. Rev. D **105**, 123032 (2022) [arXiv:2203.00623].

F. J. Zerilli, Phys. Rev. D **2**, 2141 (1970).