

AE-315: Degassing of n-Heptane

Date: 2024-07-03

Tags: Degassed AE

Status: Done

Created by: Alexander Eith

Reaction scheme/sample structure

Reagents

Name	Amount [mmol]	Equivalents	Mass _{theo} [mg]	Mass _{exp} [mg]	Molar mass [g/mol]	Volume [ml]
n-Heptane	/	/	/	/	/	approx. 400 mL

Procedure/observations

All steps (unless stated otherwise) were carried out under argon using either standard Schlenk technique or in an argon filled glovebox (MBRAUN).

Date	Time	Step	Observations
03.07	15:00	n-heptane was distilled (roti, 50 °C, settings for heptane)	
		A 500 mL Young type schlenk flask was prepared	
	15:15	Molar sieve was added to the flask (approx. 1/5)	
	15:30	n-Heptane was transferred into a prepared Schlenk Flask (not dry n-heptane)	
	15:50 - 16:25	The n-Hexane was degassed according to Protocol - Degassing of organic solvents	

Analysis

Date	Time	Sample name	Analysis method	Solvent	Raw Data	Processed Data	Interpretation

Product characterization

Linked resources

Protocol - [Degassing of organic solvents in Young-type flask](#)

Protocol - [KFT \(Karl-Fischer-Titration\)](#)

Protocol - [Collecting of SPS solvents](#)



Unique eLabID: 20240704-a3973d890f0a3140c17b00e68cad04cca2a70bed
Link: <https://elab.water-splitting.org/experiments.php?mode=view&id=1178>