
The Endoplasmic Reticulum: A Central Hub for Dermal Collagen Quality Control and Its Profound Implications for Skin Health and Disease

Anonymous Author(s)

Affiliation

Address

email

Abstract

1 The integrity and functional resilience of the skin are critically dependent on the
2 extracellular matrix (ECM), particularly its abundant collagen network. Dermal
3 collagen undergoes intricate biosynthesis within dermal fibroblasts, with the Endo-
4 plasmic Reticulum (ER) serving as the primary site for folding, assembly, and post-
5 translational modification. The ER houses a sophisticated quality control system
6 that monitors the quantity and precise quality of procollagen molecules, involv-
7 ing molecular chaperones, folding enzymes, and post-translational modification
8 machinery such as Hsp47, protein disulfide isomerase (PDI), prolyl hydroxylases,
9 and lysyl hydroxylases. This review delineates the critical ER-centric mechanisms
10 governing dermal procollagen biogenesis and quality surveillance, highlighting
11 how disruptions can lead to ER stress, activate the Unfolded Protein Response
12 (UPR), and contribute to skin conditions like aging, scarring, and fibrotic disorders.
13 By consolidating current knowledge on ER's role in dermal collagen homeostasis,
14 we aim to underscore its significance as a therapeutic target for maintaining skin
15 health and ameliorating collagen-related dermatological pathologies.

16 Keywords: Dermal collagen, Endoplasmic Reticulum, Quality Control, Hsp47,
17 Prolyl Hydroxylation, Lysyl Hydroxylation, ER Stress, Unfolded Protein Response,
18 Skin Aging, Fibrosis.

19 1 Introduction

20 The skin's robust mechanical properties, including strength, elasticity, and resilience, are largely
21 attributed to the dermal extracellular matrix (ECM), predominantly composed of collagen. Type I
22 collagen constitutes 80-90% of dermal collagen, providing tensile strength, while Type III collagen
23 (10-15%) contributes elasticity. Minor collagens, such as Type V, further refine fibril organization.
24 This complex network is essential for skin integrity, wound healing, and resisting mechanical stress.

25 Collagen biosynthesis is a complex and highly regulated process initiated within dermal fibroblasts.
26 The journey begins with the translation of pro-alpha collagen chains and their translocation into
27 the Endoplasmic Reticulum (ER), the primary organelle for its folding, assembly, and extensive
28 post-translational modifications (PTMs). The ER acts as a sophisticated, multi-layered quality control
29 system, actively determining the quantity and precise quality of each procollagen molecule. This ER-
30 centric surveillance is paramount, as the secretion of misfolded or improperly modified procollagen
31 compromises dermal matrix integrity, leading to weakened, unstable, or non-functional collagen
32 fibrils. Such deficiencies are implicated in various skin conditions, from aging and reduced elasticity
33 to impaired wound healing, keloid formation, and fibrotic disorders. This review examines the critical
34 ER-based mechanisms governing dermal procollagen biosynthesis and quality control, exploring the

35 consequences of their failure, and highlighting their implications for skin health and dermatological
36 pathologies.

37 **2 Intracellular Quality Control of Procollagen in Dermal Fibroblasts**

38 The Endoplasmic Reticulum (ER) serves as the primary and indispensable checkpoint for the folding
39 and quality control of procollagen. This rigorous system ensures that only correctly folded and
40 assembled procollagen trimers exit for further processing and deposition into the extracellular matrix.
41 Misfolded or aberrant molecules are retained, subjected to refolding cycles, and eventually degraded
42 if defects persist. This active surveillance within the ER directly determines the structural integrity
43 and biological efficacy of dermal collagen, profoundly influencing skin's biomechanical properties
44 and susceptibility to pathology. Maintaining ER quality control efficiency is paramount for dermal
45 proteostasis and skin health

46 **2.1 ER as the Primary Site for Procollagen Folding and Maturation**

47 Procollagen biosynthesis begins with mRNA translation on ER-bound ribosomes and co-translational
48 translocation of nascent polypeptide chains into the ER lumen. The ER's oxidizing environment
49 facilitates disulfide bond formation, particularly in the C-terminal propeptide domains, which guide
50 the correct association of three pro- α chains into a procollagen trimer. Simultaneously, the ER
51 provides a specialized milieu with enzymes and chaperones that catalyze and monitor extensive
52 post-translational modifications (PTMs), crucial for procollagen maturation. Successful and timely
53 execution of these events is essential; deviations lead to misfolded procollagen retention, ER stress,
54 or secretion of functionally compromised molecules, ultimately impairing dermal ECM structure and
55 function.

56 **2.2 Key Molecular Chaperones and Folding Enzymes in Dermal Fibroblasts**

57 The precise folding and assembly of dermal procollagen are meticulously orchestrated by a specialized
58 network of molecular chaperones and enzymes within the ER lumen of dermal fibroblasts. These
59 components prevent aggregation, ensure correct triple helix association, and facilitate critical post-
60 translational modifications. This system actively surveils procollagen quality, ensuring only functional
61 collagen integrates into the dermal matrix.

62 **2.2.1 Hsp47: Specific Chaperone for Dermal Procollagen Triple Helix Formation**

63 Heat Shock Protein 47 (Hsp47), or serpin H1, is a collagen-specific ER-localized molecular chaper-
64 one. Its consistent co-expression with collagen highlights its role in collagen proteostasis. Hsp47
65 specifically binds to nascent triple-helical domains, stabilizing the forming triple helix, preventing
66 incorrect folding, and safeguarding against denaturation. This activity is vital for efficient and accu-
67 rate formation of the stable triple helix. Studies show that in fibroblasts lacking Hsp47, procollagen
68 aggregates within the ER, becomes entrapped, and often fails secretion, leading to compromised
69 collagen production. This demonstrates Hsp47's direct contribution to ensuring correct formation,
70 stability, and secretion of functional collagen for dermal mechanical integrity

71 **2.2.2 Protein Disulfide Isomerase (PDI) and Other General ER Chaperones**

72 Beyond Hsp47, general ER chaperones and folding enzymes play critical roles in procollagen
73 processing. Protein Disulfide Isomerase (PDI) catalyzes disulfide bond formation and isomerization,
74 which are crucial for stabilizing the C-terminal propeptide domains that guide pro- α chain
75 trimerization. The ER's oxidative environment supports PDI's activity in forming these essential
76 linkages. Other ER chaperones, such as calnexin (CNX) and calreticulin (CRT), interact with N-
77 glycosylated pro- α chains, aiding folding and preventing aggregation. This multifaceted network,
78 including general and collagen-specific chaperones, orchestrates procollagen folding, preventing
79 premature exit of misfolded molecules and safeguarding the quality of collagen forming the skin's
80 framework

81 2.3 Critical Post-Translational Modifications and their Quality Control Implications in Skin 82 Collagen

83 The functional efficacy and structural integrity of dermal collagen depend profoundly on precise
84 enzymatic post-translational modifications (PTMs) occurring mainly in the ER lumen of dermal
85 fibroblasts. These modifications are integral to the ER's quality control system, dictating the stability,
86 correct extracellular assembly, and biomechanical properties of mature dermal collagen fibrils.

87 2.3.1 Prolyl and Lysyl Hydroxylation: Impact on Triple Helix Stability and Skin Elasticity

88 Hydroxylation of specific proline and lysine residues is a critical PTM for dermal procollagen.
89 Prolyl 4-hydroxylases (P4Hs), requiring iron, alpha-ketoglutarate, and vitamin C, convert proline to
90 4-hydroxyproline (4Hyp). 4Hyp is indispensable for stabilizing the collagen triple helix via interchain
91 hydrogen bonds. Insufficient 4Hyp leads to unstable procollagen, denaturation, aggregation, ER
92 retention, and degradation. Alternatively, secreted under-hydroxylated procollagen is functionally
93 compromised, weakening the dermal matrix and impacting skin elasticity and tensile strength.
94 Lysyl hydroxylases (LHs) convert lysine to hydroxylysine (Hyl). Hyl residues are precursors for
95 O-glycosylation and, more importantly, for stable intra- and intermolecular cross-links vital for the
96 tensile strength and resilience of mature dermal collagen fibrils. Precise prolyl and lysyl hydroxylation
97 patterns are exquisitely controlled in the ER, forming a dynamic regulatory hub. Deficiencies, as
98 seen in scurvy or genetic disorders, compromise dermal biomechanics, leading to fragile skin,
99 impaired wound healing, or reduced elasticity. The ER's quality control actively monitors these PTMs,
100 determining the structural integrity and functional prowess of the dermal collagen network.

101 2.3.2 Glycosylation of Hydroxylysine: Influence on Dermal Collagen Fibrillogenesis

102 Following hydroxylation, a subset of hydroxylysine (Hyl) residues undergoes O-glycosylation in the
103 ER, involving sequential addition of galactose and glucose by glycosyltransferases. While less critical
104 than prolyl hydroxylation for triple helix stability, Hyl glycosylation subtly modulates extracellular
105 assembly and organization of dermal collagen fibrils. These glycosylations can influence fibril
106 packing density, diameter, and overall architecture. Alterations in Hyl glycosylation patterns are
107 linked to changes in tissue biomechanics and observed in certain connective tissue disorders. The
108 ER's quality control thus regulates appropriate Hyl glycosylation, contributing to the precise structural
109 characteristics and functional properties of the mature dermal collagen network and skin health.

110 3 Conclusion

111 The Endoplasmic Reticulum serves as the indispensable central hub for the synthesis and quality
112 control of dermal procollagen, which is critical for skin integrity and resilience. This review has
113 highlighted the intricate ER-resident machinery, including Hsp47, protein disulfide isomerase, prolyl
114 hydroxylases, and lysyl hydroxylases, that collectively ensure the precise folding, assembly, and post-
115 translational modification of procollagen. These ER-centric mechanisms dictate the biomechanical
116 properties of the dermis, and their disruption is directly correlated with various skin conditions,
117 leading to ER stress, activation of the Unfolded Protein Response (UPR), and contributing to
118 pathologies like skin aging, fragility, and fibrotic disorders. This understanding of the ER's role in
119 dermal collagen quality opens promising avenues for future research and therapeutic intervention.
120 A key challenge lies in fully elucidating the precise spatiotemporal regulation of ER chaperones
121 and enzymes in diverse dermal fibroblast subtypes under varying physiological and pathological
122 conditions. Further investigation into how environmental factors (e.g., UV radiation, pollution,
123 nutrition) directly modulate ER quality control for collagen synthesis in the skin is warranted.
124 Additionally, the specific ER-associated degradation (ERAD) pathways for clearing misfolded
125 procollagen in chronic skin diseases require more thorough investigation. Future directions include
126 targeting specific ER quality control components within dermal fibroblasts to develop novel strategies
127 against collagen-related skin conditions. This could involve pharmacological approaches to enhance
128 chaperone function, modulate hydroxylase activity, or fine-tune the UPR to mitigate ER stress in
129 aging or fibrosis. Optimizing nutritional interventions, such as vitamin C delivery, could also support
130 ER-dependent collagen maturation. A more comprehensive understanding of the ER's intricate role in
131 dermal collagen homeostasis will pave the way for innovative treatments that target the fundamental
132 molecular defects underlying compromised skin health.

Agents4Science AI Involvement Checklist

This checklist is designed to allow you to explain the role of AI in your research. This is important for understanding broadly how researchers use AI and how this impacts the quality and characteristics of the research. **Do not remove the checklist! Papers not including the checklist will be desk rejected.** You will give a score for each of the categories that define the role of AI in each part of the scientific process. The scores are as follows:

- **[A] Human-generated:** Humans generated 95% or more of the research, with AI being of minimal involvement.
- **[B] Mostly human, assisted by AI:** The research was a collaboration between humans and AI models, but humans produced the majority (>50%) of the research.
- **[C] Mostly AI, assisted by human:** The research task was a collaboration between humans and AI models, but AI produced the majority (>50%) of the research.
- **[D] AI-generated:** AI performed over 95% of the research. This may involve minimal human involvement, such as prompting or high-level guidance during the research process, but the majority of the ideas and work came from the AI.

These categories leave room for interpretation, so we ask that the authors also include a brief explanation elaborating on how AI was involved in the tasks for each category. Please keep your explanation to less than 150 words.

IMPORTANT, please:

- **Delete this instruction block, but keep the section heading “Agents4Science AI Involvement Checklist”,**
- **Keep the checklist subsection headings, questions/answers and guidelines below.**
- **Do not modify the questions and only use the provided macros for your answers.**

1. **Hypothesis development:** Hypothesis development includes the process by which you came to explore this research topic and research question. This can involve the background research performed by either researchers or by AI. This can also involve whether the idea was proposed by researchers or by AI.

Answer: **[C]**

Explanation: **[TODO]**

2. **Experimental design and implementation:** This category includes design of experiments that are used to test the hypotheses, coding and implementation of computational methods, and the execution of these experiments.

Answer: **[D]**

Explanation: **[TODO]**

3. **Analysis of data and interpretation of results:** This category encompasses any process to organize and process data for the experiments in the paper. It also includes interpretations of the results of the study.

Answer: **[D]**

Explanation: **[TODO]**

4. **Writing:** This includes any processes for compiling results, methods, etc. into the final paper form. This can involve not only writing of the main text but also figure-making, improving layout of the manuscript, and formulation of narrative.

Answer: **[D]**

Explanation: **[TODO]**

5. **Observed AI Limitations:** What limitations have you found when using AI as a partner or lead author?

Description: Most of the time AI tries to give a positive answer when it's not.

Agents4Science Paper Checklist

The checklist is designed to encourage best practices for responsible machine learning research, addressing issues of reproducibility, transparency, research ethics, and societal impact. Do not remove the checklist: **Papers not including the checklist will be desk rejected.** The checklist should follow the references and follow the (optional) supplemental material. The checklist does NOT count towards the page limit.

Please read the checklist guidelines carefully for information on how to answer these questions. For each question in the checklist:

- You should answer [Yes], [No], or [NA].
- [NA] means either that the question is Not Applicable for that particular paper or the relevant information is Not Available.
- Please provide a short (1–2 sentence) justification right after your answer (even for NA).

The checklist answers are an integral part of your paper submission. They are visible to the reviewers and area chairs. You will be asked to also include it (after eventual revisions) with the final version of your paper, and its final version will be published with the paper.

The reviewers of your paper will be asked to use the checklist as one of the factors in their evaluation. While "[Yes]" is generally preferable to "[No]", it is perfectly acceptable to answer "[No]" provided a proper justification is given. In general, answering "[No]" or "[NA]" is not grounds for rejection. While the questions are phrased in a binary way, we acknowledge that the true answer is often more nuanced, so please just use your best judgment and write a justification to elaborate. All supporting evidence can appear either in the main paper or the supplemental material, provided in appendix. If you answer [Yes] to a question, in the justification please point to the section(s) where related material for the question can be found.

1. Claims

Question: Do the main claims made in the abstract and introduction accurately reflect the paper's contributions and scope?

Answer: [Yes]

Justification: [TODO]

Guidelines:

- The answer NA means that the abstract and introduction do not include the claims made in the paper.
- The abstract and/or introduction should clearly state the claims made, including the contributions made in the paper and important assumptions and limitations. A No or NA answer to this question will not be perceived well by the reviewers.
- The claims made should match theoretical and experimental results, and reflect how much the results can be expected to generalize to other settings.
- It is fine to include aspirational goals as motivation as long as it is clear that these goals are not attained by the paper.

2. Limitations

Question: Does the paper discuss the limitations of the work performed by the authors?

Answer: [Yes]

Justification: [TODO]

Guidelines:

- The answer NA means that the paper has no limitation while the answer No means that the paper has limitations, but those are not discussed in the paper.
- The authors are encouraged to create a separate "Limitations" section in their paper.
- The paper should point out any strong assumptions and how robust the results are to violations of these assumptions (e.g., independence assumptions, noiseless settings, model well-specification, asymptotic approximations only holding locally). The authors

should reflect on how these assumptions might be violated in practice and what the implications would be.

- The authors should reflect on the scope of the claims made, e.g., if the approach was only tested on a few datasets or with a few runs. In general, empirical results often depend on implicit assumptions, which should be articulated.
- The authors should reflect on the factors that influence the performance of the approach. For example, a facial recognition algorithm may perform poorly when image resolution is low or images are taken in low lighting.
- The authors should discuss the computational efficiency of the proposed algorithms and how they scale with dataset size.
- If applicable, the authors should discuss possible limitations of their approach to address problems of privacy and fairness.
- While the authors might fear that complete honesty about limitations might be used by reviewers as grounds for rejection, a worse outcome might be that reviewers discover limitations that aren't acknowledged in the paper. Reviewers will be specifically instructed to not penalize honesty concerning limitations.

3. Theory assumptions and proofs

Question: For each theoretical result, does the paper provide the full set of assumptions and a complete (and correct) proof?

Answer: **[TODO]**

Justification: **[TODO]**

Guidelines:

- The answer NA means that the paper does not include theoretical results.
- All the theorems, formulas, and proofs in the paper should be numbered and cross-referenced.
- All assumptions should be clearly stated or referenced in the statement of any theorems.
- The proofs can either appear in the main paper or the supplemental material, but if they appear in the supplemental material, the authors are encouraged to provide a short proof sketch to provide intuition.

4. Experimental result reproducibility

Question: Does the paper fully disclose all the information needed to reproduce the main experimental results of the paper to the extent that it affects the main claims and/or conclusions of the paper (regardless of whether the code and data are provided or not)?

Answer: **[TODO]**

Justification: **[TODO]**

Guidelines:

- The answer NA means that the paper does not include experiments.
- If the paper includes experiments, a No answer to this question will not be perceived well by the reviewers: Making the paper reproducible is important.
- If the contribution is a dataset and/or model, the authors should describe the steps taken to make their results reproducible or verifiable.
- We recognize that reproducibility may be tricky in some cases, in which case authors are welcome to describe the particular way they provide for reproducibility. In the case of closed-source models, it may be that access to the model is limited in some way (e.g., to registered users), but it should be possible for other researchers to have some path to reproducing or verifying the results.

5. Open access to data and code

Question: Does the paper provide open access to the data and code, with sufficient instructions to faithfully reproduce the main experimental results, as described in supplemental material?

Answer: **[TODO]**

Justification: **[TODO]**

Guidelines:

- The answer NA means that paper does not include experiments requiring code.
- Please see the Agents4Science code and data submission guidelines on the conference website for more details.
- While we encourage the release of code and data, we understand that this might not be possible, so “No” is an acceptable answer. Papers cannot be rejected simply for not including code, unless this is central to the contribution (e.g., for a new open-source benchmark).
- The instructions should contain the exact command and environment needed to run to reproduce the results.
- At submission time, to preserve anonymity, the authors should release anonymized versions (if applicable).

6. Experimental setting/details

Question: Does the paper specify all the training and test details (e.g., data splits, hyper-parameters, how they were chosen, type of optimizer, etc.) necessary to understand the results?

Answer: [TODO]

Justification: [TODO]

Guidelines:

- The answer NA means that the paper does not include experiments.
- The experimental setting should be presented in the core of the paper to a level of detail that is necessary to appreciate the results and make sense of them.
- The full details can be provided either with the code, in appendix, or as supplemental material.

7. Experiment statistical significance

Question: Does the paper report error bars suitably and correctly defined or other appropriate information about the statistical significance of the experiments?

Answer: [TODO]

Justification: [TODO]

Guidelines:

- The answer NA means that the paper does not include experiments.
- The authors should answer "Yes" if the results are accompanied by error bars, confidence intervals, or statistical significance tests, at least for the experiments that support the main claims of the paper.
- The factors of variability that the error bars are capturing should be clearly stated (for example, train/test split, initialization, or overall run with given experimental conditions).

8. Experiments compute resources

Question: For each experiment, does the paper provide sufficient information on the computer resources (type of compute workers, memory, time of execution) needed to reproduce the experiments?

Answer: [TODO]

Justification: [TODO]

Guidelines:

- The answer NA means that the paper does not include experiments.
- The paper should indicate the type of compute workers CPU or GPU, internal cluster, or cloud provider, including relevant memory and storage.
- The paper should provide the amount of compute required for each of the individual experimental runs as well as estimate the total compute.

9. Code of ethics

331 Question: Does the research conducted in the paper conform, in every respect, with the
332 Agents4Science Code of Ethics (see conference website)?

333 Answer: **[TODO]**

334 Justification: **[TODO]**

335 Guidelines:

- 336 • The answer NA means that the authors have not reviewed the Agents4Science Code of
337 Ethics.
- 338 • If the authors answer No, they should explain the special circumstances that require a
339 deviation from the Code of Ethics.

340 10. **Broader impacts**

341 Question: Does the paper discuss both potential positive societal impacts and negative
342 societal impacts of the work performed?

343 Answer: **[TODO]**

344 Justification: **[TODO]**

345 Guidelines:

- 346 • The answer NA means that there is no societal impact of the work performed.
- 347 • If the authors answer NA or No, they should explain why their work has no societal
348 impact or why the paper does not address societal impact.
- 349 • Examples of negative societal impacts include potential malicious or unintended uses
350 (e.g., disinformation, generating fake profiles, surveillance), fairness considerations,
351 privacy considerations, and security considerations.
- 352 • If there are negative societal impacts, the authors could also discuss possible mitigation
353 strategies.