


November 7, 2013



$$\binom{6}{2} = \frac{6 \cdot 5}{2} = 15 \quad \text{degree}$$



$$\binom{3}{2} = \frac{3 \cdot 2}{2} = 3 \quad \text{external degree}$$

$\{7, 8, 9, 10\} \rightarrow$ super variable (they are a clique, can treat them as one thing).

Transition from G^2 to G^3

node 3 is selected as pivot

~~elimination graph~~
quotient graph



$$L_3 = A_3 = \{5, 6, 7\}$$

represents pairwise adjacency of variables 5, 6, 7



$(5, 7)$ is now redundant, remove from A_5, A_7

* A_i : connections b/w variables

* L_i : connections from element.

Transforming from G^4 to G^5

• variable 5 is chosen as a pivot



A_5 is empty

$$\Sigma_5 = \{2, 3\}$$

$$L_5 = (A_5 \cup L_2 \cup L_3) \setminus \{5\}$$

$$= \emptyset \cup \{5, 6, 9\} \cup \{5, 6, 7\} \setminus \{5\}$$

$$= \{6, 7, 9\}$$

• The pair (7, 9) is now redundant, so eliminate from A_7, A_9

• can also eliminate elements 2+3 b/c they don't add information

In G^4 :

$$A_6 = \emptyset$$

$$\Sigma_6 = \{2, 3, 4\}$$

$$A_7 = \{9, 10\}$$

$$\Sigma_7 = \{3, 4\}$$

$$A_9 = \{7, 8, 10\}$$

$$\Sigma_9 = \{2\}$$

In G^5 :

$$A_6 = \emptyset$$

$$\Sigma_6 = \{4, 5\}$$

$$A_7 = \{10\}$$

$$\Sigma_7 = \{4, 5\}$$

$$A_9 = \{8, 10\}$$

$$\Sigma_9 = \{5\}$$

$$L_5 = \{6, 7, 9\}$$

Approximate Minimum Degree

Approximate degree of node 6:
(in G^u)

$$L_4 = \{6, 7, 8\}$$

$$L_2 = \{5, 6, 9\}$$

$$L_3 = \{5, 6, 7\}$$

$$\bar{d}_6^4 = |A_6 \setminus \{6\}| + |L_4 \setminus \{6\}| + |L_2 \setminus L_4| + |L_3 \setminus L_4|$$

allow common guys btwn L_2 & L_3

$$= \emptyset \setminus \{6\} + |\{6, 7, 8\} \setminus \{6\}| + |\{5, 6, 9\} \setminus \{6, 7, 8\}| + |\{5, 6, 7\} \setminus \{6, 7, 8\}|$$
$$= \quad \quad \quad 2 \quad \quad + \quad \quad 2 \quad \quad + \quad 1$$

$$= 5$$

The actual degree of 6 is

$$d_6^4 = 4$$

Eigenvalues:

November 12, 2013

$$Ax = b \quad \text{love}$$

$$Ax = \lambda x \quad \text{peace}$$

for linear systems

$$Ax = b$$

↳ direct - compute in finite number of operations

↳ iterative - could be infinite

now → infinite

$$Ax = \lambda x$$

↳ direct

↳ iterative

($Ax = \lambda x$: right eigenvector, $y^* A = \lambda y^*$: left eigenvector)

all of them are iterative, but direct are decompositional (QR iteration)

iterative are mat-vec product based

(Lanczos/Ronaldi, Jacobi/Davidson)

$$AX = X\Lambda$$

where

X : matrix w/ eigenvectors in its columns

Λ : diagonal

$$A = \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix}$$

$$A = X\Lambda X^{-1}$$

- diagonalizable

$$A = Q\Lambda Q^* \quad (AA^* = A^*A, \text{ normal} - \text{symm. included})$$

- unitarily diagonalizable

non-diagonalizable

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$A = \begin{pmatrix} 0 & 1 & 0 \\ & \ddots & \\ 0 & & 0 \end{pmatrix} \quad n=10, \text{ e.g.}$$

eigenvectors? → 1 eigenvector of geometric multiplicity 1

$$x = e_1$$

$\lambda = 0$ is an eigenvalue w/ algebraic multiplicity n

$$\det(\lambda I - A) = 0 = \lambda^n$$

but now, if

$$A_\varepsilon = \begin{pmatrix} 0 & 1 & & 0 \\ & 0 & & \\ & & \ddots & \\ \varepsilon & & & 0 \end{pmatrix}$$

$$\det(\lambda I - A_\varepsilon) = 0$$


$$\lambda^n - \varepsilon = 0$$

$$|\lambda| = \sqrt[n]{\varepsilon}$$

$$\text{if } \varepsilon = 10^{-10}$$

$$|\lambda| = 0.1$$

Small perturbation in the matrix \rightarrow huge perturbation in the solution



$$Ax = \lambda x, \quad y^* A = \lambda y^*$$

$$(A + \delta A)(x + \delta x) = (\lambda + \delta \lambda)(x + \delta x) \quad \text{where } \|\delta A\| \approx \varepsilon \|A\|$$

$$Ax + \delta Ax + A\delta x + \delta A\delta x = \lambda x + \delta \lambda x + \lambda \delta x + \delta \lambda \delta x$$

assume we can ignore second order terms
estimate $\delta \lambda$ and

$$\delta Ax + A\delta x = \delta \lambda x + \lambda \delta x$$

mult. by y^*

$$y^* \delta Ax + y^* A \delta x = y^* \delta \lambda x + y^* \lambda \delta x$$

$$\boxed{\delta \lambda = \frac{y^* \delta Ax}{y^* x}}$$

If y^*, x orthogonal \rightarrow blows up.

$\frac{1}{y^*x}$: condition number of the eigenvalue problem

If A is symmetric

$$\delta\lambda \sim \epsilon \|A\|$$

since

$$\|\delta\lambda\| \leq \frac{\|y\| \|\delta A\| \|x\|}{\|y^*x\|} \leq \epsilon \|A\|$$

Forward + Backward error

ex:

$$\sqrt{2} \rightarrow 1.4 \rightarrow 1.4 = \sqrt{1.96}$$

$$(x^2 - 2 = 0: \text{Newton's})$$

$$\text{Forward error} = |1.4 - 1.414...|$$

$$\text{Backward error} = |2 - 1.96|$$

Power method:

• orthogonal / simultaneous iteration (QR +)

• Lanczos / Arnoldi

• gives us the dominant eigenpair

$$Ax_i = \lambda_i x_i \quad |\lambda_1| > |\lambda_2| \geq |\lambda_3| \geq \dots \geq |\lambda_n|$$

V : initial guess

$$V = \sum_{i=1}^n \alpha_i x_i$$

$$\begin{aligned} A^k V &= \sum_{i=1}^n \alpha_i \underbrace{A^k x_i}_{\lambda_i^k x_i} = \alpha_1 \lambda_1^k x_1 + \sum_{i=2}^n \alpha_i (\lambda_i^k x_i) \\ &= \lambda_1^k \left(\alpha_1 x_1 + \sum_{i=2}^n \alpha_i \left(\frac{\lambda_i}{\lambda_1} \right)^k x_i \right) \end{aligned}$$

initial $V_0 \quad i=1, \dots$

repeat

$$V_i = A V_{i-1}$$

$$V_i \leftarrow V_i / \|V_i\|$$

$\downarrow 0$
 $k \rightarrow \infty$

eventually

$$v_i \rightarrow x_i$$

eigenvalue

$$r(v_i) = \frac{v_i^T A v_i}{v_i^T v_i}$$

Rayleigh Quotient

best estimate for an eigenvalue in the L-S sense: given x

$$\min_x \|Ax - \lambda x\|_2$$

$$= \min_x \|C - \lambda x\|_2$$

where $C = Ax$ given

$$(\text{recall } \min \|b - Ax\| \text{ NE } A^T A x = A^T b)$$

so the normal equations are

$$x^T C = x^T x \lambda$$

$$\Rightarrow \lambda = \frac{x^T C}{x^T x} = \frac{x^T A x}{x^T x}$$

Terribly slow method:

↳ converges linearly - proportional to λ_2/λ_1

↳ if initial guess has no component in the direction of x_1 ,
but → round-off errors save the day!

Shift & invert (inverse power method)

$$(A - \alpha I)^{-1}$$

$$A: \lambda_1, \dots, \lambda_n$$

eigenvalues of ↗

$$\frac{1}{\lambda_1 - \alpha}, \frac{1}{\lambda_2 - \alpha}, \dots, \frac{1}{\lambda_n - \alpha}$$

apply power method on $(A - \alpha I)^{-1}$

say $\alpha \approx \lambda_2$

$$\frac{1}{\lambda_2 - \alpha} \gg \frac{1}{\lambda_i - \alpha} \quad i \neq 2$$

• need to solve a system
• need a good way to guess α